

**Potential and Limitations of *Klebsiella pneumoniae* as a Microbial Cell Factory Utilizing  
Glycerol as the Carbon Source**

**Vinod Kumar<sup>a</sup>, Sunghoon Park<sup>b\*</sup>**

**<sup>a</sup> Bioenergy and Resource Management Centre, School of Water, Energy and  
Environment, Cranfield University, Cranfield MK43 0AL, United Kingdom**

**<sup>b</sup> School of Energy and Chemical Engineering, UNIST, 50 UNIST-gil, Ulsan 44919,  
Republic of Korea**

**\*Corresponding author**

**School of Energy and Chemical Engineering, UNIST, 50 UNIST-gil, Ulsan 44919,  
Republic of Korea**

**Tel.: +82 52 217 2565; Fax: +82 52 217 2309; E-mail address: parksh@unist.ac.kr**

## Abstract

*Klebsiella pneumoniae* is a Gram-negative facultative anaerobe that metabolizes glycerol efficiently under both aerobic and anaerobic conditions. This microbe is considered an outstanding biocatalyst for transforming glycerol into a variety of value-added products. Crude glycerol is a cheap carbon source and can be converted by *K. pneumoniae* into useful compounds such as lactic acid, 3-hydroxypropionic acid, ethanol, 1,3-propanediol, 2,3-butanediol, and succinic acid. This review summarizes glycerol metabolism in *K. pneumoniae* and its potential as a microbial cell factory for the production of commercially important acids and alcohols. Although many challenges remain, *K. pneumoniae* is a promising workhorse when glycerol is used as the carbon source.

**Key words:** *Klebsiella pneumoniae*; Glycerol; 3-Hydroxypropionic acid; 1,3-Propanediol; 2,3-Butanediol; Lactic acid; Ethanol

## 1. Introduction

Crude glycerol is produced in large quantities as a byproduct of the biodiesel industry. Global biodiesel production was 22.7 million tons in 2012 and is forecasted to reach 36.9 million tons by 2020. As a result, worldwide production of glycerol climbed from 1 million tons in 2000 to 3 million tons in 2011 and is expected to reach 6 million tons by 2025 (Ciriminna et al., 2014; OECD, 2011; Katryniok et al., 2013; Sun et al. 2017). A significant amount of glycerol is also generated during bioethanol fermentation by yeast. The liquid fraction of the stillage obtained from ethanol separation contains up to 2% glycerol. Another source of glycerol is industrial waste generated from vegetable oils and animal fats. For example, the concentration of glycerol in waste streams from the oleochemical industry is 55%–90% (da Silva et al., 2009; Yazdani and Gonzalez, 2007). The increased availability of glycerol has caused a substantial reduction in its cost. According to a recent report, the price of crude glycerol is \$0.24/kg and that of pure USP grade glycerol is \$0.9/kg (Kong et al., 2016). Extensive research has been conducted to investigate ways to utilize this surplus crude glycerol. The annual number of research articles addressing the use of glycerol has increased to >7000, doubling in number from the year 2000 to 2007. Numerous chemical conversions to valuable products have been reported (Ciriminna et al., 2014).

Glycerol can be used for the microbial production of 1,3-propanediol (PDO), ethanol, 2,3-butanediol (BDO), butanol, lactic acid, 3-hydroxypropionic acid (3-HP), and succinic acid (Fig. 1) (Clomberg and Gonzalez, 2013; Mattam et al., 2013; Yazdani and Gonzalez, 2007). Because of its reduced nature, glycerol generates twice the number of reducing equivalents per carbon than do traditional carbohydrates (glucose, xylose, and sucrose) when converted into the glycolytic intermediates phosphoenolpyruvate (PEP) or pyruvate (Dharmadi et al., 2006). As a consequence, glycerol gives higher yields of reduced metabolites (e.g., succinate, ethanol, PDO, and butanol) than does glucose (Yazdani and

Gonzalez, 2007). Table 1 compares the maximum theoretical yield of various metabolites, ATP, and redox balances between glycerol and glucose.

Despite its great potential as a carbon source, glycerol is not efficiently metabolized by many microorganisms under anaerobic or oxygen-limited conditions. The ability of *K. pneumoniae* to assimilate glycerol under oxygen-limited conditions is outstanding and has been studied extensively. Several review articles addressing this issue have also been published (Clomberg and Gonzalez, 2013; Mattam et al., 2013; Yazdani and Gonzalez, 2007), but the biotechnological potential and physiological aspects of this strain have not been fully elucidated. This review explores and discusses glycerol metabolism in *K. pneumoniae* and the production of value-added chemicals from glycerol. The potential and challenges of using *K. pneumoniae* as a microbial cell factory are also discussed.

## **2. Glycerol metabolism in *K. pneumoniae***

Many microorganisms can metabolize glycerol in the presence of external electron acceptors (respiratory metabolism), but relatively few under non-respiratory conditions (without exogenous electron acceptors). Among the latter, genera including *Citrobacter*, *Enterobacter*, *Bacillus*, *Propionibacterium*, *Anaerobiospirillum*, *Klebsiella*, *Clostridium*, and *Lactobacillus* are well documented (da Silva et al., 2009; Yazdani and Gonzalez, 2007). Some *Lactobacillus* species including *L. reuteri*, *L. brevis*, and *L. buchneri* convert glycerol to PDO but only in the presence of a major carbon source for growth. Glycerol does not support cell growth, limiting its biotechnological applications. In contrast, *Clostridium* sp. grow well on glycerol as the sole carbon source. However, the use of *Clostridium* is limited by its slow growth and requirement of strict anaerobic conditions. The lack of a convenient genetic tool box for large-scale metabolic engineering is another serious drawback of *Clostridium* (Branduardi et al., 2014; Lütke-Eversloh and Bahl, 2011).

From a biotechnological standpoint, *Klebsiella* sp., especially *K. pneumoniae*, have many advantages over the aforementioned microbes. They grow on glycerol as the sole carbon source under both aerobic and anaerobic conditions (Arasu et al., 2011; Kumar et al., 2012). Their growth is fast, typically growing at a rate >0.9/h, even on glycerol minimal medium (Arasu et al., 2011). They naturally produce coenzyme B<sub>12</sub>, an essential cofactor for the synthesis of the commercially important chemicals PDO and 3-HP (Huang et al., 2013a; Oh et al., 2012a). Furthermore, their evolutionary and biochemical proximity to *E. coli* allows for the application of most of gene manipulation methods developed for *E. coli* without much modification (Celińska, 2012; Kumar et al., 2013a).

## **2.1 Overview of glycerol metabolism in *K. pneumoniae***

In *K. pneumoniae*, glycerol is metabolized oxidatively and/or reductively (Ashok et al., 2011, Kumar et al., 2013a). The oxidative pathway provides energy and carbon constituents for the synthesis of cell biomass, while the reductive pathway reduces glycerol and regenerates NAD<sup>+</sup>, enabling the oxidative assimilation of glycerol under anaerobic conditions.

Glycerol oxidation proceeds in either a respiratory or a fermentative manner. Respiration requires an exogenous electron acceptor such as oxygen, nitrate, or fumarate (Ashok et al., 2013a,b) and is carried out by enzymes encoded by the *glp* regulon. During respiration, glycerol is first phosphorylated by glycerol kinase (*glpK*) to yield *sn*-glycerol-3-phosphate using ATP or phosphoenolpyruvate (PEP) as a phosphate donor. Next, *sn*-glycerol-3-phosphate is oxidized to dihydroxyacetone phosphate (DHAP) with electron transfer to a quinone molecule, which is linked to the electron transport chain (ETC). This transformation is catalyzed by glycerol-3-phosphate dehydrogenase, which is expressed under both aerobic (*glpD*) and anaerobic (*glpABC*) conditions. In contrast, fermentation of glycerol is conducted without a supply of external electron acceptors, and the reactions are

catalyzed by enzymes encoded by the *dha* regulon. Here, glycerol is oxidized to dihydroxyacetone (DHA) by glycerol dehydrogenase (mainly encoded by *dhaD*) using NAD<sup>+</sup> as an electron acceptor. DHA then is phosphorylated by dihydroxyacetone kinase, an enzyme encoded by the ATP-dependent *dhaK1/dhaK* and/or PEP-dependent *dhaK2/dhaKLM/dhaK123*. DHAP obtained from respiration and/or fermentation is channeled into the glycolytic pathway (Fig. 2). A variety of organic acids and alcohols, including acetic acid, lactic acid, succinic acid, ethanol, BDO, and formic acid, are produced by the downstream glycolytic pathway (Ashok et al., 2011; Kumar et al., 2013a).

The anaerobic growth capability of *K. pneumoniae* using glycerol as a carbon source is attributed to its well-established reductive metabolism. In the reductive pathway, glycerol first undergoes a difficult rearrangement reaction catalyzed by glycerol dehydratase (GDHt) to yield 3-hydroxypropionaldehyde (3-HPA). Catalysis by GDHt requires coenzyme B<sub>12</sub>, the *de novo* synthesis of which is limited to only few genera of microorganisms. The 3-HPA is subsequently reduced to PDO by at least two 1,3-propanediol oxidoreductases (1,3-PDORs), including NADH-dependent DhaT (PDOR) and NADPH-dependent hypothetical oxidoreductase (HOR), along with the regeneration of NAD(P)<sup>+</sup> (Fig. 3). *E. coli* and other *enterobacter sp.* have the oxidative (respiratory) pathway of glycerol assimilation. However, these organisms lack the reductive pathway and cannot grow on glycerol under anaerobic conditions (Dharmadi et al., 2006; Kumar et al., 2013a).

## **2.2 Genes and enzymes involved in fermentative metabolism of glycerol**

The *dha* regulon, induced by the DHA molecule and expressed in the absence of exogenous electron acceptors, encodes numerous genes needed for both the oxidative (fermentative) and reductive pathways (Celińska, 2012; Forage and Foster, 1982) (Fig. 4A). The *dha* regulon of *K. pneumoniae* contains the following genes arranged in the order as appears: *Kpk\_0615/dhaK*, *dhaM*, *dhaL*, *dhaK*, *dhaD*, *dhaR*, *orfW*, *CdAT*, *orfX (dhaG)*, *dhaT*,

*orfY*, *dhaB*, *dhaC*, *dhaE*, *orfZ* (*dhaF*), and *glpF*. Of these, *dhaB*, *dhaC*, *dhaE*, *orfX*, *orfZ*, and *dhaT* are relatively well documented for their functions in the reductive metabolism of glycerol. The *orfW* and *orfY* genes, although commonly found in the *dha* regulons of many organisms, are not as well described. The proteins encoded by the *dha* regulon of *K. pneumoniae* have high similarity (80%–95%) with those of *Citrobacter* species but not with those of *Clostridium* species (30%–80%). Surprisingly, the sequence similarity of *dha* proteins among *Clostridium* species is very low in some cases, even lower than their similarity to the corresponding proteins in *K. pneumoniae* or *C. freundii* (Celińska, 2012; Sun et al., 2003; Wei et al., 2014).

The complete *dha* regulon is found in only a few dozen species belonging to five different taxonomic groups, including four bacterial species [Actinobacteria, Firmicutes, Fusobacteria, and Proteobacteria (Gammaproteobacteria and Deltaproteobacteria)] and one archaea species, *Halalkalicoccus jeotgali* (Martins-Pinheiro et al., 2016). Interestingly, an incomplete *dha* regulon is present in more than 100 prokaryotes, suggesting that the enzymes of the regulon have other functions not restricted to the anaerobic growth on glycerol. A well-documented example is the one found in *Lactobacillus* species. *Lactobacillus* cannot grow on glycerol as the sole carbon source because it lacks the enzymes for oxidative assimilation of glycerol. However, the enzymes encoded in the incomplete *dha* regulon enable the production of PDO from glycerol, which accelerates cell growth in *Lactobacillus* grown on glucose (Årsköld et al., 2008; Martins-Pinheiro et al., 2016).

In the following sections, the major enzymes involved in fermentative glycerol metabolism and their genes are described in detail.

### **2.2.1 Glycerol dehydrogenase**

Encoded by the *dhaD* and/or *glcA* gene, glycerol dehydrogenase is an oxidoreductase that converts glycerol to DHA. The enzyme extracts electrons from glycerol and transfers

them to NAD<sup>+</sup>. DhaD plays a major role during the anaerobic growth of *K. pneumoniae* on glycerol. Quantification of mRNA expression indicates that *dhaD* is highly induced when glycerol is used as carbon source. GldA expression is lower than that of DhaD and is thought to serve as a back up to DhaD during anaerobic growth on glycerol. However, DhaD and GldA exhibit a high level of similarity in amino acid sequence and have the same substrate-binding site (Wang et al., 2014).

Recently, DhaD was reported to catalyze the interconversion of acetoin and BDO for NADH disposal and recovery (Wang et al., 2014). The promiscuity of DhaD is responsible for BDO production in the  $\Delta budC$  mutant of *K. pneumoniae*. The contribution by DhaD to or its participation in BDO production is also supported by the observed increase in BDO synthesis and increased concentration of NADH in the presence of the *dhaD* gene. NADH is generated by the oxidation of glycerol and consumed by the production of BDO from acetoin. The dual physiological functions of DhaD allow *K. pneumoniae* to control the intracellular redox level. In response to the intracellular level of NADH, DhaD is able to switch its role from NADH production (glycerol oxidation) to NADH consumption (BDO production) (Wang et al., 2014). The diversion of glycerol flux from organic acids toward alcohols such as BDO also helps the cells to counteract intracellular acidification (Petrov and Petrova, 2009). Thus, this promiscuity confers different physiological roles to the enzyme, which include regulation of the intracellular NADH/NAD<sup>+</sup> ratio, prevention of acidification, and storage of carbon and energy (Wang et al., 2014).

### **2.2.2 Dihydroxyacetone kinases**

Dihydroxyacetone kinase (DHAK) phosphorylates DHA to DHAP, an intermediate of the glycolytic pathway. Sun et al. (2003) analyzed the genome of *K. pneumoniae* MGH 78578 and found, in addition to ATP-dependent DhaK I, a PEP-dependent DhaK II. DhaK I is encoded by the *dhaK* (*Kpk\_0615*) gene, while DhaK II is encoded by three genes, *dhaK*,



*dhaL* and *dhaM*, which correspond to the medium, small, and large subunits of the enzyme, respectively (Celińska, 2012; Sun et al., 2003; Wei et al., 2014). The names *dhaKLM* and *dhaK123* are synonyms used without distinction (Martins-Pinheiro et al., 2016; Sun et al., 2003; Wei et al., 2014). DhaK enzymes are found in many microorganisms. PEP-dependent DhaK II is present in *E. coli*, *L. lactis*, and *C. butyricum*, whereas ATP-dependent DhaK I is present in *Citrobacter freundii* (Garcia-Alles et al., 2004; Raynaud et al., 2011). The PEP-dependent DhaK II of *E. coli*, encoded by *dhaKLM*, is homologous to *dhaK123* (*dhaKLM*) of *K. pneumoniae* (Gutknecht et al., 2001). In a recent study, Wei et al. (2014) further elucidated the physiological roles of DhaK I and DhaK II of the *dha* regulon of *K. pneumoniae*. They found that disruption of ATP-dependent DhaK I had no significant effect on glycerol uptake, indicating that PEP-dependent DhaK II is the major contributor to the conversion of DHA to DHAP. As in *E. coli*, the subunits of DhaK II regulate expression of the *dha* regulon; *dha* regulon expression was suppressed by the disruption of *dhaK1* (*dhaK*) and *dhaK2* (*dhaL*). However, in contrast to *E. coli*, mutation of *dhaK3* (*dhaM*) in *K. pneumonia* did not upregulate the *dha* regulon but rather decreased cell growth and glycerol uptake.

### 2.2.3 Glycerol dehydratase and reactivating factor

Glycerol dehydratase (GDHt) is the first enzyme in the reductive pathway and catalyzes the dehydration of glycerol to 3-HPA. The same reaction can be carried out by diol dehydratase (DDHt), although the latter has a higher activity on 1,2-propanediol than on glycerol. GDHt requires coenzyme B<sub>12</sub> or S-adenosyl methionine (SAM) as a cofactor. *K. pneumoniae* has B<sub>12</sub>-dependent GDHt only, while strict anaerobes such as *clostridia* sp. have SAM-dependent GDHt. The three subunits of GDHt are encoded by the genes *dhaB1* (~1670 bp), *dhaB2* (~590 bp) and *dhaB3* (~430 bp), for which the gene products are the  $\alpha$  (large, 60.7 kDa),  $\beta$  (medium, 21.3 kDa), and  $\gamma$  (small, 16.1 kDa) subunits, respectively (Xu et al., 2009a). GDHt of *K. pneumoniae* is known to be present as a dimer of heterotrimers, ( $\alpha\beta\gamma$ )<sub>2</sub>.

The subunits of GDHt have high homology among *K. pneumoniae* strains; for example, *K. pneumoniae* XJPD-Li and U30909 strains showed >99% identity.

Coenzyme B<sub>12</sub> is present in the reaction center and plays an essential role in the catalysis of GDHt. The catalytic cycle often results in the mechanism-based inactivation of coenzyme B<sub>12</sub> due to the irreversible breakage of the chemical bond between cobalt (Co) and a carbon of the adenosyl moiety. Oxygen accelerates breakage of the Co–C bond. Oxygen is also known to inactivate GDHt, even in the absence of the substrate glycerol. The gene products of *orfZ/dhaG* (small subunit) and *orfX/dhaF* (large subunit) act as reactivating factors of GDHt (GdrAB) (Celińska, 2012; Sun et al., 2003). They reactivate GDHt by catalyzing the exchange of damaged for intact coenzyme B<sub>12</sub> in the presence of ATP and Mg<sup>2+</sup>/Mn<sup>2+</sup> (Celińska, 2012; Kumar et al., 2013a; Maervoet et al., 2011; Sun et al., 2003; Wei et al., 2014). The reactivase is a molecular chaperone that functions as a heterotetramer containing two elongated  $\alpha$  subunits (63 kDa) and two globular  $\beta$  subunits (14 kDa). Structurally, the  $\alpha$  subunit resembles both GroEL and Hsp70 chaperones, while the  $\beta$  subunit resembles that of the  $\beta$  subunit of glycerol dehydratase, except that it lacks some of the amino acids responsible for coenzyme B<sub>12</sub> binding (Liao et al., 2003).

In some organisms, the GDHt enzyme varies in terms of the number of subunits and their copies (Fig. 4B) (Liu et al., 2010). For example, in *Mesorhizobium loti* and *Mesorhizobium opportunistum*, the large and medium subunits (*dhaB1* and *dhaB2*) are fused together and encoded by a single gene. The gene fusion may arise by frameshift mutations. The active site of the fused enzyme differs slightly from that of unfused enzyme. Moreover, these microorganisms lack the genes encoding the reactivation factor, indicating that the reactivation process may not exist or is carried out by a different route. *Mycobacterium smegmatis* has two subunits (fused *dhaB1* + *dhaB2* and *dhaB3*) that are similar to those of *M. loti* and *M. opportunistum*; however, the reactivating factor differs in that it has one (large)

subunit. No small subunit has been identified in *M. smegmatis*. *Fusobacterium Ilyobacter polytropus* is unique in that it has genes coding for both B<sub>12</sub>-independent (SAM-dependent) and B<sub>12</sub>-dependent GDHt. The genes for B<sub>12</sub>-independent GDHt are present in the genomic DNA, while those encoding B<sub>12</sub>-dependent GDHt along with its reactivation factor are present in two copies, one in the chromosome and the other in a plasmid. Surprisingly, even with this gene redundancy, no significant production of 3-HPA and/or PDO has been reported in *I. polytropus* (Stieb and Schink, 1984). The B<sub>12</sub>-independent proteins showed no homology with B<sub>12</sub>-dependent GDHt but significant similarity with the pyruvate formate lyases (PFL) and PFL-activating enzymes and their homologues (Raynaud et al., 2003).

#### **2.2.4 1,3-Propanediol oxidoreductase and hypothetical oxidoreductase**

The 1,3-propanediol oxidoreductase (PDOR), responsible for the second step of the reductive pathway of glycerol, catalyzes the conversion of 3-HPA to PDO with the oxidation of NAD(P)H. 3-HPA is a toxic intermediate, and PDOR protects the cell from deleterious damage by reducing 3-HPA to PDO (Celińska, 2012). In *K. pneumoniae*, PDOR is encoded by the *dhaT* gene and is strictly NADH dependent. Active PDOR is a homo-octamer with a monomeric molecular mass of 43 kDa. Despite its importance, the kinetic properties of DhaT were elucidated only recently (Lama et al., 2015). DhaT has high specificity for 3-HPA among various aldehydes. Under physiological conditions at pH 7.0 and 37°C, the catalytic efficiency ( $k_{cat}/K_m$ ) of the forward reaction (3-HPA to PDO) was found to be 77.2/s. mM, which is 57-fold higher than that of the reverse reaction (1.3/s. mM; PDO to 3-HPA). The  $K_m$  (mM) values for 3-HPA and NADH (forward reaction) are 0.77 and 0.03, respectively, while those for PDO and NAD<sup>+</sup> (backward reaction) are 7.4 and 0.23, respectively. Furthermore, the maximum activity of DhaT for the reverse reaction was obtained at pH 9.0 and 55°C. These results strongly suggest that, under physiological conditions, DhaT catalyzes the reduction of 3-HPA to PDO rather than the oxidation of PDO to 3-HPA. Nevertheless, the

reverse reaction (PDO→3-HPA) becomes significant as the PDO concentration increases. This factor is considered a serious drawback of using DhaT for PDO production from glycerol.

*K. pneumoniae* has several oxidoreductases in addition to PDOR that can reduce 3-HPA to PDO. One such enzyme, designated as hypothetical oxidoreductase (HOR), has been studied extensively and shares 89% identity with YqhD of *E. coli* (Zhu et al., 2009; Zhuge et al., 2010). Expression of *yqhD* in wild-type *K. pneumoniae* is not high, with an mRNA level ~20-fold lower than that of *dhaT*. However, deletion of *dhaT* increases the transcription of *yqhD* by more than 10-fold (Ko et al., 2015). This observation suggests that in addition to its wide-ranging activity on short-chain aldehydes, YqhD is a detoxification enzyme for many aldehydes (Jarboe, 2011). YqhD exhibits negligible activity on PDO (backward reaction), which is an important advantage over DhaT for use in the production of PDO from glycerol (Celińska, 2012; Chen et al., 2011; Zheng et al., 2006). However, YqhD is kinetically inferior to DhaT because of its low catalytic activity ( $k_{cat}/K_m$ ) on 3-HPA and the requirement of NADPH as a cofactor. The  $k_{cat}/K_m$  value of YqhD on 3-HPA (pH 7.0 and 37°C) is estimated to be 2.1/s. mM, which is 36-fold lower than that of DhaT (unpublished data). The use of NADPH as a cofactor prohibits this enzyme from being directly coupled to the reactions catalyzed by DhaD and other enzymes in glycolysis and the TCA cycle, by which NADH is mainly produced. Furthermore, excessive use of NADPH for PDO production can significantly reduce the cellular NADPH level and this can disturb many anabolic reactions requiring NADPH (Wang et al., 2003; Jarboe, 2011). NADH and NADPH differ in a phosphate group only, and it has been suggested that electrostatic interaction of this phosphate is the major factor distinguishing NADH from NADPH for enzymes using these cofactors. Comparison of the amino acid sequences of DhaT and HOR from *K. pneumoniae* and YqhD from *E. coli* has revealed that DhaT contains aspartic acid (Asp) at position 41,

while HOR and YqhD have valine (Val) at this position. The steric hindrance and electrostatic repulsion between Asp in the active site and the phosphate group of NADPH impede their binding. In contrast, the hydrophobic amino acid Val poses no such obstruction; thus, YqhD uses NADPH as a cofactor (Ma et al., 2010).

Despite their difference in cofactor preference, DhaT and YqhD are similar (44%) in amino acid sequences. *K. pneumoniae* has five enzymes that have at least 40% similarity in amino acid sequence with DhaT and YqhD: putative alcohol dehydrogenase (PaldH), L-1,2-propanediol oxidoreductase (FucO), ethanolamine-utilization enzyme (EutG), propanediol-utilization propanol dehydrogenase (PduQ), and bifunctional alcohol dehydrogenase (AdhE) (Ko et al., 2015). The roles of these enzymes in PDO production have been investigated.

### 2.2.5 Regulatory protein

The *dhaR* gene product is an important transcription factor that is responsible for expression of the *dhaT* and *dhaB* genes (Sun et al., 2003). This gene is found in *K. pneumoniae*, *C. freundii*, and *Clostridium botulinum* but not in every organism containing a complete *dha* regulon (Martins-Pinheiro et al., 2016). The presence of DhaR is thought to allow the *dha* operon to function as a separate regulatory system, independent of fumarate nitrate reduction (FNR). Under anaerobic conditions, FNR acquires DNA-binding properties and induces the expression of many anaerobiosis-related genes. DhaR contains domains for GAF (52-199), PAS (203-267),  $\sigma^{54}$  factor interaction, and histidine HTH-8. The PAS senses internal levels of energy charge, light, oxygen, and redox potential. The HTH-8 domain has a helix-turn-helix conformation and acts as a DNA binding structure. The  $\sigma^{54}$  factor interaction domain interacts with the  $\sigma^{54}$  factor of RNA polymerase and activates RNA transcription from the  $\sigma^{54}$  promoters (Buck et al., 2000; Taylor and Zhulin, 1999). Thus, the DhaR protein triggers the transcription of *dha* genes in response to intracellular levels of several important physiological parameters (Celińska, 2012; Sun et al., 2003). Zheng et al. (2006) investigated

the effect of overexpression of the putative regulatory gene *dhaR* in the *dha* regulon on glycerol metabolism in *K. pneumoniae*. *dhaR* overexpression increased PDOR activity up to 6.7-fold over that of the control wild-type strain, confirming the role of DhaR as a positive regulator of the *dhaT* gene. However, *dhaR* overexpression did not increase the activity of glycerol dehydratase (DhaB). Furthermore, despite the significant increase in PDOR activity, PDO production was unexpectedly low compared with that of the wild type (440 vs. 656.23 mM, respectively). More studies are required to determine the precise role of DhaR and its regulatory function in the *dha* regulon.

DhaR in *E. coli* (70% identity to that of *K. pneumoniae*) has also been studied. This protein stimulated transcription of the *dhaKLM* operon from a  $\sigma^{70}$  promoter. Interestingly, in *E. coli*, DhaL was a positive regulator of the *dhaKLM* operon while phosphorylated DhaM and DhaK were negative regulators of the same operon. DhaK contains the DHA-binding site. DhaL uses ADP as a cofactor for the double displacement of phosphate from DhaM to DHA. DHA is the oxidation product of glycerol metabolism; binding of DHA to DhaK reduces the affinity of DhaK for DhaR. In the presence of DHA, DhaL::ATP is dephosphorylated, leading to the displacement of DhaK by DhaL::ADP, which stimulates DhaR activity. In the absence of DHA, DhaL::ADP is phosphorylated back to DhaL::ATP by the phototransferase system (PTS), thereby inhibiting the binding of the phosphorylated complex (DhaL::ATP) to DhaR. This phosphorylation is mediated by DhaM, providing a phospho-histidine relay between the PTS and DhaL::ADP. This double-check mechanism of binding and turnover increases the selectivity such that the binding of nonphosphorylated compounds does not induce the *dha* operon (Bächler et al., 2005).

### **2.3 Genes and enzymes involved in respiratory assimilation of glycerol**

The *glp* regulon comprises the genes encoding proteins required for the conversion of glycerol to DHAP via sn-glycerol-3-phosphate. The five operons that constitute the *glp*

regulon are *glpFK*, *glpD* (anaerobic glycerol-3-phosphate dehydrogenase), *glp ACB* (aerobic glycerol-3-phosphate dehydrogenase), *glpTQ* (glycerol-phosphate permease/glycerophosphodiesterase), and *glpEGR*. The *glpF* gene encodes a cytoplasmic membrane protein that facilitates the diffusion of glycerol into the cell. The *glpE* gene encodes an acidic, cytoplasmic protein of 108 amino acids with a molecular weight of 12,082. The *glpG* gene encodes a basic, cytoplasmic membrane-associated protein of 276 amino acids with a molecular weight of 31,278 (Lin, 1976; Yang and Larson, 1998; Zeng et al., 1996). The biochemical functions of GlpE and GlpG are unknown. The *glpR* gene encodes the GlpR repressor protein. These five operons are located at three different positions on the chromosome. Transcription of these operons is subject to multiple controls, including catabolite repression mediated by cAMP-CRP and respiratory control mediated by the FNR and ArcA/ArcB systems. Moreover, each of the operons is negatively controlled by a repressor specific for the regulon (Lin, 1987; Zeng et al., 1996).

### **2.3.1 GlpR and other regulatory proteins**

The GlpR repressor protein, predicted to contain 252 amino acids with a molecular weight of 28,048, belongs to the DeoR bacterial regulatory factor super family. Although not studied extensively, GlpR of *K. pneumonia* is expected to have the same biochemical properties and regulatory functions as that of *E. coli*. GlpR negatively controls all the *glp* operons by binding to operators that overlap or are close to the *glp* promoters. In case of *E. coli*, glycerol-3-phosphate acts as an inducer and has a high affinity for GlpR ( $K_d = 31 \mu\text{M}$ ) (Lin, 1987; Weissenborn et al., 1992; Zeng et al., 1996). When complexed with glycerol-3-phosphate, GlpR cannot bind to the operator sites, and repression is relieved. Glycerol-3-phosphate is a product of GlpK and a substrate of GlpD/GlpABC; thus, the absence of GlpD causes a substantial increase in the level of glycerol-3-phosphate, resulting in induction of the *glp* regulon (Fig. 2). The GlpR protein exhibits differential binding affinity for the operators

of corresponding operons, as follows: *glpFK* > *glpD* > *glpTQ* > *glpACB*. The *glpFK* promoter appears to be the strongest among the *glp* promoters. Therefore, the maximum strength of the *glpFK* promoter combined with the high affinity of GlpK for glycerol ( $K_m = 10 \mu\text{M}$ ) should result in sufficient accumulation of the inducer, glycerol-3-phosphate, in the presence of glycerol to relieve repression by GlpR. The relatively lower strength of the *glpD* promoter along with the low affinity of the GlpD enzyme for glycerol-3-phosphate ( $K_m = 1 \text{ mM}$ ) prevents excessive degradation of glycerol-3-phosphate that is needed for phospholipid synthesis when the exogenous source of glycerol-3-phosphate becomes limiting, until after accumulation of the inducer (Lin, 1987; Weissenborn et al., 1992). Recently, Jung et al. (2014) investigated the role of GlpR in the aerobic production of 3-HP from glycerol in *E. coli*. They found that elimination of this regulatory factor caused increased assimilation of glycerol and higher production of 3-HP and suggested that the deletion of *glpR* led to the upregulation of genes involved in glycerol transport and assimilation. *K. pneumoniae* has several other regulatory genes in addition to *glpR*: the global regulatory genes *hdeB*, *hdeD*, and *yfdX*, with unknown functions and *kvgS* and *kvgA*, encoding two proteins comprising a two-component signal transduction system (Sun et al., 2003).

### **2.3.2 Glycerol uptake/transport facilitator and other components**

The glycerol facilitator GlpF is an integral membrane protein that forms aqueous pores. The GlpF pores selectively allow passive transport of glycerol and other molecules such as DHA and urea across the cytoplasmic membrane, but not water or ions (Heller et al., 1980; Reizer et al., 1993). The GlpF protein contains six membrane-spanning helices that are unique among prokaryotic transport proteins. These transmembrane helices are arranged in two bundles. The GlpF protein may function as a homodimer with the two six-member domains arranged in the membrane to form a channel (Weissenborn et al., 1992). *K. pneumoniae* has two glycerol transporter genes, one next to *orfZ/dhaF* in the fermentative



*dha* regulon (see Fig. 4A) and the other in the respiratory *glpFK* operon. Both *glpF* genes have a high similarity (85 %) to each other. When one but not both of the *glpF* genes is disrupted, the *glpF* deletion mutants of *K. pneumoniae* can consume glycerol and convert it to PDO at a high rate, similar to the wild type (unpublished data; Supplementary Fig. S1). This observation indicates that both *glpF* genes encode active glycerol transporter proteins. However, their presence is not essential for glycerol utilization in *K. pneumoniae* because the double-deletion mutant for both facilitators can still uptake glycerol and convert it to PDO. In *E. coli*, only one *glpF* gene is present in the *glpFK* operon; the GlpF protein shows a high similarity (80.9%) to that (encoded from *glpFK*) of *K. pneumoniae* (Sun et al., 2003). The importance of GlpF in *E. coli* also seems to be limited only at low levels of glycerol (<8 mM), as glycerol itself may pass through the membrane by diffusion at high concentrations (Richey and Lin, 1972).

### **2.3.3 Simultaneous operation of fermentative and respiratory pathways**

The presence of the *dha* and *glp* regulons affords *K. pneumoniae* metabolic flexibility. The *dha* regulon enables fermentative utilization of glycerol, while the *glp* regulon enables respiratory metabolism of glycerol. During fermentative metabolism of glycerol, redox constraints force the production of PDO from glycerol, with a low ratio of carbon conversion yield to cell biomass. Moreover, ATP is formed through substrate-level phosphorylation, and NADH oxidation occurs through the formation of reduced metabolites. In the presence of oxygen, the redox constraints forcing the production of PDO are relieved; therefore, the carbon loss required for regeneration of NAD<sup>+</sup> is greatly reduced. As a consequence, glycerol utilization, biomass synthesis, and even the formation of many metabolites increase (Durnin et al., 2008). Interestingly, the fermentative route (*dha* regulon) is in operation under a wide range of oxygen levels, from microaerobic to highly aerobic conditions, as evidenced by PDO formation. For example, a high titer PDO (102.7 g/L) was produced under the condition

of 2.0 vvm aeration and even at 3.5 vvm, although the production was a little lowered in the latter case (Oh et al., 2012a). These results suggest that the fermentative and respiratory pathways might have evolved to complement each other for optimal control of cell growth under a wide range of aeration conditions except the strictly anaerobic one.

## **2.4 Carbon catabolite repression**

Carbon catabolite repression (CCR) refers to the suppression of the use of less-preferred carbon sources, when a more preferred carbon, most prominently glucose, is present. Cyclic AMP, cAMP receptor protein (CRP), and the enzyme EIIA<sup>Glc</sup>, an intermediate in the phosphorylation cascade of the PTS, are key players in the CCR of enterobacteria. The EIIA<sup>Glc</sup>, a cytosolic protein, exists in phosphorylated and unphosphorylated forms. The phosphorylated form stimulates membrane-bound adenylate cyclase, which catalyzes the formation of cAMP; the cAMP-CRP complex then induces expression of the catabolic genes of less-preferred carbon sources. The unphosphorylated form of EIIA<sup>Glc</sup> is also responsible for inhibiting the transport of less-preferred carbon sources by inhibitory binding to permeases, causing so-called inducer exclusion (Eppler and Boos, 1999; Eppler et al., 2002). Glycerol assimilation in *K. pneumoniae* is greatly inhibited by CCR in the presence of glucose. This repression is related to not only low cAMP and cAMP-CRP but also inhibition of GlpK (which functions mainly under aerobic conditions) by fructose 1,6-bisphosphate, a metabolite produced from glucose. The unphosphorylated EIIA<sup>Glc</sup> is also known to allosterically inhibit GlpK (Holtman et al., 2001). Anaerobic glycerol metabolism mediated by the *dha* operons is also suppressed by the presence of glucose. Enzyme-level inhibition similar to that of the aerobic GlpK has not been reported, but we have observed that, in *K. pneumoniae*, transcription of the *dha* operons is greatly reduced in the presence of glucose (Suman et al., 2017).

Interestingly, glycerol exerts catabolite repression on the assimilation of maltose in *E. coli*. To exert catabolite repression, glycerol must be phosphorylated to glycerol-3-phosphate (Hogema et al., 1998; Eppler et al. 2002). However, unlike other carbon sources, further metabolism of glycerol is not required. According to Eppler and Boos (1999), glycerol-3-phosphate reduces expression of MalT, a positive activator of all *mal* genes. The key players in this repression are adenylate cyclase, EIIA<sup>Glc</sup>, and CRP. The growth on maltose of the mutants lacking EIIA<sup>Glc</sup> or containing truncated adenylate cyclase was no longer repressed by glycerol, and CRP-independent transcription of *malT* was also not influenced by glycerol either. In addition, stimulation of adenylate cyclase by phosphorylated EIIA<sup>Glc</sup> was controlled by glycerol-3-phosphate. The mutation in adenylate cyclase relieved the repression caused by glycerol-3-phosphate (Eppler et al., 2002). Although not investigated, we assume that the glycerol repression on maltose catabolism also hold true in *K. pneumoniae*.

Glycerol also exerts catabolite repression on the assimilation of citrate in *K. pneumoniae*. Citrate can be used as the sole carbon and energy source during anaerobic growth of *K. pneumoniae*, but the presence of glycerol suppresses the expression of the citrate fermentation genes. A reduced concentration of the cAMP–CRP complex has been postulated to be the cause of this repression (Meyer et al., 2001). Further research is needed to test the hypothesis.

### **3. *K. pneumoniae* as a microbial cell factory for the production of commodity chemicals**

During glycerol fermentation, *K. pneumoniae* produces many metabolites, including PDO, BDO, and lactic acid. This section describes the biotechnological production of bulk chemicals from glycerol in *K. pneumoniae*. 3-HP is not a natural product during the growth of *K. pneumoniae* on glycerol. However, its production is possible using a minor modification of the PDO synthetic pathway and is included here. The biochemical pathways

used for the synthesis of these chemicals are explained, and the metabolic engineering used to improve their production is discussed.

### 3.1 1,3-Propanediol

PDO, also known as trimethylene glycol, is a promising platform chemical that has two hydroxyl groups. With terephthalic acid, it can be co-polymerized to the novel co-polymer polytrimethylene terephthalate (PTT). PDO also has applications in the food and cosmetic industries. Currently, commercial PDO is mainly produced by recombinant *E. coli* (Maervoet et al., 2011; Saxena et al., 2009). The global demand for PDO was 60.2 kilotons in 2012 and is predicted to reach approximately 150 kilotons by 2019. The market for PDO is growing rapidly, from \$157 million in 2012 with projections up to \$560 million in 2019 (MarketsANDMarkets, 2012; Lee et al., 2015).

*K. pneumoniae* is one of the best native producers of PDO from glycerol. GDHt and PDOR are two important enzymes for the conversion of glycerol to PDO (Fig. 3). Physiologically, the synthesis of PDO from glycerol is essential for the anaerobic growth of *K. pneumoniae*, as described previously. The production of PDO requires two cofactors, coenzyme B<sub>12</sub> and NAD(P)H. *K. pneumoniae* has a *de novo* pathway for the biosynthesis of coenzyme B<sub>12</sub> that comprises more than 20 genes. NAD(P)H is generated through the oxidative metabolism of glycerol, during the production of cell biomass and/or oxidized (pyruvate and acetate) or partially reduced (BDO, lactic acid) metabolites. The production yield of PDO from glycerol varies depending on the cost of carbon for NAD(P)H production; the lower the cost, the higher the yield of PDO from glycerol. NADPH can also be the electron donor for the conversion of 3-HPA to PDO when YqhD is present. However, the contribution of NADPH to PDO production in native *K. pneumoniae* seems to be limited because the PP pathway, which is the main source of NADPH, is not active when glycerol is

the sole carbon source. Furthermore, *K. pneumoniae* has very low transhydrogenase activity for the conversion of NADH to NADPH (Zhang and Xiu, 2009; Chen et al., 2011).

Several metabolic engineering strategies have been employed to improve PDO production by *K. pneumonia* (Table 2). Amplification of the reductive pathway by overexpressing DhaB, PDOR, or both has been attempted by many research groups. As indicated previously, 3-HPA is highly toxic, and its accumulation is detrimental to cell viability and toxic to PDO production. To reduce 3-HPA accumulation, overexpression of *dhaT* has been attempted. In one study, 3-HPA accumulation was reduced by *dhaT* overexpression, but no improvement in PDO production resulted (Hao et al., 2008). Overexpression of both DhaT and DhaD increased PDO production (56.3%) in batch culture but no increase in fed-batch culture (Chen et al., 2009). With another *K. pneumoniae* strain, DSM 2026, the same experiments were repeated but also with no increase in PDO production (Zheng et al., 2006). These results suggest that the reductive pathway in *K. pneumoniae* is evolutionarily well established (perhaps in concert with the oxidative pathway) and that PDO production is not limited by expression of the *dhaB* and *dhaT* genes of the reductive pathway. However, in one study where the *E. coli yqhD* gene was highly overexpressed to give ~10-fold higher PDOR activity (110 IU/mg total protein), a significant increase in PDO titer (25%; 67.6 g/L) and yield (from 0.53 to 0.62 mol/mol), along with less 3-HPA accumulation, was achieved (Zhu et al., 2009). This observation suggests that in the recombinant strain, NADPH can play an important role as a cofactor and that PDO production can be increased by modifying the reductive pathway. It is also probable that the effect of DhaB and PDOR on PDO production is highly dependent on the strain and/or culture conditions adopted.

Lactic acid, BDO, ethanol, and acetic acid are the major byproducts of PDO synthesis in *K. pneumoniae*. Inactivation of the metabolic pathways that give rise to these products has resulted in increased PDO synthesis. In particular, deletion of the *ldhA* gene encoding lactate

dehydrogenase has been highly beneficial. For example, by disrupting the *ldhA* gene, Xu et al. (2009b) significantly increased PDO production without lactic acid production. They observed increases in the PDO titer (95.4–102.1 g/L), conversion yield (0.48–0.52 mol/mol), and productivity (1.98–2.13 g/L h; Table 2). Similarly, Oh et al. (2012a) achieved the high PDO titer of 102.7 g/L by deleting *ldhA* in their own *K. pneumoniae* isolate. When lactic acid production was eliminated, more ethanol and succinic acid were generated from the pyruvate node. Reduction of these compounds has also been attempted by deleting *adhE* (encoding alcohol dehydrogenase) and *frdA* (encoding fumarate reductase). Although the production of ethanol and succinate was substantially reduced in the triple mutant ( $\Delta ldhA \Delta adhE \Delta frdA$ ), PDO production was only marginally increased compared with that of the single  $\Delta ldhA$  mutant. The failure of the triple mutant ( $\Delta ldhA \Delta adhE \Delta frdA$ ) to increase PDO production was mainly due to an increase in BDO production. Disruption of the BDO pathway encoded by the *bud* operon (*budA*, *budB*, and *budC*) has also been studied (Fig. 5). Although partially effective, individual inactivation of each of the three genes did not successfully decrease BDO production (Oh et al., 2012a; Wu et al., 2013). Deletion of the entire *bud* operon completely eliminated BDO synthesis (Kumar et al., 2016), but seriously hampered cell growth and glycerol consumption. Consequently, no increase in PDO production was resulted. Wu et al. (2013) attempted to decrease BDO production while increasing PDO production by inserting the *fdh* gene from *Candida boidinii* (NADH-forming formate dehydrogenase) into the *budC* locus. This protocol increased the PDO titer (62.3–72.2 g/L) and yield (0.47–0.57 mol/mol) and decreased the production of BDO by 52.2% and formic acid by 73.4%.

Acetic acid is one of the most toxic metabolites, accumulating in large quantities during glycerol fermentation by *K. pneumoniae* (Celińska, 2012). The toxic effects of acetate is realized even at the low concentration of 20 mM. Physiologically, acetate is a preferred metabolite for many *Enterobacter* sp. because its production is accompanied by ATP

generation. In a typical bioreactor run for PDO production by *K. pneumoniae*, the accumulation of acetic acid to >300 mM by the end is not uncommon, which often completely stops glycerol fermentation. Acetate is produced by Pta-Ack and PoxB when pyruvate formation is faster than its consumption, which is so-called ‘overflow metabolism’ (Fig. 6) (De Mey et al., 2007). In *K. pneumoniae*, the contribution of the *poxB* gene is not significant and its deletion does not much affect acetate production. In comparison, deletion of the *pta-ack* genes greatly reduces cell growth and results in high accumulation of pyruvate and pyruvate-based metabolites (Ko et al., 2017). Extensive studies to reduce or eliminate acetate production have been conducted in *E. coli*. Compared to *poxB*, *ackA* and/or *pta* were much more significant in reducing acetate production. However, deletion of the latter genes was not beneficial because the glycolytic flux and cell yield were seriously reduced and the accumulation of other byproducts such as pyruvate, lactate, and formate was greatly increased (Causey et al., 2004; De Mey et al., 2007). Several indirect approaches to decreasing pyruvate/acetate accumulation were investigated: stimulation of the anaplerotic pathway by overexpression of pyruvate carboxylase, PEP carboxylase, or PEP carboxykinase (Abdel-Hamid et al., 2001; Causey et al., 2004; Chao and Liao, 1993; Sanchez et al., 2005); overexpression of acetyl-CoA synthetase (ACS); increase of the TCA cycle throughput by deleting the transcriptional repressor *arcA* gene; and stimulation of the glyoxylate shunt by deleting the transcriptional repressor *iclR* gene (De Mey et al., 2007; Jeong et al., 2004; Lin et al., 2006). Moreover, the use of a mutant pyruvate dehydrogenase which is less sensitive to inhibition by NADH has also been examined (Kim et al., 2008). In *E. coli*, these approaches proved to be effective to varying extents under properly selected culture conditions. However, they have not yet been fully investigated in *K. pneumoniae*.

To eliminate byproduct formation, reduction of glycerol flux through the oxidative pathway at the glycerol node has also been attempted. The deletion of glycerol

dehydrogenase (*dhaD*) and/or dihydroxyacetone kinase (*dhaK*) resulted in decrease of the oxidative flux and increase of the conversion yield of glycerol to PDO. In addition, production of the byproducts lactate, ethanol, and succinate but not acetate decreased significantly, even without blocking the pathways leading to these byproducts (Seo et al. 2009; Horng et al. 2010). However, cell growth and glycerol consumption rate were also reduced. Interestingly, in these mutant strains, deletion of *dhaT* has almost no effect on PDO production. In the absence of *dhaT*, expression of an HOR, highly homologous to YqhD, was upregulated (see section 2.2.4).

The theoretical maximum yields of PDO on glycerol are 0.875 and 0.844 mol/mol under aerobic and anaerobic conditions, respectively (Celińska, 2012). However, experimentally, the yields are reported between 0.35 and 0.65 mol/mol (Saxena et al., 2009; Sun et al., 2008). The low yield is mainly attributed to the accumulation of unnecessary byproducts and inefficient metabolism for generating reducing equivalents (NADH) and ATP. If acetyl-CoA is fully oxidized through the TCA cycle (without byproducts formation) and the activity of the electron transport chain (ETC) oxidizing NADH is properly controlled, a significant increase in PDO yield is expected. Attaining this goal requires extensive pathway engineering of cellular metabolism. In bioreactor operation, the aeration rate should be optimized. Oxygen regulates gene expression and the enzymatic reactions of glycerol metabolism and determines the diversion of DHA to the PP pathway. Regeneration of NADH by the TCA cycle and its oxidation by ETC are also controlled by oxygen levels.

### **3.2 3-Hydroxypropionic acid**

As with PDO, 3-HP is an important platform chemical and has been selected by the US Department of Energy (DOE) as one of the top 12 chemicals obtainable from biomass. 3-HP is a C3 bifunctional molecule and a structural isomer of lactic acid (Werpy and Petersen, 2004). 3-HP can be converted into a variety of useful chemicals such as acrylic acid and acryl



amide (Pina et al., 2011; Kumar et al., 2013a). 3-HP can be synthesized by chemical routes from acrylic acid, 3-propiolactone, 3-hydroxypropionitrile, allyl alcohol, vinyl acetate, and PDO. However, none of the chemical processes is commercially feasible at present due to the high cost of the starting materials, toxicity of intermediates, and/or the environmental incompatibility of the processes (Jiang et al., 2009; Pina et al., 2011; Kumar et al., 2013a). Many microorganisms, including both prokaryotes and eukaryotes, can naturally synthesize 3-HP as either an intermediate or end product through a range of metabolic pathways. However, the production of 3-HP by these native microorganisms is too low to be commercially meaningful (Jiang et al., 2009; Kumar et al., 2013a).

In studies pursuing commercial production of 3-HP, glycerol is first converted to 3-HPA by GDHt. 3-HPA then is converted to 3-HP via the coenzyme A (CoA)-dependent or CoA-independent pathway (Fig. 3). In the CoA-independent pathway, 3-HPA is oxidized to 3-HP by aldehyde dehydrogenase (ALDH); in the CoA-dependent pathway, 3-HPA is converted to 3-HP via 3-HP-CoA and 3-HP-phosphate. In *K. pneumoniae*, both the CoA-dependent and CoA-independent pathways are present. However, the titer of 3-HP produced by the CoA-dependent pathway is low as <5.0 g/L (Luo et al., 2011 and 2012). High production of 3-HP requires the disruption of oxidoreductases (e.g., DhaT, YqhD) and the overexpression of a proper aldehyde dehydrogenase (ALDH). Ashok et al. (2013a) developed a recombinant *K. pneumoniae* in which *dhaT* and *yqhD* were deleted, and the homologous *puuC* gene encoding the NAD<sup>+</sup>-dependent  $\gamma$ -glutamyl- $\gamma$ -aminobutyraldehyde dehydrogenase was overexpressed. The recombinant *K. pneumoniae* produced 3.8 g/L 3-HP in 12 h of flask culture, but only under appropriate aeration (i.e., microaerobic) conditions. Under anaerobic conditions, PDO (instead of 3-HP) was obtained as the main product, even though two oxidoreductases were disrupted and *puuC* was highly overexpressed. In contrast, highly aerobic conditions produced a high cell mass without much accumulation of either 3-HP or

1,3-PDO. In a glycerol-fed batch bioreactor experiment under a constant dissolved oxygen (DO) concentration of 5% (considered “proper” microaerobic condition), the recombinant *K. pneumoniae*  $\Delta dhaT \Delta yqhD$  overexpressing both PuuC and DhaB produced >28 g/L 3-HP in 48 h, with a yield of >40%. In contrast, the same strain produced only 10.5 g/L 3-HP when cultivated under the higher 10% DO conditions. Several important conclusions regarding 3-HP production by *K. pneumoniae* were made, as follows. First, in addition to DhaT and YqhD, *K. pneumoniae* has more unidentified oxidoreductases that can produce PDO from 3-HPA. Second, to produce 3-HP rather than PDO, proper aeration to regenerate  $NAD^+$  is essential. Third, excessively high aeration decreases the expression of the Dha regulon and synthesis of coenzyme B<sub>12</sub>, an essential cofactor for GDHt catalysis.

To eliminate PDO production even under limited aeration conditions, Ko et al. attempted to identify and disrupt other potential PDORs from *K. pneumoniae* (Ko et al., 2015). A mutant strain devoid of *dhaT*, *yqhD*, *ahpF*, and *adhE* genes was developed; however, the mutant neither eliminated 1,3-PDO production nor increased 3-HP production. Again, this result suggests that *K. pneumoniae* has more unidentified oxidoreductases, and thus, the complete elimination of 1,3-PDO production during 3-HP production is highly challenging. To address the important problem, of  $NAD^+$  regeneration and coenzyme B<sub>12</sub> production, Ashok et al. (2013b) studied anaerobic respiration using nitrate as an external electron acceptor. They attempted to regenerate  $NAD^+$  from NADH by nitrate reduction while maintain anaerobic conditions to synthesize and stabilize the oxygen-sensitive coenzyme B<sub>12</sub>. Disruption of the *glpK* gene (encoding for glycerol kinase) was also necessary because with *glpK* intact, the rate of anaerobic respiration was too fast, and most of the glycerol was metabolized through GlpK to produce biomass, as in the case of highly aerobic cultivation. The anaerobic glycerol fermentation in the presence of nitrate was successful: 1.74 g/L 3-HP was obtained from a 12-h flask culture and 22 g/L 3-HP was produced in a 48-

h fed-batch bioreactor culture (Ashok et al., 2013b) (Table 3). However, disadvantages were noted, including the requirement for large amounts of nitrate and the toxic effect of nitrite generated as a reduced intermediate of nitrate. In a recent study, Li et al. (2016) optimized a promoter for ALDH expression, culture conditions, and metabolic flux to achieve high-level production of 3-HP in their *K. pneumoniae* isolate. One recombinant strain, *K. pneumoniae* (pTAC-*puuC*) expressing *puuC* under the IPTG-inducible *tac* promoter, produced 73.4 g/L 3-HP in a bioreactor with a molar yield of glycerol of 0.52 and productivity of 1.53 g/L h. Further, elimination of the *ldh1*, *ldh2*, and *pta* genes elevated the titer and molar yield to 83.8 g/L 3-HP and 0.54 mol/mol, respectively, with decreased cell growth and productivity. This titer of 3-HP is the highest produced by *K. pneumoniae* to date.

As another approach to addressing the problem associated with NAD<sup>+</sup> regeneration and coenzyme B<sub>12</sub> synthesis in 3-HP production, co-production of 3-HP and PDO has been investigated. The simultaneous production of 3-HP and PDO can eliminate the dependency on oxidative metabolism of glycerol (and electron transport chain), as the cofactor required for 3-HP production is regenerated by PDO production or vice versa. Furthermore, the problems associated with the expression of genes for the assimilation of vitamin B<sub>12</sub> and glycerol can be alleviated substantially because co-production can be performed under anaerobic or microaerobic conditions (Kumar et al., 2012, 2013b). Ashok et al. (2011) developed a recombinant strain of *K. pneumoniae* DSMZ by overexpressing ALDH and deleting *dhaT*. With *dhaT* intact, PDOR activity was too high compared with that of ALDH, even with overexpression of the latter enzyme from a multicopy plasmid. The recombinant *K. pneumoniae* DSMZ accumulated 16.0 g/L 3-HP and 16.8 g/L PDO in 24 h, and the cumulative yield of these two metabolites on glycerol was 51% (Table 3). Huang et al. (2012) studied the same co-production using their *K. pneumoniae* isolate overexpressing ALDH and obtained 24.4 3-HP and 49.9 g/L PDO, respectively, after 24 h under anaerobic conditions.

The cumulative molar yield of the two metabolites reached 0.61 (0.18 for 3-HP and 0.43 for PDO). In a following study, they showed that the maximum titer of 48.9 g/L 3-HP (along with 25.3 g/L PDO) was achieved at 1.5 vvm, while the highest concentration of 38.4 g/L PDO (along with 16.6 g/L 3-HP) was obtained at 0.1 vvm (Huang et al., 2013a).

During 3-HP production from glycerol by *K. pneumoniae*, the accumulation of such byproducts as lactic acid, ethanol, acetic acid, formic acid, and 2,3-BDO is a serious consideration, as in the case of 1,3-PDO production (Ashok et al., 2013a,b; Li et al., 2016). Deletion of *ldhA* reduces lactic acid formation and increases the 3-HP yield. However, deletion of other genes such as *adhE*, *pfl*, *budBAC*, and *pta-ack* had little desired effect and seriously hampered cell growth and glycerol assimilation (unpublished data). Even with the co-production of 3-HP and 1,3-PDO, where oxidative glycerol assimilation was greatly reduced, accumulation of these byproducts was a serious drawback (Ko et al., 2017). To prevent the formation of lactic acid along with other byproducts, Kumar et al. (2013b) employed resting cells of recombinant *K. pneumoniae* J2B overexpressing ALDH and devoid of lactate dehydrogenase (*ldhA*). For this strain, final titers of 3-HP and PDO of 22.7 and 23.5 g/L, respectively, were obtained without lactic acid accumulation. The cumulative product yield increased to 0.77.

In addition to *K. pneumoniae*, *E. coli* has been extensively studied for 3-HP production from glycerol. Several successful results have been reported by Samsung Ltd. Co. Korea. One study (Chu et al., 2015) used an engineered strain (*E. coli* W3110  $\Delta$ *ackA-pta*  $\Delta$ *yqhD\_dhaB\_mutant gabD4*) harboring an active ALDH mutant (designated as GabD4) from *Cupriavidus necator*, producing 71.9 g/L 3-HP with a productivity of 1.8 g/L h. However, unlike *K. pneumoniae*, *E. coli* does not synthesize vitamin B<sub>12</sub> naturally, and it was necessary to add the coenzyme externally. Furthermore, a significant amount of glucose for cell growth was required because for *E. coli*, glycerol is a less-preferred carbon source than glucose.

Consequently, the researchers at Samsung initially grew cells to a high density (40–300 OD<sub>600</sub>) on glucose and then produced 3-HP by feeding glycerol (Chu et al., 2015; Kim et al., 2014a). In comparison, *K. pneumoniae* can produce a high concentration of 3-HP growing purely on glycerol at a much lower cell concentration (OD<sub>600</sub>, <20) (Huang et al., 2013a,b; Li et al., 2016). Together, the use of glucose for cell proliferation, exogenous addition of expensive cofactor vitamin B<sub>12</sub>, and low cumulative yield of 3-HP (on glucose plus glycerol) increase the production cost. Thus, *K. pneumoniae* can be considered a better biocatalyst than *E. coli*. However, for the commercial production of 3-HP, many other factors are important, including biosafety, process stability, and downstream processing. Further studies to assess the potential of these strains as hosts for 3-HP production are needed.

### 3.3 2,3-Butanediol

BDO has many applications in the pharmaceutical, biomedical, and other chemical industries for the production of printing inks, perfumes, fumigants, spandex, moistening and softening agents, and plasticizers (e.g., cellulose nitrate, polyvinyl chloride, polyacrylates) (Celińska and Grajek, 2009; Ji et al., 2011). BDO can also be used as an antifreezing agent and octane booster for petrol as is or can be converted to useful derivatives such as 1,3-butadiene, and diacetyl and methyl ethyl ketone.

Unlike 3-HP and PDO, BDO is a product of oxidative metabolism. Of the three stereoisomers of BDO, *K. pneumoniae* mainly produces the *meso* form (Ji et al., 2011). BDO is synthesized from pyruvate via  $\alpha$ -acetolactate and acetoin. The C5 intermediate,  $\alpha$ -acetolactate, is formed from two molecules of pyruvate by self-condensation catalyzed by  $\alpha$ -acetolactate synthase (ALS; *budB*).  $\alpha$ -Acetolactate is decarboxylated to acetoin by  $\alpha$ -acetolactate decarboxylase (*budA*), and acetoin is reduced to BDO by 2,3-butanediol dehydrogenase/acetoin reductase (*budC*) using NADH as a reductant. In the presence of oxygen,  $\alpha$ -acetolactate is spontaneously decarboxylated to diacetyl, further reduced to acetoin

by action of diacetyl reductase, and then reduced to BDO.  $\alpha$ -Acetolactate is also the precursor of branched-chain amino acids (valine, leucine, and isoleucine); thus, elimination of the *bud* operon can arrest *K. pneumoniae* growth when cultured on glycerol minimal medium (Kumar et al., 2016). In the anaerobic process, one NADH molecule is required for the conversion of two pyruvates to one BDO, while the aerobic process requires two NADH (Fig. 5). The enzyme DhaD may also contribute to BDO formation (see Section 2.2.1). In the BDO pathway itself, the conversion of two pyruvates to BDO regulates or is regulated by the intracellular NADH/NAD<sup>+</sup> ratio, similar to other fermentative pathways. Another important role of the BDO pathway is to reduce intracellular acidification by converting acids to the neutral metabolite, BDO (Celińska and Grajek, 2009; Ji et al., 2011).

Although BDO is one of the main byproducts of glycerol fermentation by *K. pneumoniae* (Petrov and Petrova, 2009; Durgapal et al. 2014), its production from glycerol has not been studied extensively. In few studies using glycerol as the carbon source, Petrov and Petrova (2009) focused on the effects of pH and aeration with their *K. pneumoniae* isolate G31. BDO production was greatest at alkaline initial pH when tested at a pH range of 5–8. Without pH control a sharp pH drop occurred initially due to the production of organic acids, and this pH drop triggered the synthesis of BDO, which is considered the only non-inhibitory compound produced by the oxidative metabolism of glycerol. Once neutral pH is restored with the rising concentration of BDO, catabolism shifted toward the production of ATP-generating acetic acid. The amount of BDO generated in each cycle (pH drop and neutralization) was dependent on the extent of the pH drop. In one study, a BDO concentration of 49.8 g/L was reached when the initial pH was set at 8.0 and no pH control was given during the rest of fermentation. PDO was the major byproduct (11.6 g/L) (Table 4) and its production decreased by exclusion of Co<sup>2+</sup>, an essential component of coenzyme B<sub>12</sub>. They also found that intensive aeration led to a significant increase in BDO production; an

increase in aeration rate from 1.1 to 2.2 vvm enhanced the yield from 0.38 to 0.46 mol/mol and productivity from 0.24 to 0.36 g/L h. BDO production was also increased by forced pH fluctuations of the culture medium with discrete  $\Delta\text{pH}$  values (1.0, 2.0, and 3.0) at predetermined time intervals (6, 12, and 24 h). The highest BDO concentration (70.0 g/L) was produced by increasing the pH by one unit ( $\Delta\text{pH} = 1.0$ ) every 12 h (Petrov and Petrova, 2010). The mechanism behind the effect of this peculiar pH change on BDO production has not been elucidated. Although BDO is readily produced from glycerol by *K. pneumoniae*, the volumetric productivity is low due to the long fermentation time (generally >150 h). Huang et al. (2013b) isolated two *Klebsiella* strains that produce BDO plus acetoin with a total yield of 0.44–0.45 (g/g). One isolate utilized 64.5 g/L glycerol in 78 h and produced 24.9 g/L BDO, 3.9 g/L acetoin, and 3.2 g/L ethanol. Again, the volumetric productivity was not high.

Metabolic engineering efforts with *K. pneumoniae* to improve BDO production from glycerol are limited. Using glucose as carbon source, Kim et al. (2014b) reported that overexpression of the BDO pathway (*budA*, *budB*) in the *ldhA* deficient *K. pneumoniae* strain KCTC2242 improved BDO production and resulted in 90 g/L BDO, with a productivity of 2.75 g/L. h. Because both glucose and glycerol are converted through the common intermediate pyruvate, BDO production from glycerol is also expected to increase by amplification of the *bud* operon. Further work with glycerol as carbon source is required to prove this effect. Recently, co-production of BDO and PDO from glycerol has also been attempted (Park et al., 2017). In glycerol fermentation, both alcohols always appear simultaneously and co-production seems to be beneficial for cells to maintain redox balance during anaerobic growth. Park et al. (2017) found that by using the *K. pneumoniae* mutant deficient of *ldhA* and *mdh*, 50.1 g/L BDO was produced along with 63.8 g/L PDO in 75 h. The ratio of the two alcohols varied according to aeration; with increasing aeration, the ratio of [BDO]/[PDO] increased as observed by Petrov and Petrova (2010). In the 5,000 L pilot-

scale fed-batch fermentation using crude glycerol, they could produce 114 g/L diols (70 g/L PDO and 44 g/L BDO) with the yield of 0.60 diols/glycerol (g/g) and a productivity of 2.2 g/L. h of diols. If separation is not an issue, co-production of the two diols can be a good option.

### 3.4 Lactic acid

Lactic acid has applications in the food, chemical, cosmetic, and pharmaceutical industries. Lactic acid is a monomer that can be polymerized to yield the biodegradable plastics polylactic acid (PLA) and poly(3-hydroxybutyrate-*co*-lactate). Two optical isomers of lactic acid, D- and L-lactic acid, are produced in racemic mixtures depending on the chemical route used. Microbial fermentation can yield optically pure isomers. Currently, nearly all commercial lactic acid comes from microbial fermentation (Abdel-Rahman et al., 2013; Abdel-Rahman and Sonomoto, 2016).

Because of the availability of highly efficient lactic-acid-producing microbes derived from *E. coli*, lactic acid bacteria, and even acid-tolerant yeasts (Abdel-Rahman et al., 2013), *K. pneumoniae* has received little attention as a lactic acid producer. However, as previously described, lactic acid is a major byproduct of glycerol fermentation by *K. pneumoniae*, and its potential as a producer is high. This bacterium contains metabolic pathways for the production of both L- and D-isomers (Fig. 7). They are synthesized through the pyruvate and/or methylglyoxal pathways in *K. pneumoniae* (Ashok et al., 2011). When produced from pyruvate, lactate dehydrogenase (*ldhA*) reduces pyruvate using NADH as reductant. In the methylglyoxal route, DHAP is converted to methylglyoxal, which can be transformed to both D- and L-lactic acid. For the production of L-lactic acid, methylglyoxal is reduced to L-lactaldehyde, which is subsequently oxidized to L-lactic acid. Using glutathione, D-lactic acid is obtained through simultaneous aldehyde group oxidation and keto group reduction of methylglyoxal (Booth et al., 2003; Booth, 2005). The methylglyoxal route has several



disadvantages. Methylglyoxal is highly toxic, and its accumulation can severely impair other metabolisms and even lead to cell death. Second, the route is energetically inefficient, with an ATP yield of  $-1$ . The L- or D-lactic acid product can be converted back into pyruvate by respiratory lactate dehydrogenase (Mazumdar et al., 2010, 2013).

Several studies showing the potential of *K. pneumoniae* as a lactic acid producer are available (Table 4). For example, *K. pneumoniae* DSMZ is reported to accumulate 22.7 g/L lactic acid in 24 h during co-production of 3-HP and PDO from glycerol. The titer of lactic acid was more than that of any of the targeted products, and the yield of lactic acid on glycerol alone was 0.31 (Ashok et al., 2011). Similarly, fed-batch cultivation of *K. pneumoniae* BLh-1 under oxygen-limited conditions gave lactic acid as the main product, outcompeting PDO; 59 g/L lactic acid was produced in 40 h when crude glycerol from biodiesel synthesis without any purification was used (Rossi et al., 2013). Durgapal et al. (2014) also found that *K. pneumoniae* J2B and DSMZ strains produced lactic acid at 37.0 and 47.8 g/L, respectively (Table 4); in both strains, lactic acid levels surpassed those of the target product PDO. Recently, Feng et al. (2014) engineered *K. pneumoniae* ATCC25955 by overexpressing *ldhA* and deleting *dhaT* and *yqhD* (recall that these two genes are mainly responsible for PDO production). They found that the recombinant produced 142.1 g/L of optically pure D-lactic acid from glycerol in fed-batch cultivation under microaerobic conditions (2.5 vvm and 400 rpm). This result is the highest lactic acid concentration produced from glycerol reported to date. *E. coli* has also been engineered for the production of D- and L-lactate from glycerol (Mazumdar et al., 2010, 2013). For diverting glycerol flux toward L-lactic acid in *E. coli*, the chromosomal copy of D-lactate dehydrogenase was replaced with *Streptococcus bovis* L-lactate dehydrogenase (Mazumdar et al., 2013). Moreover, gene encoding *lldD* (respiratory L-lactate dehydrogenase) was inactivated to prevent the consumption of L-lactic acid, and the methylglyoxal route ( $\Delta mgsA$ ) was blocked

to prevent production of toxic methylglyoxal and formation of racemic mixtures of D- and L-lactic acid (Fig. 7). Furthermore, the respiratory route for glycerol assimilation was overexpressed (*glpK* and *glpD*) to improve the ATP yield of the metabolic pathway through transfer of electrons from glycerol-3-phosphate to oxygen (i.e., oxidative phosphorylation). The engineered *E. coli* strain ( $\Delta pflB \Delta pta \Delta adhE \Delta frdA \Delta mgsA \Delta lldD \Delta ldhA::ldhA\_glpK\_glpD$ ) produced 50 g/L of L-lactate from 56 g/L of crude glycerol, at a yield 93% of the theoretical maximum and with high optical (99.9%) and chemical (97%) purity (Mazumdar et al., 2013). The approaches proven successful in *E. coli* should be applicable to *K. pneumoniae*. These studies, along with the results by Feng et al. (2014), suggest that *K. pneumoniae* is a good host for the production of D- and L-isomers of lactic acid, with high optical purity, from glycerol.

### 3.5 Ethanol and other metabolites

*K. pneumoniae* has also been studied for the production of ethanol, succinic acid, 2-ketogluconic acid, catechol, *cis,cis*-muconic acid and 2-butanol although its potential for such production has not been fully explored (Cheng et al., 2013; Jung et al., 2015; Oh et al., 2014; Wang et al. 2015; Wei et al., 2013). Of these chemicals, ethanol production has been relatively well studied. Ethanol is a renewable energy source and is widely used as a fuel additive for partial gasoline replacement. Currently, commercial ethanol is produced from the fermentation of sugar and starch. However, glycerol is one of the cheapest substrates for ethanol production, estimated as nearly 40% cheaper than sugar-based production (Yazdani and Gonzalez, 2007). This factor has encouraged the development of microbial strains for fermentative conversion of glycerol to ethanol. The metabolic route by which ethanol is synthesized from glycerol is shown in Fig. 6. Oh et al. (2011, 2012b) constructed a mutant strain of *K. pneumoniae* (termed GEM167) through  $\gamma$ -irradiation. PDO synthesis in this mutant strain was lower than that of wild type (0.2 vs. 7.93 g/L, respectively), while ethanol

accumulation was higher (8.6 vs. 1.1 g/L, respectively). The mechanism underlying these changes remains unclear. Introduction of the pyruvate decarboxylase (*pdc*) and aldehyde dehydrogenase (*adhII*) genes into mutant strains lacking lactate dehydrogenase (*ldhA*) resulted in the production of 31.0 g/L (673.9 mM) ethanol, with a yield of 0.89 (mol/mol) and productivity of 1.2 g/L h (Oh et al., 2012b), the highest level of ethanol production from glycerol reported to date. More extensive studies including estimation of the maximum achievable titer and microbial tolerance to high-concentration ethanol, both of which are key factors for commercialization, are required to assess the potential to produce ethanol from glycerol by the use of *K. pneumoniae*.

#### **4. Challenges to bioconversion of glycerol using *K. pneumoniae* as a biocatalyst**

The design of microbial cell factories is gaining unprecedented momentum as metabolic engineering is progressively aided by advances in synthetic biology and multi-omics analyses. Several bio-based chemicals have already been launched successfully in the marketplace, and others are in the pipeline. For a bulk chemical to be considered for commercial production, the production process should achieve a product concentration of 100 g/L, carbon yield of 0.50 (g/g), and volumetric productivity of 2.0 g/L/h. The aforementioned products produced by *K. pneumoniae* from glycerol are no exception. Challenges to the use of *K. pneumoniae* as a microbial cell factory using glycerol as a carbon source are described.

##### **4.1 Pathogenicity**

The pathogenicity of *K. pneumoniae* is one of the major obstacles to its commercial application. *K. pneumoniae* is a Bio-safety Level 2 microorganism and causes nosocomial and urinary tract infections. Moreover, capsulated *K. pneumoniae* cells are difficult to separate from culture broth, complicating the downstream processing. The biotechnological application of this bacterium will require attenuation of its pathogenicity and reduction of biosafety concerns. *K. pneumoniae* produces a number of virulence factors that contribute to

its pathogenicity, including lipopolysaccharide (LPS), capsular antigens, fimbrial adhesins, siderophores, and O antigens (Albertí et al., 1993; Fang et al., 2004; Podschun and Ullmann 1998). The mechanism by which this bacterium causes disease is still obscure because most studies have been conducted on a limited number of virulence factors (El Fertat-Aissani et al., 2013).

In the last decade, efforts have been made to overcome the pathogenicity of *K. pneumoniae* (Shrivastav et al., 2013), suggesting hope for the generation of nonpathogenic *K. pneumoniae*. The two major pathogenic determinants in *K. pneumoniae* are the capsular polysaccharides (CPS), which cover the entire bacterial surface, and LPS, expressed on the outer bacterial membrane. LPS consists of lipid A, a core oligosaccharide, and an O-antigenic polysaccharide. In a highly virulent strain of *K. pneumoniae*, mutations in genes involved in LPS biosynthesis (*waaC*, *waaF*, and *wabG*) resulted in a decrease in bacterial colonization and virulence (Izquierdo et al., 2003; Jung et al., 2013). Mutant strains devoid of the outer core LPS were also found to be deficient in the “cell-attached” capsular polysaccharides covering the bacterial surface. These mutations abolished the highly virulent characteristics of pathogenic *K. pneumoniae* when tested in different animal models. Furthermore, these mutants were more sensitive to several hydrophobic compounds than were wild-type strains. Reintroduction of the *waaC*, *waaF*, and *wabG* genes into *K. pneumoniae* rescued the pathogenic properties, confirming the role of these genes in pathogenicity.

Another important study was conducted by Lin et al. (2012), in which eight highly conserved residues of the MagA protein (G308, G310, G334, G337, R290, P305, H323, and N324) encoded by the mucoviscosity-associated gene A (*magA*) of *K. pneumoniae* were subjected to site-directed mutagenesis. The *magA* gene contributes to the biosynthesis of K1 capsular polysaccharide (CPS), which correlates with pathogenic phenotypes including mucoviscosity, serum and phagocytosis resistance, and virulence. Alanine substitutions at

R290A or H323A abolished MagA function, with annihilation of CPS production, serum resistance, and virulence. Some *K. pneumoniae* strains showing less pathogenicity were also isolated. For example, *K. pneumoniae* strain J2B produced less LPS and demonstrated a high sensitivity to many antibiotics and lower pathogenicity. Moreover, its biomass was readily separated from fermentation broth by centrifugation (Arasu et al., 2011). Future studies should include further engineering of these less pathogenic strains to make them completely nonpathogenic, followed by careful evaluation of their toxicity.

#### **4.2 Complex glycerol metabolism**

Although glycerol metabolism has been extensively studied, the mechanism and regulation of its gene expression in *K. pneumoniae* have not been fully elucidated. The diversity of glycerol metabolism should be a great advantage for *K. pneumoniae* survival in different environments and for its biotechnological utilization as well. However, a high level of complexity poses difficulty in the engineering of a strain to suit our purposes. Metabolic complexity is likely responsible in part for the fact that, despite serious attempts, the molar yield of many important products such as PDO and 3-HP is still far below the theoretical maximum. Efforts based on our current knowledge, such as the overexpression of DhaB, DhaT, YqhD and AldH and inactivation/deletion of GlpK, DhaD, DhaK and DhaKLM, were not satisfactory (Maervoet et al. 2011; Kumar et al. 2013a; Kumar et al., 2016). Another challenge is to understand how the ratio of glycerol distribution between respiratory and fermentative routes is controlled. Because the  $K_M$  of glycerol kinase toward glycerol is low, it has been speculated that a major fraction of glycerol flows through the respiratory route when oxygen is present. However, some experimental results do not support this hypothesis. For example, deletion of *glpK* does not decrease the glycerol flow through the oxidative pathway in the presence of oxygen (Ashok et al. 2013a). In fact, wild-type and mutant ( $\Delta glpK$ ) strains of *K. pneumoniae* demonstrate similar cell growth, glycerol consumption, and PDO

production profiles under (micro)aerobic conditions. These observations suggest the presence and/or involvement of other kinases.

Another challenging issue for the use of *K. pneumoniae* is its inefficient TCA cycle (Cabelli, 1955). Several genes, including isocitrate dehydrogenase (*icd*), fumarase (*fumA*), and malate dehydrogenase (*mdh*), in *K. pneumoniae* are significantly less transcribed than those in *E. coli* (unpublished data). Furthermore, the activity of isocitrate dehydrogenase was more than sevenfold lower in *K. pneumoniae* than in *E. coli*. If the TCA cycle is inefficient, achieving a high cell density, which is essential for improving the productivity of target metabolites, is difficult. Moreover, achieving high-yield generation of NADH with little consumption of glycerol carbon is difficult. Furthermore, the overflow metabolism that produces highly toxic acetate becomes more serious. These problems have been well documented in PDO production. In a recent study, Kumar et al. (2016) attempted to increase PDO production yield by blocking the lactate and BDO pathways, while diverting carbon flux through the TCA cycle for efficient regeneration of NADH. However, this effort resulted in the detrimental accumulation of pyruvate, acetate, and ethanol, causing low NADH regeneration and low PDO production. Neither increased aeration nor the addition of good nitrogen sources (complex nitrogen sources) alleviated the accumulation of these intermediates or activated the TCA cycle. Furthermore, the accumulation of pyruvate quickly terminated glycerol assimilation. *K. pneumoniae* has a well-established BDO production pathway that likely compensates for its inefficient TCA cycle, avoiding the overflow metabolism and carbon traffic at the pyruvate node. If this is the case, co-production of BDO along with PDO or 3-HP, rather than single production of PDO or 3-HP, should be more appropriate, which suits the nature of *K. pneumoniae*. In fact, when BDO and PDO were co-produced, more PDO was generated than during PDO production alone. More studies are required to gain a better understanding of glycerol metabolism, the slow operation of the

TCA cycle in *K. pneumoniae*, and its effect on the selection of target products and/or development of suitable strains and processes. Otherwise, it will be difficult to exploit this potential biocatalyst for commercial manufacturing of valuable chemicals.

#### **4.3 Toxicity of intermediate metabolites and end-products**

A general but major challenge to the use of microbial cell factories for producing bulk chemicals is the toxic effect of target products and metabolic intermediates at high concentrations. In particular, during the production of PDO or 3-HP from glycerol, the highly toxic intermediate 3-HPA is inevitably generated. Aldehydes generally damage DNA, inactivate enzymes, and inhibit DNA synthesis. 3-HPA causes irreversible cessation of metabolic activity and cellular growth, even at very low concentrations of 15–30 mM (Hao et al., 2008; Zheng et al., 2008). Native *K. pneumoniae* does not accumulate 3-HPA at toxic levels during PDO production, but in engineered strains that overexpress DhaB, high 3-HPA accumulation is often experienced. To avoid 3-HPA accumulation, enzymes that consume 3-HPA, ALDH (for 3-HP production), and PDOR (for PDO production) are expressed at sufficiently (sometimes excessively) high levels compared with that of DhaB (Lim et al., 2016). Intermediates such as methylglyoxal, DHA, DHAP, and glyceraldehyde-3-phosphate are also known to be toxic.

Organic acids and alcohols are toxic at high concentrations. Generally, organic acids are more toxic than their corresponding alcohols because they disturb the intracellular pH and have anion-specific effects on metabolism (Chun et al., 2014; Warnecke and Gill, 2005). When accumulated in cell membranes, organic compounds damage membrane integrity and inhibit the activity of membrane-bound enzymes. In *K. pneumoniae*, cell growth and glycerol assimilation are severely inhibited by high concentrations of PDO or 3-HP. Furthermore, enzymes that play major roles in 3-HP and PDO production, such as DhaB, DhaT, and AldH, are inhibited by 3-HP and PDO (Kumar et al. 2013b). Furthermore, *K. pneumoniae* is

sensitive to 24 g/L acetic acid, 26 g/L lactic acid, and 17 g/L ethanol under aerobic conditions and 15, 19, and 26 g/L, respectively, under anaerobic condition (Celińska, 2012; Cheng et al., 2005). The inhibition of pathways by target product(s) and/or unwanted byproducts often places a limit on the maximum attainable titer of bulk chemicals.

Several studies have reported engineered improvements in microbial tolerance against biochemicals and biofuels, especially in *E. coli*. Modifications of exporter proteins, heat shock proteins, membrane composition, and stress responses have been attempted. Moreover, *in situ* recovery of target products has been attempted for hydrophobic targets. Some tolerant strains have been developed, but unfortunately, without appreciable increases in productivity (Dunlop, 2011; Dunlop et al., 2011; Jarboe et al., 2011). Such efforts to improve tolerance to chemicals have not been reported for *K. pneumoniae*.

#### **4.4 Availability of glycerol**

The global market for glycerol is known to be unpredictable and complex. Glycerol in the current market mainly is from the biodiesel industry, and its supply is directly affected by the use of biodiesel. In contrast, glycerol demand depends on many industries that use it as feedstock. Between 2001 and 2011, the price varied significantly, from \$0.70/kg to \$1.70/kg for pure glycerol and from \$0.04/kg to \$0.33/kg for crude glycerol (Ayoub and Abdullah, 2012; Ciriminna et al., 2014; Kong et al., 2016; Quispe et al., 2013). Nevertheless, glycerol prices, for pure as well as crude, have come down significantly since 2004, when many biodiesel production plants initiated operation. The countries producing biodiesel in large quantities (in million liters) include Germany (2900), Brazil (2300), Argentina (2100), France (2000), USA (1200), Spain (1100), Italy (800), Indonesia (700), and Thailand (600) (Kong et al., 2016). The Asia-Pacific region overtook Europe as the largest glycerol market in the world in 2009 and has remained so ever since. Market growth is driven by increases in the



uses for glycerol in such sectors as pharmaceuticals, personal hygiene, and food and beverage production (Quispe et al., 2013).

The production of crude glycerol is expected to reach 6 million tons by 2025 (Ciriminna et al., 2014). At present, the surplus of glycerol from biodiesel and oleochemicals is assumed to be sufficient to meet these new demands. However, the scenario may change in the future with rapid growth of glycerol-based industries and a limited supply of crude glycerol. For example, if 3-HP produced from glycerol were used as a substitute for the chemical acrylic acid (global production, 5.85 million tons as of 2014), approximately 7 million tons of glycerol/year would be needed. If we include other platform chemicals such as PDO, BDO, and ethanol, the demand will be even higher, much more than the expected supply. The growth of glycerol-based biotechnology will eventually be constrained by the supply and price of crude glycerol as a carbon source.

## **5. Concluding remarks**

Glycerol is an excellent substrate for the production of biochemicals and biofuels. Although the biodiesel industry has grown rather slowly in recent years, crude glycerol is still cheap and generated in large quantities. *K. pneumoniae* has already been successfully employed for the production of PDO, 3-HP, BDO, lactic acid, and ethanol from glycerol at industrially attractive levels. Other metabolites such as acetic acid, pyruvic acid, malic acid, fumaric acid, alanine, *n*-butanol, 1,2-propanediol, propionic acid, formic acid, and hydrogen can also be produced from glycerol by *K. pneumoniae*, although these have not yet been extensively explored. At present, sugar-based bioprocesses are prevalent, and no bioprocesses employing *K. pneumoniae* and glycerol are commercially available. The pathogenicity of *K. pneumoniae* and other technical issues are existing challenges to its use. However, advancements in the fields of metabolic engineering, synthetic biology, systems biology, and evolutionary engineering will enable us to further exploit the advantageous nature of *K.*

1000 *pneumoniae* as a biocatalyst and glycerol as a carbon source. A better understanding of  
1001 glycerol metabolism in *K. pneumoniae* and improvements in its performance as a microbial  
1002 cell factory should make this organism an attractive alternative to current sugar-based  
1003 methods for the production of biochemicals and biofuels.

#### 1004 **Acknowledgement**

1005 This study was supported financially by the Advanced Biomass R&D Center (ABC) of  
1006 Global Frontier Project funded by the Korean Ministry of Science, ICT and Future planning  
1007 (ABC-2011-0031361).

## References

- Abdel-Hamid, A., Attwood, M., Guest, J., 2001. Pyruvate oxidase contributes to the aerobic growth efficiency of *Escherichia coli*. *Microbiology* 147, 1483–1498.
- Abdel-Rahman, M.A., Tashiro, Y., Sonomoto, K., 2013. Recent advances in lactic acid production by microbial fermentation processes. *Biotechnol. Adv.* 31, 877–902.
- Abdel-Rahman, M.A., Sonomoto, K., 2016. Opportunities to overcome the current limitations and challenges for efficient microbial production of optically pure lactic acid. *J. Biotech.* 236, 176–192.
- Albertí, S., Marqués, G., Camprubi, S., Merino, S., Tomás, J.M., Vivanco, F., Benedí, V.J., 1993. C1q binding and activation of the complement classical pathway by *Klebsiella pneumoniae* outer membrane proteins. *Infect. Immun.* 61, 852–860.
- Arasu, M.V., Kumar, V., Ashok, S., Song, H., Rathnasingh, C., Lee, H.J., Seung, D., Park, S., 2011. Isolation and characterization of the new *Klebsiella pneumoniae* J2B strain showing improved growth characteristics with reduced lipopolysaccharide formation. *Biotechnol. Bioprocess Eng.* 16, 1134–1143.
- Årsköld, E., Lohmeier-Vogel, E., Cao, R., Roos, S., Rådström, P., van Niel, E.W.J., 2008. Phosphoketolase Pathway Dominates in *Lactobacillus reuteri* ATCC 55730 Containing Dual Pathways for Glycolysis. *J. Bacteriol.* 190:206-212.
- Ashok, S., Raj, S.M., Rathnasingh, C., Park, S., 2011. Development of recombinant *Klebsiella pneumoniae*  $\Delta dh a T$  strain for the co-production of 3-hydroxypropionic acid and 1,3-propanediol from glycerol. *Appl. Microbiol. Biotechnol.* 90, 1253–1265.
- Ashok, S., Sankaranarayanan, M., Ko, Y., Jae, K.-E., Ainala, S.K., Kumar, V., Park, S., 2013a. Production of 3-hydroxypropionic acid from glycerol by recombinant *Klebsiella pneumoniae*  $\Delta dh a T \Delta y q h D$  which can produce vitamin B<sub>12</sub> naturally. *Biotechnol. Bioeng.* 110(2), 511–524.
- Ashok, S., Raj, S.M., Ko, Y., Sankaranarayanan, M., Zhou, S., Kumar, V., Park, S., 2013b. Effect of *puuC* overexpression and nitrate addition on glycerol metabolism and anaerobic 3-

1038 hydroxypropionic acid production in recombinant *Klebsiella pneumoniae*  $\Delta glpK\Delta dhaT$ .  
1039 Metab. Eng. 15, 10–24.

1040 Ayoub, M., Abdullah, A.Z., 2012. Critical review on the current scenario and sig- nificance  
1041 of crude glycerol resulting from biodiesel industry towards more sustainable renewable  
1042 energy industry. Renew. Sustain. Energy Rev. 16, 2671–2686.

1043 Bächler, C., Schneider, P., Bähler, P., Lustig, A., Erni, B., 2005. *Escherichia*  
1044 *coli* dihydroxyacetone kinase controls gene expression by binding to transcription factor  
1045 DhaR. EMBO J. 24, 283-293.

1046 Booth, I.R., Ferguson, G.P., Miller, S., Li, C., Gunasekera, B., Kinghorn, S., 2003. Bacterial  
1047 production of methylglyoxal: a survival strategy or death by misadventure? Biochem Soc  
1048 Trans. 31, 1406–1408.

1049 Booth, I.R., 2005. Glycerol and methylglyoxal metabolism. In: (Ed.) Bock, A., Curtiss III, R.,  
1050 Kaper, J. B., Neidhardt, F. C., Nystrom, T., Rudd, K. E., Squires, C. L., EcoSal—*Escherichia*  
1051 *coli* and *Salmonella*: cellular and molecular biology. ASM press, Washington, D.C.

1052 Branduardi, P., de Ferra, F., Longo, V., Porro, D., 2014. Microbial *n*-butanol production from  
1053 Clostridia to non-Clostridial hosts. Eng. Life Sci. 14, 16–26.

1054 Buck, M., Gallegos, M.T., Studholme, D.J., Guo, Y., Gralla, J.D., 2000. The bacterial  
1055 enhancer-dependent  $\sigma^{54}$  ( $\sigma^N$ ) transcription factor. J. Bacteriol. 182, 4129-4136.

1056 Cabelli, V.J., 1955. The tricarboxylic acid cycle in the oxidative and synthetic metabolism of  
1057 *Klebsiella pneumoniae*. J. Bacteriol. 70, 23-29.  
1058

1059 Causey, T.B., Shanmugam, K.T., Yomano, L.P., Ingram, L.O., 2004. Engineering  
1060 *Escherichia coli* for efficient conversion of glucose to pyruvate. Proc Natl Acad Sci USA  
1061 101, 2235–2240.  
1062

1063 Celińska, E., Grajek, W., 2009. Biotechnological production of 2,3-butanediol—current state  
1064 and prospects. Biotechnol. Adv. 27, 715–725.

1065 Celińska, E., 2012. *Klebsiella* spp. as a 1, 3-propanediol producer—the metabolic  
1066 engineering approach. Crit. Rev. Biotechnol. 32, 274–288.

1067 Chao, Y.-P., Liao, J.C., 1993. Alteration of growth yield by overexpression of  
 1068 phosphoenolpyruvate carboxylase and phosphoenolpyruvate carboxykinase in *Escherichia*  
 1069 *coli*. Appl. Environ. Microbiol. 59, 4261–4265.

1070

1071 Chen, Z., Liu, H.J., Liu, D.H., 2009. Regulation of 3-hydroxypropionaldehyde accumulation  
 1072 in *Klebsiella pneumoniae* by overexpression of *dhaT* and *dhaD* genes. Enzyme Microb.  
 1073 Technol. 45 (4), 305–309.

1074 Chen, Z., Liu, H., Liu, D., 2011. Metabolic pathway analysis of 1,3-propanediol production  
 1075 with a genetically modified *Klebsiella pneumoniae* by overexpressing an endogenous  
 1076 NADPH-dependent alcohol dehydrogenase. Biochemical Eng. J. 54, 151–157.

1077 Cheng, K.K., Liu, H.J., Liu, D.H., 2005. Multiple growth inhibition of *Klebsiella pneumoniae*  
 1078 in 1,3-propanediol fermentation. Biotechnol. Lett. 27, 19–22.

1079 Cheng, K.K., Zhang, J.A., Liu, D.H., Sun, Y., Liu, H.-J., Yang, M.-D., Xu, J.M., 2007. Pilot-  
 1080 scale production of 1,3- propanediol using *Klebsiella pneumoniae*. Process Biochem. 2007;  
 1081 42, 740–744.

1082 Cheng, K.K., Wu, J., Wang, G.Y., Li, W.Y., Feng, J., Zhang, J.A., 2013. Effects of pH and  
 1083 dissolved CO<sub>2</sub> level on simultaneous production of 2,3-butanediol and succinate using  
 1084 *Klebsiella pneumoniae*. Bioresour. Technol. 135:500–503.

1085 Chu, H.S., Kim, Y.S., Lee, C.M., Lee, J.H., Jung, W.S., Ahn, J.-H., Song, S.H., Choi, I.S.,  
 1086 Cho, K.M., 2015. Metabolic engineering of 3-hydroxypropionic acid biosynthesis in  
 1087 *Escherichia coli*. Biotechnol. Bioeng. 112(2), 356-364.

1088 Chun, A., Yunxiao, L., Ashok, S., Seol, E., Park, S., 2014. Elucidation of toxicity of organic  
 1089 acids inhibiting growth of *Escherichia coli* W. Biotechnol. Bioprocess Eng. 19, 858–865.

1090 Ciriminna, R., Pina C.D., Rossi, M., Pagliaro, M., 2014. Understanding the glycerol market.  
 1091 Eur. J. Lipid Sci. Technol. 116, 1432–1439.

1092 Clomburg, J.M., Gonzalez, R., 2013. Anaerobic fermentation of glycerol: a platform for  
 1093 renewable fuels and chemicals. Trends Biotechnol. 31(1), 20-28.

1094 da Silva, G.P., Mack, M., Contiero, J., 2009. Glycerol: A promising and abundant carbon  
 1095 source for industrial microbiology. Biotechnol. Adv. 27 (1), 30–39.

1096 De Mey, M., De Maeseneire, S., Soetaert, W., Vandamme, E., 2007. Minimizing acetate  
1097 formation in *E. coli* fermentations. J. Ind. Microbiol. Biotechnol. 34, 689-700.  
1098

1099 Dharmadi, Y., Murarka, A., Gonzales, R., 2006. Anaerobic fermentation of glycerol by  
1100 *Escherichia coli*: a new platform for metabolic engineering. Biotechnol. Bioeng. 94, 821–829.

1101 Dugar, D., Stephanopoulos, G., 2011. Relative potential of biosynthetic pathways for biofuels and  
1102 bio-based products. Nature Biotechnol. 29, 1074–1078.

1103 Dunlop, M.J., 2011. Engineering microbes for tolerance to next generation biofuel.  
1104 Biotechnol Biofuels 4:32.

1105 Dunlop, M.J., Dossani, Z.Y., Szmidt, H.L., Chu, H.C., Lee, T.S., Keasling, J.D., Hadi, M.Z.,  
1106 Mukhopadhyay, A., 2011. Engineering microbial biofuel tolerance and export using efflux  
1107 pumps. Mol. Syst. Biol. 7:487.

1108 Durgapal, M., Kumar, V., Yang, T.H., Lee, H.J., Seung, D., Park S., 2014. Production of 1,3-  
1109 propanediol from glycerol using the newly isolated *Klebsiella pneumoniae* J2B. Bioresour.  
1110 Technol. 159, 223–231.

1111 Durnin, G., Clomburg, J., Yeates, Z., Alvarez, P.J., Zygorakis, K., Campbell, P., Gonzalez,  
1112 R., 2008. Understanding and harnessing the microaerobic metabolism of glycerol  
1113 in *Escherichia coli*. Biotechnol. Bioeng. 103, 148–161.

1114 El Fertas-Aissani, R., Messai, Y., Alouache, S., Bakour, R., 2013. Virulence profiles and  
1115 antibiotic susceptibility patterns of *Klebsiella pneumoniae* strains isolated from different  
1116 clinical specimens. Pathol. Biol. 61(5), 209-216.

1117 Eppler, T., Boos, W., 1999. Glycerol-3-phosphate-mediated repression of *maltT* in  
1118 *Escherichia coli* does not require metabolism, depends on enzyme IIAGlc and is mediated by  
1119 cAMP levels. Mol. Microbiol. 33, 1221–1231.  
1120

1121 Eppler, T., Postma, P., Schutz, A., Volker, U., Boos, W., 2002. Glycerol-3-phosphate-  
1122 induced catabolite repression in *Escherichia coli*. J. Bacteriol., 184, 3044-3052.

1123 Fang, C.T., Chuang, Y.P., Shun, C.T., Chang, C.S., Wang, J.T., 2004. A novel virulence gene  
1124 in *Klebsiella pneumoniae* strains causing primary liver abscess and septic metastatic  
1125 complications. J. Exp. Med. 199, 697–705.

1126 Feng, X., Ding, Y., Xian, M., Xu, X., Zhang, R., Zhao, G., 2014. Production of optically pure  
 1127 D-lactate from glycerol by engineered *Klebsiella pneumoniae* strain. *Bioresour. Technol.* 172,  
 1128 269–275.

1129 Forage, R.G., Foster, M.A., 1982. Glycerol fermentation in *Klebsiella pneumoniae*: functions  
 1130 of the coenzyme B12-dependent glycerol and diol dehydratases. *J. Bacteriol.* 149, 413–419.

1131 Garcia-Alles, L.F., Siebold, C., Nyffeler, T.L., Flukiger-Bruhwiller, K., Schneider, P., Burgi,  
 1132 H.-B., Baumann, U., Erni, B., 2004. Phosphoenolpyruvate- and ATP-dependent  
 1133 dihydroxyacetone kinases: covalent substrate-binding and kinetic mechanism. *Biochemistry*  
 1134 43, 13037–13045.

1135 Gutknecht, R., Beutler, R., Garcia-Alles, L., Baumann, U., Erni, B., 2001. The dihydro-  
 1136 xyacetone kinase of *Escherichia coli* utilizes a phosphoprotein instead of ATP as phosphoryl  
 1137 donor. *EMBO J.* 20, 2480–2486.

1138 Hao, J., Wang, W., Tian, J., Li, J., Liu, D., 2008. Decrease of 3-hydroxypropionaldehyde  
 1139 accumulation in 1,3-propanediol production by over-expressing *dhaT* gene in *Klebsiella*  
 1140 *pneumoniae* TUAC01. *J. Ind. Microbiol. Biotechnol.* 35, 735–741.  
 1141

1142 Heller, K.B., Lin, E.C., Wilson, T.H., 1980. Substrate specificity and transport properties of  
 1143 the glycerol facilitator of *Escherichia coli*. *J. Bacteriol.* 144, 274–278.  
 1144

1145 Hogema, B. M., Arents, J. C., Bader, R., Eijkemans, K., Yoshida, H., Takahashi, H., Aiba,  
 1146 H., Postma, P. W., 1998. Inducer exclusion in *Escherichia coli* by non-PTS substrates: the  
 1147 role of the PEP to pyruvate ratio in determining the phosphorylation state of enzyme IIAGlc.  
 1148 *Mol. Microbiol.* 30, 487–498.  
 1149

1150 Holtman, C. K., Pawlyk, A. C., Meadow, N. D., Pettigrew, D. W., 2001. Reverse genetics of  
 1151 *Escherichia coli* glycerol kinase allosteric regulation and glucose control of glycerol  
 1152 utilization in vivo. *J. Bacteriol.* 183, 3336–3344.  
 1153

1154 Horng, Y.T., Chang, K.C., Chou, T.C., Yu, C.J., Chien, C.C., Wei, Y.H., Soo, P.C., 2010.  
 1155 Inactivation of *dhaD* and *dhaK* abolishes by-product accumulation during 1,3-propanediol  
 1156 production in *Klebsiella pneumoniae*. *J. Ind. Microbiol. Biotechnol.* 37(7), 707–716.  
 1157

1158 Huang, Y., Li, Z., Shimizu, K., Ye, Q., 2012. Simultaneous production of 3-  
1159 hydroxypropionic acid and 1,3-propanediol from glycerol by a recombinant strain of  
1160 *Klebsiella pneumoniae*. *Bioresour. Technol.* 103, 351–359.

1161

1162 Huang, Y., Li, Z., Shimizu, K., Ye, Q., 2013a. Co-production of 3-hydroxypropionic acid and  
1163 1,3-propanediol by *Klebsiella pneumoniae* expressing *aldH* under Microaerobic conditions.  
1164 *Bioresour. Technol.* 128, 505–512.

1165

1166 Huang, C.-F., Jiang, Y.-F., Guo, G.-L., Hwang, W.-S., 2013b. Method of 2,3-butanediol  
1167 production from glycerol and acid-pretreated rice straw hydrolysate by newly isolated strains:  
1168 Pre-evaluation as an integrated biorefinery process. *Bioresour. Technol.* 135, 446–453.

1169

1170 Izquierdo, L., Coderch, N., Pique, N., Bedini, E., Corsaro, M.M., Merino, S., Fresno, S.,  
1171 Tomás, J.M., Regué, M., 2003. The *Klebsiella pneumoniae wabG* gene: role in biosynthesis  
1172 of the core lipopolysaccharide and virulence. *J. Bacteriol.* 185, 7213–7221.

1173 Jarboe, L.R. 2011. YqhD: a broad-substrate range aldehyde reductase with various  
1174 applications in production of biorenewable fuels and chemicals. *Appl. Microbiol.*  
1175 *Biotechnol.* 89, 249–257.

1176 Jarboe, L.R., Liu, P., Royce, L.A., 2011. Engineering inhibitor tolerance for the production of  
1177 biorenewable fuels and chemicals. *Curr. Opin. Chem. Eng.* 1, 38–42.

1178 Jeong, J.-Y., Kim, Y.-J., Cho, N., Shin, D., Nam, T.-W., Ryu, S., Seok YJ., 2004. Expression  
1179 of *ptsG* encoding the major glucose transporter is regulated by *arcA* in *Escherichia coli*. *J.*  
1180 *Biol. Chem.* 279, 38513–38518.

1181

1182 Ji, X.-J., Huang, H., Ouyang, P.-K., 2011. Microbial 2,3-butanediol production: A state-of-  
1183 the-art review. *Biotechnol. Adv.* 29, 351–364.

1184 Jiang, X., Meng, X., Xian, M., 2009. Biosynthetic pathways for 3-hydroxypropionic acid  
1185 production. *Appl. Microbiol. Biotechnol.* 82, 995–1003.

1186 Jung, S.G., Jang, J.H., Kim, A.Y., Lim, M.C., Kim, B., Lee, J., Kim, Y.-R., 2013. Removal of  
1187 pathogenic factors from 2,3-butanediol producing *Klebsiella* species by inactivating virulence  
1188 related *wabG* gene. *Appl. Microbiol. Biotechnol.* 97, 1997–2007.



1189

1190 Jung, W.S., Kang, J.H., Chu, H.S., Choi, I.S., Cho, K.M., 2014. Elevated production of 3-

1191 hydroxypropionic acid by metabolic engineering of the glycerol metabolism in *Escherichia*

1192 *coli*. Metab. Eng. 23, 116–122.

1193 Jung, H.-W., Jung, M.-W., Oh, M.-K., 2015. Metabolic engineering of *Klebsiella*

1194 *pneumoniae* for the production of cis,cis-muconic acid. Appl. Microbiol. Biotechnol. 99,

1195 5217–5225.

1196 Katryniok, B., Paul, S., F. Dumeignil, F., 2013. Recent Developments in the Field of Catalytic

1197 Dehydration of Glycerol to Acrolein. ACS Catal. 3, 1819-1834.

1198 Kim, Y., Ingram, L.O., Shanmugam, K.T., 2008. Dihydrolipoamide dehydrogenase mutation

1199 alters the NADH sensitivity of pyruvate dehydrogenase complex of *Escherichia coli* K-12. J.

1200 Bacteriol. 190, 3851–3858.

1201

1202 Kim, K., Kim, S.K., Park, Y.C., Seo, J.H., 2014a. Enhanced production of 3-

1203 hydroxypropionic acid from glycerol by modulation of glycerol metabolism in

1204 recombinant *Escherichia coli*. Bioresour. Technol. 156, 170-175.

1205 Kim, B., Lee, S., Jeong, D., Yang, J., Oh, M.-K., Lee, J., 2014b. Redistribution of Carbon

1206 Flux toward 2,3-Butanediol Production in *Klebsiella pneumoniae* by Metabolic Engineering.

1207 PLoS One 9(10), e105322.

1208 Ko, Y., Ashok, S., Zhou, S., Kumar, V., Park, S., 2012. Aldehyde dehydrogenase activity is

1209 important to the production of 3-hydroxypropionic acid from glycerol by recombinant

1210 *Klebsiella pneumoniae*. Process Biochem. 47, 1135–1143.

1211

1212 Ko, Y., Ashok, S., Seol, E., Ainala, S., Park, S., 2015. Deletion of putative oxidoreductases

1213 from *Klebsiella pneumoniae* J2B could reduce 1,3-propanediol during the production of 3-

1214 hydroxypropionic acid from glycerol. Biotechnol. Bioprocess Eng. 20, 834–843.

1215

1216 Ko, Y., Seol, E., Sekar, B.S., Kwon, S., Lee, J., Park, S., 2017. Metabolic engineering of

1217 *Klebsiella pneumoniae* J2B for co-production of 3- hydroxypropionic acid and 1,3-

1218 propanediol from glycerol: Reduction of acetate and other by-products. Bioresour. Technol.

1219 244, 1096–1103.

1220 Kong P.S., Aroua M.K., Daud, W.M.A.D. 2016. Conversion of crude and pure glycerol into  
1221 derivatives: A feasibility evaluation. *Renew. Sustain. Energy Rev.* 63, 533-555.

1222 Kraus, G.A., 2008. Synthetic methods for the preparation of 1, 3-propanediol. *Clean* 36 (8),  
1223 648–651.

1224 Kumar, V., Mugesh, S., Jae, K., Durgapal, M., Ashok, S., Ko, Y., Sarkar, R., Park, S., 2012.  
1225 Co-production of 3-hydroxypropionic acid and 1,3-propanediol from glycerol using resting  
1226 cells of *Klebsiella pneumoniae* J2B strain with overexpression of KGSADH. *Appl. Microbiol.*  
1227 *Biotechnol.* 96, 373–383.

1228 Kumar, V., Ashok, S., Park, S., 2013a. Recent advances in biological production of 3-  
1229 hydroxypropionic acid. *Biotechnol. Adv.* 31(6), 945–961

1230 Kumar, V., Mugesh, S., Durgapal, M., Zhou, S., Ko, Y., Ashok, S., Sarkar, R., Park, S.,  
1231 2013b. Simultaneous production of 3-hydroxypropionic acid and 1,3-propanediol from  
1232 glycerol using resting cell system of D-lactic acid deficient mutant of *Klebsiella pneumonia*  
1233 with overexpression of KGSADH. *Bioresour. Technol.* 135, 555-563.

1234 Kumar, V., Durgapal, M., Sankaranarayanan, M., Somasundar, A., Rathnasingh, C., Song,  
1235 H., Seung, D., Park, S., 2016. Effects of mutation of 2,3-butanediol formation pathway on  
1236 glycerol metabolism and 1,3-propanediol production by *Klebsiella pneumoniae* J2B.  
1237 *Bioresour. Technol.* 214, 432–440.

1238 Lama, S., Ro, S. M., Seol, E., Sekar, B.S., Ainala, S.K., Thangappan, J., Song, H., Seung,  
1239 D., Park, S., 2015. Characterization of 1,3-propanediol oxidoreductase (DhaT)  
1240 from *Klebsiella pneumoniae* J2B. *Biotechnol. Bioproc. Eng.* 20: 971–979.

1241 Lama, S. , Seol, E., Park, S., 2017. Metabolic engineering of *Klebsiella pneumoniae* J2B for  
1242 the production of 1,3-propanediol from glucose. *Bioresour. Technol.* doi:  
1243 10.1016/j.biortech.2017.05.052.

1244 Lee, C.S., Aroua M.K. Aroua, Daud, W.M.A.W., Cognet, P., Pérès-Lucchese, Y., Fabre, P.-  
1245 L., Reynes, O., Latapie, L., 2015. A review: Conversion of bioglycerol into 1,3-propanediol  
1246 via biological and chemical method. *Renew. Sustain. Energy Rev.* 42, 963-972.

1247 Li, Y., Wang, X., Ge, X. & Tian, P., 2016. High production of 3-hydroxypropionic acid  
1248 in *Klebsiella pneumoniae* by systematic optimization of glycerol metabolism. *Sci. Rep.* 6.  
1249 26932.

1250 Liao, D.-I., Reiss, L., Turner, Jr. I., Dotson, G., 2003. Structure of Glycerol Dehydratase  
1251 Reactivase: A New Type of Molecular Chaperone. *Structure* 11, 109–119.

1252 Lim, H.G., Noh, M.H., Jeong, J.H., Park, S., Jung, G.Y., 2016. Optimum rebalancing of the  
1253 3-hydroxypropionic acid production pathway from glycerol in *Escherichia coli*. *ACS Synth.*  
1254 *Biol.* **5** (11), 1247–1255.

1255

1256 Lin, E. C., 1976. Glycerol dissimilation and its regulation in bacteria. *Annu. Rev. Microbiol.*  
1257 **30**, 535-578.

1258

1259 Lin, E. C. C., 1987. *Escherichia coli* and *Salmonella typhimurium*: Cellular and Molecular  
1260 Biology. In: Neidhardt, F. C. (Ed.), Vol. 1. American Society for Microbiology, Washington,  
1261 D. C, pp. 244-284.

1262

1263 Lin, H., Castro, N.M., Bennett, G.N., San, K.-Y., 2006. Acetyl-CoA synthetase  
1264 overexpression in *Escherichia coli* demonstrates more efficient acetate assimilation and lower  
1265 acetate accumulation: a potential tool in metabolic engineering. *Appl. Microbiol. Biotechnol.*  
1266 **71**(6), 870–874.

1267

1268 Lin, T.-L., Yang, F.-L., Yang, A.-S., Peng, H.-P., Li, T.-L., Tsai, M.-D., Wu, S.-H., Wang, J.-  
1269 T., 2012. Amino acid substitutions of *MagA* in *Klebsiella pneumoniae* affect the biosynthesis  
1270 of the capsular polysaccharide. *PLoS One* **7**:e46783.

1271 Liu, H.J., Zhang, D.J., Xu, Y.H., Mu, Y., Sun, Y.Q., Xiu, Z.L., 2007. Microbial production of  
1272 1, 3-propanediol from glycerol by *Klebsiella pneumoniae* under micro-aerobic conditions up  
1273 to a pilot scale. *Biotechnol. Lett.* **29**:1281–1285.

1274

1275 Liu, Y., Gallo, A.A., Bajpai, R.K., Chistoserdov, A., Nelson, A., Segura, L., Xu W., 2010.  
1276 The diversity and molecular modelling analysis of B-12 and B-12-independent glycerol  
1277 dehydratases. *Int. J. Bioinform. Res. Appl.* **6**(5), 484–507.

1278

1279 Luo, L.H., Seo, J.W., Baek, J.O., Oh, B.R., Heo, S.Y., Hong, W.K., Kim, D.H., Kim, C.H.,  
1280 2011. Identification and characterization of the propanediol utilization protein PduP of  
1281 *Lactobacillus reuteri* for 3-hydroxypropionic acid production from glycerol. Appl. Microbiol.  
1282 Biotechnol. 89, 697–703.

1283 Luo, L.H., Kim, C.H., Heo, S.Y., Oh, B.R., Hong, W.K., Kim, S., Kim, D.H., Seo, J.W.,  
1284 2012. Production of 3-hydroxypropionic acid through propionaldehyde dehydrogenase PduP  
1285 mediated biosynthetic pathway in *Klebsiella pneumoniae*. Bioresour. Technol. 103, 1–6.

1286 Lütke-Eversloh, T., Bahl, H., 2011. Metabolic engineering of *Clostridium acetobutylicum*:  
1287 Recent advances to improve butanol production. Curr. Opin. Biotechnol. 22, 634–647.

1288 Ma, B.B., Xu, X.L., Zhang, G.L., Wang, L.W., Wu, M., Li, C., 2009. Microbial production of  
1289 1,3-propanediol by *Klebsiella pneumoniae* XJPD-Li under different aeration strategies. Appl.  
1290 Biochem. Biotechnol. 152(1), 127–134.

1291 Ma, C., Zhang, L., Dai, J., Xiu, Z., 2010. Relaxing the coenzyme specificity of 1,3-  
1292 propanediol oxidoreductase from *Klebsiella pneumoniae* by rational design. J. Biotechnol.  
1293 146(4), 173–178.

1294 Maervoet, V.E.T., Mey, M.D., Beauprez, J., Maeseneire, S.D., Soetaert, W.K., 2011.  
1295 Enhancing the microbial conversion of glycerol to 1,3-propanediol using metabolic  
1296 engineering. Org. Process Res. Dev. 15, 189–202.

1297 MarketsANDMarkets, 2012. Global 1,3-propanediol (1,3-PDO) market worth \$560 million  
1298 by 2019.

1299 Martins-Pinheiro, M., Lima, W.C., Asif, H., Oller, C.A., Menck, C.F.M., 2016. Evolutionary  
1300 and Functional Relationships of the dha Regulon by Genomic Context Analysis. PLOS  
1301 ONE 11, e0150772.

1302 Mattam, A.J., James, M., Clomburg, J.M., Gonzalez, R., Yazdani, S.S., 2013. Fermentation  
1303 of glycerol and production of valuable chemical and biofuel molecules. Biotechnol. Lett. 35,  
1304 831–842.

1305 Mazumdar, S., Clomburg, J.M., Gonzalez, R., 2010. *Escherichia coli* strains engineered for  
1306 homofermentative production of D-lactic acid from glycerol. Appl. Environ. Microbiol. 76,  
1307 4327–4336.

1308 Mazumdar, S., Blankschien, M.D., Clomburg, J.M., Gonzalez R., 2013. Efficient synthesis of  
1309 L-lactic acid from glycerol by metabolically engineered *Escherichia coli*. Microb. Cell Fact.  
1310 12:7.

1311 Meyer, M., Dimroth, P., Bott, M., 2001. Catabolite repression of the citrate fermentation  
1312 genes in *Klebsiella pneumoniae*: evidence for involvement of the cyclic AMP receptor  
1313 protein. J. Bacteriol. 183, 5248–5256.

1314

1315 Organisation for Economic Co-operation and Development (OECD), Food and Agriculture  
1316 Organization of the United Nations (FAO), Biofuels, in OECD-FAO Agricultural Outlook  
1317 2011–2020, OECD, Paris, 2011, pp. 77–94.

1318

1319 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Joe, M., Park, D.H., Kim, C.H., 2011.  
1320 Efficient production of ethanol from crude glycerol by a *Klebsiella pneumonia* mutant strain.  
1321 Bioresour. Technol. 102, 3918-3922.

1322 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Kim, S.H., Park, D.H., Kim, C.H.,  
1323 2012a. Optimization of culture conditions for 1,3-propanediol production from glycerol using  
1324 a mutant strain of *Klebsiella pneumoniae*. Appl. Biochem. Biotechnol. 166, 127–137.

1325 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Kim, S., Kwon, O., Sohn, J.-H., Joe,  
1326 M.-H., Park, D.-H., Kim, C.H., 2012b. Enhancement of ethanol production from glycerol in a  
1327 *Klebsiella pneumoniae* mutant strain by the inactivation of lactate dehydrogenase. Process  
1328 Biochem. 47, 156–159.

1329 Oh, B.R., Heo, S.Y., Lee, S.M., Hong, W.K., Park, J.M., Jung, Y.R., Kim, D.H., Sohn, J.H.,  
1330 Seo, J.W., Kim, C.H., 2014. Production of 2- butanol from crude glycerol by a genetically-  
1331 engineered *Klebsiella pneumoniae* strain. Biotechnol. Lett. 36 (1), 57–62.

1332 Park, J.M., Rathnasingh, C., Song H., 2017. Metabolic engineering of *Klebsiella pneumoniae*  
1333 based on in silico analysis and its pilot-scale application for 1,3-propanediol and  
1334 2,3-butanediol co-production Jong. J. Ind. Microbiol. Biotechnol. 44, 431–441.

1335 Petrov, K., Petrova, P., 2009. High production of 2,3-butanediol from glycerol by *Klebsiella*  
1336 *pneumoniae* G31. Appl. Microbiol. Biotechnol. 84, 659–665.

1337 Petrov, K., Petrova, P., 2010. Enhanced production of 2,3-butanediol from glycerol by forced  
1338 pH fluctuations. *Appl. Microbiol. Biotechnol.* 87, 943–949.

1339 Pina, C.D., Falletta, E., Rossi, M., 2011. A green approach to chemical building blocks. The  
1340 case of 3-hydroxypropanoic acid. *Green Chem.* 13, 1624–1632.

1341 Podschun, R., Ullmann, U., 1998. *Klebsiella* spp. as nosocomial pathogens: epidemiology,  
1342 taxonomy, typing methods, and pathogenicity factors. *Clin. Microbiol. Rev.* 11, 589–603.

1343 Quispe C. A. G., Coronado, C.J.R., Carvalho, Jr.J.A., Glycerol: Production, consumption,  
1344 prices, characterization and new trends in combustion. *Renew. Sustain. Energy Rev.* 27, 475–  
1345 493.

1346 Raynaud C., Sarcabal P., Meynial-Salles I., Croux C., Soucaille P. 2003. Molecular  
1347 characterization of the 1,3-propanediol (1,3-PD) operon of *Clostridium butyricum*. *Proc. Natl.*  
1348 *Acad. Sci. USA* 100, 5010–5015.

1349 Raynaud, C., Lee, J., Sarcabal, P., Croux, C., Meynial-Salles, I., Soucaille, P., 2011.  
1350 Molecular characterization of the glycerol-oxidative pathway of *Clostridium butyricum* VPI  
1351 1718. *J. Bacteriol.* 193, 3127–3134.

1352 Reizer, J., Reizer, A., Saier, M.H. Jr., 1993. The MIP family of integral membrane channel  
1353 proteins: sequence comparisons, evolutionary relationships, reconstructed pathway of  
1354 evolution, and proposed functional differentiation of the two repeated halves of the proteins.  
1355 *Crit Rev Biochem. Mol. Biol.* 28:235-257.

1356 Richey, D.P., Lin, E.C., 1972. Importance of facilitated diffusion for effective utilization of  
1357 glycerol by *Escherichia coli*. *J. Bacteriol.* 112, 784-790.

1358 Rossi, D.M., de Souza, E.A., Ayub, M.A., 2013. Biodiesel residual glycerol metabolism by  
1359 *Klebsiella pneumoniae*: pool of metabolites under anaerobiosis and oxygen limitation as a  
1360 function of feeding rates. *Appl. Biochem. Biotechnol.* 169, 1952–1964.

1361 Sanchez, A.M., Bennett, G.N., San, K.Y., 2005. Efficient succinic acid production from  
1362 glucose through overexpression of pyruvate carboxylase in an *Escherichia coli* alcohol  
1363 dehydrogenase and lactate dehydrogenase mutant. *Biotechnol. Prog.* 21, 358–365.

1364

1365 Saxena, R.K., Anand, P., Saran, S., Isar, J., 2009. Microbial production of 1,3-propanediol:  
 1366 Recent developments and emerging opportunities. *Biotechnol. Adv.* 27, 895–913.  
 1367

1368 Seo, M.Y., Seo, J.W., Heo, S.Y., Baek, J.O., Rairakhwada, D., Oh, B.R., Seo, P.S., Choi,  
 1369 M.H., Kim, C.H., 2009. Elimination of by-product formation during production of 1,3-  
 1370 propanediol in *Klebsiella pneumoniae* by inactivation of glycerol oxidative pathway. *Appl.*  
 1371 *Microbiol. Biotechnol.* 84, 527–534.  
 1372

1373 Shrivastav, A., Lee, J., Kim, H.-Y., Kim, Y.-R., 2013. Recent insights in the removal  
 1374 of *Klebsiella* pathogenicity factors for the industrial production of 2,3-butanediol. *J.*  
 1375 *Microbiol. Biotechnol.* 23, 885–896.  
 1376

1377 Stieb, M., Schink, B., 1984. A new 3-hydroxybutyrate fermenting anaerobe, *Ilyobacter*  
 1378 *polytropus*, gen nov, sp nov, possessing various fermentation pathways. *Arch Microbiol.*  
 1379 140(2), 139–146.  
 1380

1381 Sun, J., van den Heuvel, J., Soucaille, P., Qu, Y., Zeng, A., 2003. Comparative genomic  
 1382 analysis of *dha* regulon and related genes for anaerobic glycerol metabolism in bacteria.  
 1383 *Biotechnol. Progr.* 19, 263–272.  
 1384

1385 Sun, Y.Q., Qi, W.T., Teng, H., Xiu, Z.L., Zeng, A.P., 2008. Mathematical modelling of  
 1386 glycerol fermentation by *Klebsiella pneumoniae*: Concerning enzyme-catalytic reductive  
 1387 pathway and transport of glycerol and 1,3-propanediol across cell membrane. *Biochem. Eng.*  
 1388 *J.* 38, 22–32.  
 1389

1390 Sun, D., Yamada, Y., Sato, S., Uedab, W., 2017. Glycerol as a potential renewable raw  
 1391 material for acrylic acid production. *Green Chem.* DOI: 10.1039/C7GC00358G.  
 1392

1393 Taylor, B.L., Zhulin, I.B., 1999. PAS domains: internal sensors of oxygen, redox potential,  
 1394 and light. *Microbiol. Mol. Biol. Rev.* 63, 479-506.  
 1395

1396 Trinh, C.T., Sreenc, F., 2009. Metabolic Engineering of *Escherichia coli* for efficient  
 1397 conversion of glycerol to ethanol. *Appl. Environ. Microbiol.* 75, 6696–6705.  
 1398

1399 Wang, W., Sun, J., Hartlep, M., Deckwer, W.D., Zeng, A.P., 2003. Combined use of  
1400 proteomic analysis and enzyme activity assays for metabolic pathway analysis of glycerol  
1401 fermentation by *Klebsiella pneumoniae*. Biotechnol. Bioeng. 83, 525–536.  
1402

1403 Wang, Y., Tao, F., Xu P., 2014. Glycerol dehydrogenase plays a dual role in glycerol  
1404 metabolism and 2,3-butanediol formation in *Klebsiella pneumoniae*. J. Biol. Chem. 113, 525–  
1405 535.  
1406

1407 Wang, M., Hu, L., Fan, L., Tan, T., 2015. Enhanced 1-Butanol Production in Engineered  
1408 *Klebsiella pneumoniae* by NADH Regeneration. Energ. Fuel 29, 1823–1829.  
1409

1410 Warnecke, T., Gill, R.T., 2005. Organic acid toxicity, tolerance, and production in  
1411 *Escherichia coli* biorefining applications. Microb. Cell Fact. 4, 425.  
1412

1413 Wei, D., Xu, J., Sun, J., Shi, J., Hao, J., 2013. 2-Ketogluconic acid production by *Klebsiella*  
1414 *pneumoniae* CGMCC 1.6366. J. Ind. Microbiol. Biotechnol. 40, 561–570.

1415 Wei., D., Wang, M., Jiang, B., Shi, J., Hao, J., 2014. Role of dihydroxyacetone kinases I and  
1416 II in the *dha* regulon of *Klebsiella pneumoniae*. J. Biotechnol. 177, 13-19.

1417 Weissenborn, D. L., Wittekindt, N., Larson, T.J., 1992. Structure and regulation of the *glpFK*  
1418 operon encoding glycerol diffusion facilitator and glycerol kinase of *Escherichia coli* K-12. J.  
1419 Biol. Chem. 267, 6122–6131.  
1420

1421 Werpy, T, Petersen, G., 2004. Top value added chemicals from biomass, vol 1: results of  
1422 screening for potential candidates from sugars and synthesis gas. US Department of Energy  
1423 [<http://www.osti.gov/bridge>].

1424 Wu, Z., Wang, Z., Wang, G., Tan, T., 2013. Improved 1,3-propanediol production by  
1425 engineering the 2,3-butanediol and formic acid pathways in integrative recombinant  
1426 *Klebsiella pneumoniae*. J. Biotechnol. 168, 194–200.

1427 Xu, X.L., Zhang, G.L., Wang, L.W., Ma, B.B., Li C., 2009a. Quantitative analysis on  
1428 inactivation and reactivation of recombinant glycerol dehydratase from *Klebsiella*  
1429 *pneumoniae* XJPD-Li. J Mol. Catal. B Enzym. 56, 108–114.



1430 Xu, Y.Z., Guo, N.N., Zheng, Z.M., Ou, X.J., Liu, H.J., Liu, D.H., 2009b. Metabolism in 1,3-  
1431 Propanediol Fed-Batch Fermentation by a D-Lactate Deficient Mutant of *Klebsiella*  
1432 *pneumoniae*. Biotechnol. Bioeng. 104, 965–972.

1433 Yang, B., Larson, T.J., 1998. Multiple promoters are responsible for transcription of the  
1434 *glpEGR* operon of *Escherichia coli*. Biochim. Biophys. Acta 1396, 114–126.

1435 Yazdani, S.S., Gonzalez, R., 2007. Anaerobic fermentation of glycerol: a path to economic  
1436 viability for the biofuels industry. Curr. Opin. Biotechnol. 18(3), 213–219.

1437 Zeng, G., Ye, S., Larson, T.J., 1996. Repressor for the sn-glycerol 3-phosphate regulon of  
1438 *Escherichia coli* K-12: primary structure and identification of the DNA-binding domain. J.  
1439 Bacteriol. 178, 7080–7089.

1440 Zhang, Y., Li, Y., Du, C., Liu, M., Cao, Z., 2006. Inactivation of aldehyde dehydrogenase: a  
1441 key factor for engineering 1,3-propanediol production by *Klebsiella pneumoniae*. Metab. Eng.  
1442 8, 578–586.

1443 Zhang, Q.R., Xiu, Z.L., 2009. Metabolic pathway analysis of glycerol metabolism in  
1444 *Klebsilla pneumonia* incorporating oxygen regulatory system. Biotechnol. Prog. 25, 103–115.

1445 Zhao, L., Zheng, Y., Ma, X., Wei, D., 2009. Effects of over-expression of glycerol  
1446 dehydrogenase and 1,3-propanediol oxidoreductase on bioconversion of glycerol into 1,3-  
1447 propandediol by *Klebsiella pneumoniae* under micro-aerobic conditions. Bioprocess Biosyst.  
1448 En. 32(3), 313–320.

1449 Zheng, P., Wereath, K., Sun, J., Van den Heuvel, J., Zeng AP., 2006. Overexpression of  
1450 genes of the *dha* regulon and its effects on cell growth, glycerol fermentation to 1,3-  
1451 propanediol and plasmid stability in *Klebsiella pneumoniae*. Process Biochem. 41, 2160–  
1452 2169.

1453 Zheng, Z.M., Cheng, K.K., Hu, Q.L., Liu, H.J., Guo, N.N., Liu, D., 2008. Effect of culture  
1454 conditions on 3-hydroxypropionaldehyde detoxification in 1,3-propanediol fermentation by  
1455 *Klebsiella pneumoniae*. Biochem. Eng. J. 39(2), 305–310.

1456 Zhu, J.G., Li, S., Ji, X.J., Huang, H., Hu, N., 2009. Enhanced 1,3-propanediol production in  
1457 recombinant *Klebsiella pneumoniae* carrying the gene *yqhD* encoding 1,3-propanediol  
1458 oxidoreductase isoenzyme. W. J. Microbiol. Biotechnol. 25, 1217–1223.

Zhuge, B., Zhang, C., Fang, H., Zhuge, J., Permaul, K., 2010. Expression of 1,3-propanediol oxidoreductase and its isoenzyme in *Klebsiella pneumoniae* for bioconversion of glycerol into 1,3-propanediol. Appl. Microbiol. Biotechnol. 87(6), 2177–2184.

## Figure captions

**Figure 1:** Transformation of glycerol into a number of valuable chemicals. The solid line represent single step while broken one indicate multiple steps.

**Figure 2:** Oxidative metabolism of glycerol in *K. pneumoniae* (Ashok et al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrows. Solid lines indicate single steps; broken lines indicate multiple steps.

**Figure 3:** Metabolic pathways for the production of PDO and 3-HP from glycerol (Ashok et al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrows.

**Figure 4 (A):** Arrangement of genes in the *dha* regulon of *K. pneumoniae*. 1, *dhaK* (ATP-dependent dihydroxyacetone kinase); 2, *dhaK3/dhaM* (large subunit of ATP-dependent dihydroxyacetone kinase); 3, *dhaK2/dhaL* (small subunit of ATP-dependent dihydroxyacetone kinase); 4, *dhaK1/dhaK* (medium subunits of ATP-dependent dihydroxyacetone kinase); 5, *orfW* (hypothetical protein); 6, *dhaD* (glycerol dehydrogenase); 7, *dhaR* (transcription regulator); 8, *orfW/CdAT* [cob(I)yrinic acid a,c-diamide adenosyltransferase]; 9, *orfX/dhaG* (small subunit of reactivating factor of glycerol dehydratase); 10, *dhaT* (1,3-propanediol oxidoreductase); 11, *orfY* (hypothetical protein); 12, *dhaB/B1* (large subunits of glycerol dehydratase); 13, *dhaC/B2* (medium subunit of glycerol dehydratase); 14, *dhaE/B3* (small subunit of glycerol dehydratase); 15, *orfZ/dhaF* (large subunit of reactivating factor of glycerol dehydratase); 16, *glpF* (glycerol uptake facilitator); 17, 22- *hdeB*; 18, *hdeD* (*hdeBD*-genes related to global regulation); 19, *yfdX*; 20, *kvgS*; and 21, *kvgA* (*kvgS* and *kvgA* encode for a two-component signal transduction system) (Celińska, 2012; Sun et al., 2003; Wei et al., 2014). **(B):** Glycerol dehydratase gene arrangements and reactivation factors in different organisms (Martins-Pinheiro et al., 2016).

**Figure 5:** Metabolic pathway for the synthesis of BDO from glycerol (Ashok et al., 2011; Celińska and Grajek, 2009; Ji et al., 2011). Gene names are shown in italics on arrows. Solid lines indicate single steps; broken lines indicate multiple steps.

1488 **Figure 6:** Biochemical routes for ethanol and acetic acid production from glycerol (Ashok et  
1489 al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrow. Solid lines  
1490 indicate single steps; broken lines indicate multiple steps.

1491 **Figure 7:** Lactic acid production from glycerol via methylglyoxal and pyruvate routes  
1492 (Ashok et al., 2011; Mazumdar et al., 2010; 2013). Gene names are shown in italics on arrow.  
1493 Solid lines indicate single steps; broken lines indicate multiple steps.

1494 **Fig. S1** Effect of deletion of glycerol transporters, *glpF1* and *glpF2*, on glycerol consumption  
1495 and PDO production in *K. pneumoniae* J2B3.

1496

1    **Potential and Limitations of *Klebsiella pneumoniae* as a Microbial Cell Factory Utilizing**  
2    **Glycerol as the Carbon Source**

3  
4    **Vinod Kumar<sup>a</sup>, Sunghoon Park<sup>b\*</sup>**

5  
6    **<sup>a</sup> Bioenergy and Resource Management Centre, School of Water, Energy and**  
7    **Environment, Cranfield University, Cranfield MK43 0AL, United Kingdom**

8    **<sup>b</sup> School of Energy and Chemical Engineering, UNIST, 50 UNIST-gil, Ulsan 44919,**  
9    **Republic of Korea**

10  
11    **\*Corresponding author**

12    **School of Energy and Chemical Engineering, UNIST, 50 UNIST-gil, Ulsan 44919,**  
13    **Republic of Korea**

14    **Tel.: +82 52 217 2565; Fax: +82 52 217 2309; E-mail address: parksh@unist.ac.kr**  
15

16  
17

## Abstract

*Klebsiella pneumoniae* is a Gram-negative facultative anaerobe that metabolizes glycerol efficiently under both aerobic and anaerobic conditions. This microbe is considered an outstanding biocatalyst for transforming glycerol into a variety of value-added products. Crude glycerol is a cheap carbon source and can be converted by *K. pneumoniae* into useful compounds such as lactic acid, 3-hydroxypropionic acid, ethanol, 1,3-propanediol, 2,3-butanediol, and succinic acid. This review summarizes glycerol metabolism in *K. pneumoniae* and its potential as a microbial cell factory for the production of commercially important acids and alcohols. Although many challenges remain, *K. pneumoniae* is a promising workhorse when glycerol is used as the carbon source.

**Key words:** *Klebsiella pneumoniae*; Glycerol; 3-Hydroxypropionic acid; 1,3-Propanediol; 2,3-Butanediol; Lactic acid; Ethanol

## 1. Introduction

Crude glycerol is produced in large quantities as a byproduct of the biodiesel industry. Global biodiesel production was 22.7 million tons in 2012 and is forecasted to reach 36.9 million tons by 2020. As a result, worldwide production of glycerol climbed from 1 million tons in 2000 to 3 million tons in 2011 and is expected to reach 6 million tons by 2025 (Ciriminna et al., 2014; OECD, 2011; Katryniok et al., 2013; Sun et al. 2017). A significant amount of glycerol is also generated during bioethanol fermentation by yeast. The liquid fraction of the stillage obtained from ethanol separation contains up to 2% glycerol. Another source of glycerol is industrial waste generated from vegetable oils and animal fats. For example, the concentration of glycerol in waste streams from the oleochemical industry is 55%–90% (da Silva et al., 2009; Yazdani and Gonzalez, 2007). The increased availability of glycerol has caused a substantial reduction in its cost. According to a recent report, the price of crude glycerol is \$0.24/kg and that of pure USP grade glycerol is \$0.9/kg (Kong et al., 2016). Extensive research has been conducted to investigate ways to utilize this surplus crude glycerol. The annual number of research articles addressing the use of glycerol has increased to >7000, doubling in number from the year 2000 to 2007. Numerous chemical conversions to valuable products have been reported (Ciriminna et al., 2014).

Glycerol can be used for the microbial production of 1,3-propanediol (PDO), ethanol, 2,3-butanediol (BDO), butanol, lactic acid, 3-hydroxypropionic acid (3-HP), and succinic acid (Fig. 1) (Clomberg and Gonzalez, 2013; Mattam et al., 2013; Yazdani and Gonzalez, 2007). Because of its reduced nature, glycerol generates twice the number of reducing equivalents per carbon than do traditional carbohydrates (glucose, xylose, and sucrose) when converted into the glycolytic intermediates phosphoenolpyruvate (PEP) or pyruvate (Dharmadi et al., 2006). As a consequence, glycerol gives higher yields of reduced metabolites (e.g., succinate, ethanol, PDO, and butanol) than does glucose (Yazdani and

Gonzalez, 2007). Table 1 compares the maximum theoretical yield of various metabolites, ATP, and redox balances between glycerol and glucose.

Despite its great potential as a carbon source, glycerol is not efficiently metabolized by many microorganisms under anaerobic or oxygen-limited conditions. The ability of *K. pneumoniae* to assimilate glycerol under oxygen-limited conditions is outstanding and has been studied extensively. Several review articles addressing this issue have also been published (Clomberg and Gonzalez, 2013; Mattam et al., 2013; Yazdani and Gonzalez, 2007), but the biotechnological potential and physiological aspects of this strain have not been fully elucidated. This review explores and discusses glycerol metabolism in *K. pneumoniae* and the production of value-added chemicals from glycerol. The potential and challenges of using *K. pneumoniae* as a microbial cell factory are also discussed.

## **2. Glycerol metabolism in *K. pneumoniae***

Many microorganisms can metabolize glycerol in the presence of external electron acceptors (respiratory metabolism), but relatively few under non-respiratory conditions (without exogenous electron acceptors). Among the latter, genera including *Citrobacter*, *Enterobacter*, *Bacillus*, *Propionibacterium*, *Anaerobiospirillum*, *Klebsiella*, *Clostridium*, and *Lactobacillus* are well documented (da Silva et al., 2009; Yazdani and Gonzalez, 2007). Some *Lactobacillus* species including *L. reuteri*, *L. brevis*, and *L. buchneri* convert glycerol to PDO but only in the presence of a major carbon source for growth. Glycerol does not support cell growth, limiting its biotechnological applications. In contrast, *Clostridium* sp. grow well on glycerol as the sole carbon source. However, the use of *Clostridium* is limited by its slow growth and requirement of strict anaerobic conditions. The lack of a convenient genetic tool box for large-scale metabolic engineering is another serious drawback of *Clostridium* (Branduardi et al., 2014; Lütke-Eversloh and Bahl, 2011).

From a biotechnological standpoint, *Klebsiella sp.*, especially *K. pneumoniae*, have many advantages over the aforementioned microbes. They grow on glycerol as the sole carbon source under both aerobic and anaerobic conditions (Arasu et al., 2011; Kumar et al., 2012). Their growth is fast, typically growing at a rate  $>0.9/h$ , even on glycerol minimal medium (Arasu et al., 2011). They naturally produce coenzyme B<sub>12</sub>, an essential cofactor for the synthesis of the commercially important chemicals PDO and 3-HP (Huang et al., 2013a; Oh et al., 2012a). Furthermore, their evolutionary and biochemical proximity to *E. coli* allows for the application of most of gene manipulation methods developed for *E. coli* without much modification (Celińska, 2012; Kumar et al., 2013a).

## **2.1 Overview of glycerol metabolism in *K. pneumoniae***

In *K. pneumoniae*, glycerol is metabolized oxidatively and/or reductively (Ashok et al., 2011, Kumar et al., 2013a). The oxidative pathway provides energy and carbon constituents for the synthesis of cell biomass, while the reductive pathway reduces glycerol and regenerates NAD<sup>+</sup>, enabling the oxidative assimilation of glycerol under anaerobic conditions.

Glycerol oxidation proceeds in either a respiratory or a fermentative manner. Respiration requires an exogenous electron acceptor such as oxygen, nitrate, or fumarate (Ashok et al., 2013a,b) and is carried out by enzymes encoded by the *glp* regulon. During respiration, glycerol is first phosphorylated by glycerol kinase (*glpK*) to yield *sn*-glycerol-3-phosphate using ATP or phosphoenolpyruvate (PEP) as a phosphate donor. Next, *sn*-glycerol-3-phosphate is oxidized to dihydroxyacetone phosphate (DHAP) with electron transfer to a quinone molecule, which is linked to the electron transport chain (ETC). This transformation is catalyzed by glycerol-3-phosphate dehydrogenase, which is expressed under both aerobic (*glpD*) and anaerobic (*glpABC*) conditions. In contrast, fermentation of glycerol is conducted without a supply of external electron acceptors, and the reactions are



catalyzed by enzymes encoded by the *dha* regulon. Here, glycerol is oxidized to dihydroxyacetone (DHA) by glycerol dehydrogenase (mainly encoded by *dhaD*) using NAD<sup>+</sup> as an electron acceptor. DHA then is phosphorylated by dihydroxyacetone kinase, an enzyme encoded by the ATP-dependent *dhaK1/dhaK* and/or PEP-dependent *dhaK2/dhaKLM/dhaK123*. DHAP obtained from respiration and/or fermentation is channeled into the glycolytic pathway (Fig. 2). A variety of organic acids and alcohols, including acetic acid, lactic acid, succinic acid, ethanol, BDO, and formic acid, are produced by the downstream glycolytic pathway (Ashok et al., 2011; Kumar et al., 2013a).

The anaerobic growth capability of *K. pneumoniae* using glycerol as a carbon source is attributed to its well-established reductive metabolism. In the reductive pathway, glycerol first undergoes a difficult rearrangement reaction catalyzed by glycerol dehydratase (GDHt) to yield 3-hydroxypropionaldehyde (3-HPA). Catalysis by GDHt requires coenzyme B<sub>12</sub>, the *de novo* synthesis of which is limited to only few genera of microorganisms. The 3-HPA is subsequently reduced to PDO by at least two 1,3-propanediol oxidoreductases (1,3-PDORs), including NADH-dependent DhaT (PDOR) and NADPH-dependent hypothetical oxidoreductase (HOR), along with the regeneration of NAD(P)<sup>+</sup> (Fig. 3). *E. coli* and other *enterobacter sp.* have the oxidative (respiratory) pathway of glycerol assimilation. However, these organisms lack the reductive pathway and cannot grow on glycerol under anaerobic conditions (Dharmadi et al., 2006; Kumar et al., 2013a).

## 2.2 Genes and enzymes involved in fermentative metabolism of glycerol

The *dha* regulon, induced by the DHA molecule and expressed in the absence of exogenous electron acceptors, encodes numerous genes needed for both the oxidative (fermentative) and reductive pathways (Celińska, 2012; Forage and Foster, 1982) (Fig. 4A). The *dha* regulon of *K. pneumoniae* contains the following genes arranged in the order as appears: *Kpk\_0615/dhaK*, *dhaM*, *dhaL*, *dhaK*, *dhaD*, *dhaR*, *orfW*, *CdAT*, *orfX (dhaG)*, *dhaT*,

*orfY*, *dhaB*, *dhaC*, *dhaE*, *orfZ* (*dhaF*), and *glpF*. Of these, *dhaB*, *dhaC*, *dhaE*, *orfX*, *orfZ*, and *dhaT* are relatively well documented for their functions in the reductive metabolism of glycerol. The *orfW* and *orfY* genes, although commonly found in the *dha* regulons of many organisms, are not as well described. The proteins encoded by the *dha* regulon of *K. pneumoniae* have high similarity (80%–95%) with those of *Citrobacter* species but not with those of *Clostridium* species (30%–80%). Surprisingly, the sequence similarity of *dha* proteins among *Clostridium* species is very low in some cases, even lower than their similarity to the corresponding proteins in *K. pneumoniae* or *C. freundii* (Celińska, 2012; Sun et al., 2003; Wei et al., 2014).

The complete *dha* regulon is found in only a few dozen species belonging to five different taxonomic groups, including four bacterial species [Actinobacteria, Firmicutes, Fusobacteria, and Proteobacteria (Gammaproteobacteria and Deltaproteobacteria)] and one archaea species, *Halalkalicoccus jeotgali* (Martins-Pinheiro et al., 2016). Interestingly, an incomplete *dha* regulon is present in more than 100 prokaryotes, suggesting that the enzymes of the regulon have other functions not restricted to the anaerobic growth on glycerol. A well-documented example is the one found in *Lactobacillus* species. *Lactobacillus* cannot grow on glycerol as the sole carbon source because it lacks the enzymes for oxidative assimilation of glycerol. However, the enzymes encoded in the incomplete *dha* regulon enable the production of PDO from glycerol, which accelerates cell growth in *Lactobacillus* grown on glucose (Årsköld et al., 2008; Martins-Pinheiro et al., 2016).

In the following sections, the major enzymes involved in fermentative glycerol metabolism and their genes are described in detail.

### **2.2.1 Glycerol dehydrogenase**

Encoded by the *dhaD* and/or *gldA* gene, glycerol dehydrogenase is an oxidoreductase that converts glycerol to DHA. The enzyme extracts electrons from glycerol and transfers

them to  $\text{NAD}^+$ . DhaD plays a major role during the anaerobic growth of *K. pneumoniae* on glycerol. Quantification of mRNA expression indicates that *dhaD* is highly induced when glycerol is used as carbon source. GldA expression is lower than that of DhaD and is thought to serve as a back up to DhaD during anaerobic growth on glycerol. However, DhaD and GldA exhibit a high level of similarity in amino acid sequence and have the same substrate-binding site (Wang et al., 2014).

Recently, DhaD was reported to catalyze the interconversion of acetoin and BDO for NADH disposal and recovery (Wang et al., 2014). The promiscuity of DhaD is responsible for BDO production in the  $\Delta budC$  mutant of *K. pneumoniae*. The contribution by DhaD to or its participation in BDO production is also supported by the observed increase in BDO synthesis and increased concentration of NADH in the presence of the *dhaD* gene. NADH is generated by the oxidation of glycerol and consumed by the production of BDO from acetoin. The dual physiological functions of DhaD allow *K. pneumoniae* to control the intracellular redox level. In response to the intracellular level of NADH, DhaD is able to switch its role from NADH production (glycerol oxidation) to NADH consumption (BDO production) (Wang et al., 2014). The diversion of glycerol flux from organic acids toward alcohols such as BDO also helps the cells to counteract intracellular acidification (Petrov and Petrova, 2009). Thus, this promiscuity confers different physiological roles to the enzyme, which include regulation of the intracellular  $\text{NADH}/\text{NAD}^+$  ratio, prevention of acidification, and storage of carbon and energy (Wang et al., 2014).

### 2.2.2 Dihydroxyacetone kinases

Dihydroxyacetone kinase (DHAK) phosphorylates DHA to DHAP, an intermediate of the glycolytic pathway. Sun et al. (2003) analyzed the genome of *K. pneumoniae* MGH 78578 and found, in addition to ATP-dependent DhaK I, a PEP-dependent DhaK II. DhaK I is encoded by the *dhaK* (*Kpk\_0615*) gene, while DhaK II is encoded by three genes, *dhaK*,

*dhaL* and *dhaM*, which correspond to the medium, small, and large subunits of the enzyme, respectively (Celińska, 2012; Sun et al., 2003; Wei et al., 2014). The names *dhaKLM* and *dhaK123* are synonyms used without distinction (Martins-Pinheiro et al., 2016; Sun et al., 2003; Wei et al., 2014). DhaK enzymes are found in many microorganisms. PEP-dependent DhaK II is present in *E. coli*, *L. lactis*, and *C. butyricum*, whereas ATP-dependent DhaK I is present in *Citrobacter freundii* (Garcia-Alles et al., 2004; Raynaud et al., 2011). The PEP-dependent DhaK II of *E. coli*, encoded by *dhaKLM*, is homologous to *dhaK123* (*dhaKLM*) of *K. pneumoniae* (Gutknecht et al., 2001). In a recent study, Wei et al. (2014) further elucidated the physiological roles of DhaK I and DhaK II of the *dha* regulon of *K. pneumoniae*. They found that disruption of ATP-dependent DhaK I had no significant effect on glycerol uptake, indicating that PEP-dependent DhaK II is the major contributor to the conversion of DHA to DHAP. As in *E. coli*, the subunits of DhaK II regulate expression of the *dha* regulon; *dha* regulon expression was suppressed by the disruption of *dhaK1* (*dhaK*) and *dhaK2* (*dhaL*). However, in contrast to *E. coli*, mutation of *dhaK3* (*dhaM*) in *K. pneumonia* did not upregulate the *dha* regulon but rather decreased cell growth and glycerol uptake.

### 2.2.3 Glycerol dehydratase and reactivating factor

Glycerol dehydratase (GDHt) is the first enzyme in the reductive pathway and catalyzes the dehydration of glycerol to 3-HPA. The same reaction can be carried out by diol dehydratase (DDHt), although the latter has a higher activity on 1,2-propanediol than on glycerol. GDHt requires coenzyme B<sub>12</sub> or S-adenosyl methionine (SAM) as a cofactor. *K. pneumoniae* has B<sub>12</sub>-dependent GDHt only, while strict anaerobes such as *clostridia* sp. have SAM-dependent GDHt. The three subunits of GDHt are encoded by the genes *dhaB1* (~1670 bp), *dhaB2* (~590 bp) and *dhaB3* (~430 bp), for which the gene products are the  $\alpha$  (large, 60.7 kDa),  $\beta$  (medium, 21.3 kDa), and  $\gamma$  (small, 16.1 kDa) subunits, respectively (Xu et al., 2009a). GDHt of *K. pneumoniae* is known to be present as a dimer of heterotrimers, ( $\alpha\beta\gamma$ )<sub>2</sub>.

204 The subunits of GDHt have **high homology among *K. pneumoniae* strains**; for example, *K.*  
205 *pneumoniae* XJPD-Li and U30909 strains showed >99% identity.

206 Coenzyme B<sub>12</sub> is present in the reaction center and plays an essential role in the  
207 catalysis of GDHt. The catalytic cycle often results in the mechanism-based inactivation of  
208 coenzyme B<sub>12</sub> due to the irreversible breakage of the chemical bond between cobalt (Co) and  
209 a carbon of the adenosyl moiety. Oxygen accelerates breakage of the Co–C bond. Oxygen is  
210 also known to inactivate GDHt, even in the absence of the substrate glycerol. The gene  
211 products of *orfZ/dhaG* (small subunit) and *orfX/dhaF* (large subunit) act as reactivating  
212 factors of GDHt (GdrAB) (Celińska, 2012; Sun et al., 2003). They reactivate GDHt by  
213 catalyzing the exchange of damaged for intact coenzyme B<sub>12</sub> in the presence of ATP and  
214 Mg<sup>2+</sup>/Mn<sup>2+</sup> (Celińska, 2012; Kumar et al., 2013a; Maervoet et al., 2011; Sun et al., 2003; Wei  
215 et al., 2014). The reactivase is a molecular chaperone that functions as a heterotetramer  
216 containing two elongated  $\alpha$  subunits (63 kDa) and two globular  $\beta$  subunits (14 kDa).  
217 Structurally, the  $\alpha$  subunit resembles both GroEL and Hsp70 chaperones, while the  $\beta$  subunit  
218 resembles that of the  $\beta$  subunit of glycerol dehydratase, except that it lacks some of the amino  
219 acids responsible for coenzyme B<sub>12</sub> binding (Liao et al., 2003).

220 In some organisms, the GDHt enzyme varies in terms of the number of subunits and  
221 their copies (Fig. 4B) (Liu et al., 2010). For example, in *Mesorhizobium loti* and  
222 *Mesorhizobium opportunistum*, the large and medium subunits (*dhaB1* and *dhaB2*) are fused  
223 together and encoded by a single gene. The gene fusion may arise by frameshift mutations.  
224 The active site of the fused enzyme differs slightly from that of unfused enzyme. Moreover,  
225 these microorganisms lack the genes encoding the reactivation factor, indicating that the  
226 reactivation process may not exist or is carried out by a different route. *Mycobacterium*  
227 *smegmatis* has two subunits (fused *dhaB1* + *dhaB2* and *dhaB3*) that are similar to those of *M.*  
228 *loti* and *M. opportunistum*; however, the reactivating factor differs in that it has one (large)

subunit. No small subunit has been identified in *M. smegmatis*. *Fusobacterium Ilyobacter polytropus* is unique in that it has genes coding for both B<sub>12</sub>-independent (SAM-dependent) and B<sub>12</sub>-dependent GDHt. The genes for B<sub>12</sub>-independent GDHt are present in the genomic DNA, while those encoding B<sub>12</sub>-dependent GDHt along with its reactivation factor are present in two copies, one in the chromosome and the other in a plasmid. Surprisingly, even with this gene redundancy, no significant production of 3-HPA and/or PDO has been reported in *I. polytropus* (Stieb and Schink, 1984). The B<sub>12</sub>-independent proteins showed no homology with B<sub>12</sub>-dependent GDHt but significant similarity with the pyruvate formate lyases (PFL) and PFL-activating enzymes and their homologues (Raynaud et al., 2003).

#### **2.2.4 1,3-Propanediol oxidoreductase and hypothetical oxidoreductase**

The 1,3-propanediol oxidoreductase (PDOR), responsible for the second step of the reductive pathway of glycerol, catalyzes the conversion of 3-HPA to PDO with the oxidation of NAD(P)H. 3-HPA is a toxic intermediate, and PDOR protects the cell from deleterious damage by reducing 3-HPA to PDO (Celińska, 2012). In *K. pneumoniae*, PDOR is encoded by the *dhaT* gene and is strictly NADH dependent. Active PDOR is a homo-octamer with a monomeric molecular mass of 43 kDa. Despite its importance, the kinetic properties of DhaT were elucidated only recently (Lama et al., 2015). DhaT has high specificity for 3-HPA among various aldehydes. Under physiological conditions at pH 7.0 and 37°C, the catalytic efficiency ( $k_{cat}/K_m$ ) of the forward reaction (3-HPA to PDO) was found to be 77.2/s. mM, which is 57-fold higher than that of the reverse reaction (1.3/s. mM; PDO to 3-HPA). The  $K_m$  (mM) values for 3-HPA and NADH (forward reaction) are 0.77 and 0.03, respectively, while those for PDO and NAD<sup>+</sup> (backward reaction) are 7.4 and 0.23, respectively. Furthermore, the maximum activity of DhaT for the reverse reaction was obtained at pH 9.0 and 55°C. These results strongly suggest that, under physiological conditions, DhaT catalyzes the reduction of 3-HPA to PDO rather than the oxidation of PDO to 3-HPA. Nevertheless, the

reverse reaction (PDO→3-HPA) becomes significant as the PDO concentration increases.

This factor is considered a serious drawback of using DhaT for PDO production from glycerol.

*K. pneumoniae* has several oxidoreductases in addition to PDOR that can reduce 3-HPA to PDO. One such enzyme, designated as hypothetical oxidoreductase (HOR), has been studied extensively and shares 89% identity with YqhD of *E. coli* (Zhu et al., 2009; Zhuge et al., 2010). Expression of *yqhD* in wild-type *K. pneumoniae* is not high, with an mRNA level ~20-fold lower than that of *dhaT*. However, deletion of *dhaT* increases the transcription of *yqhD* by more than 10-fold (Ko et al., 2015). This observation suggests that in addition to its wide-ranging activity on short-chain aldehydes, YqhD is a detoxification enzyme for many aldehydes (Jarboe, 2011). YqhD exhibits negligible activity on PDO (backward reaction), which is an important advantage over DhaT for use in the production of PDO from glycerol (Celińska, 2012; Chen et al., 2011; Zheng et al., 2006). However, YqhD is kinetically inferior to DhaT because of its low catalytic activity ( $k_{cat}/K_m$ ) on 3-HPA and the requirement of NADPH as a cofactor. The  $k_{cat}/K_m$  value of YqhD on 3-HPA (pH 7.0 and 37°C) is estimated to be 2.1/s. mM, which is 36-fold lower than that of DhaT (unpublished data). The use of NADPH as a cofactor prohibits this enzyme from being directly coupled to the reactions catalyzed by DhaD and other enzymes in glycolysis and the TCA cycle, by which NADH is mainly produced. Furthermore, excessive use of NADPH for PDO production can significantly reduce the cellular NADPH level and this can disturb many anabolic reactions requiring NADPH (Wang et al., 2003; Jarboe, 2011). NADH and NADPH differ in a phosphate group only, and it has been suggested that electrostatic interaction of this phosphate is the major factor distinguishing NADH from NADPH for enzymes using these cofactors. Comparison of the amino acid sequences of DhaT and HOR from *K. pneumoniae* and YqhD from *E. coli* has revealed that DhaT contains aspartic acid (Asp) at position 41,

while HOR and YqhD have valine (Val) at this position. The steric hindrance and electrostatic repulsion between Asp in the active site and the phosphate group of NADPH impede their binding. In contrast, the hydrophobic amino acid Val poses no such obstruction; thus, YqhD uses NADPH as a cofactor (Ma et al., 2010).

Despite their difference in cofactor preference, DhaT and YqhD are similar (44%) in amino acid sequences. *K. pneumoniae* has five enzymes that have at least 40% similarity in amino acid sequence with DhaT and YqhD: putative alcohol dehydrogenase (PaldH), L-1,2-propanediol oxidoreductase (FucO), ethanolamine-utilization enzyme (EutG), propanediol-utilization propanol dehydrogenase (PduQ), and bifunctional alcohol dehydrogenase (AdhE) (Ko et al., 2015). The roles of these enzymes in PDO production have been investigated.

### 2.2.5 Regulatory protein

The *dhaR* gene product is an important transcription factor that is responsible for expression of the *dhaT* and *dhaB* genes (Sun et al., 2003). This gene is found in *K. pneumoniae*, *C. freundii*, and *Clostridium botulinum* but not in every organism containing a complete *dha* regulon (Martins-Pinheiro et al., 2016). The presence of DhaR is thought to allow the *dha* operon to function as a separate regulatory system, independent of fumarate nitrate reduction (FNR). Under anaerobic conditions, FNR acquires DNA-binding properties and induces the expression of many anaerobiosis-related genes. DhaR contains domains for GAF (52-199), PAS (203-267),  $\sigma^{54}$  factor interaction, and histidine HTH-8. The PAS senses internal levels of energy charge, light, oxygen, and redox potential. The HTH-8 domain has a helix-turn-helix conformation and acts as a DNA binding structure. The  $\sigma^{54}$  factor interaction domain interacts with the  $\sigma^{54}$  factor of RNA polymerase and activates RNA transcription from the  $\sigma^{54}$  promoters (Buck et al., 2000; Taylor and Zhulin, 1999). Thus, the DhaR protein triggers the transcription of *dha* genes in response to intracellular levels of several important physiological parameters (Celińska, 2012; Sun et al., 2003). Zheng et al. (2006) investigated



the effect of overexpression of the putative regulatory gene *dhaR* in the *dha* regulon on glycerol metabolism in *K. pneumoniae*. *dhaR* overexpression increased PDOR activity up to 6.7-fold over that of the control wild-type strain, confirming the role of DhaR as a positive regulator of the *dhaT* gene. However, *dhaR* overexpression did not increase the activity of glycerol dehydratase (DhaB). Furthermore, despite the significant increase in PDOR activity, PDO production was unexpectedly low compared with that of the wild type (440 vs. 656.23 mM, respectively). More studies are required to determine the precise role of DhaR and its regulatory function in the *dha* regulon.

DhaR in *E. coli* (70% identity to that of *K. pneumoniae*) has also been studied. This protein stimulated transcription of the *dhaKLM* operon from a  $\sigma^{70}$  promoter. Interestingly, in *E. coli*, DhaL was a positive regulator of the *dhaKLM* operon while phosphorylated DhaM and DhaK were negative regulators of the same operon. DhaK contains the DHA-binding site. DhaL uses ADP as a cofactor for the double displacement of phosphate from DhaM to DHA. DHA is the oxidation product of glycerol metabolism; binding of DHA to DhaK reduces the affinity of DhaK for DhaR. In the presence of DHA, DhaL::ATP is dephosphorylated, leading to the displacement of DhaK by DhaL::ADP, which stimulates DhaR activity. In the absence of DHA, DhaL::ADP is phosphorylated back to DhaL::ATP by the phototransferase system (PTS), thereby inhibiting the binding of the phosphorylated complex (DhaL::ATP) to DhaR. This phosphorylation is mediated by DhaM, providing a phospho-histidine relay between the PTS and DhaL::ADP. This double-check mechanism of binding and turnover increases the selectivity such that the binding of nonphosphorylated compounds does not induce the *dha* operon (Bächler et al., 2005).

### **2.3 Genes and enzymes involved in respiratory assimilation of glycerol**

The *glp* regulon comprises the genes encoding proteins required for the conversion of glycerol to DHAP via sn-glycerol-3-phosphate. The five operons that constitute the *glp*

regulon are *glpFK*, *glpD* (anaerobic glycerol-3-phosphate dehydrogenase), *glp ACB* (aerobic glycerol-3-phosphate dehydrogenase), *glpTQ* (glycerol-phosphate permease/glycerophosphodiesterase), and *glpEGR*. The *glpF* gene encodes a cytoplasmic membrane protein that facilitates the diffusion of glycerol into the cell. The *glpE* gene encodes an acidic, cytoplasmic protein of 108 amino acids with a molecular weight of 12,082. The *glpG* gene encodes a basic, cytoplasmic membrane-associated protein of 276 amino acids with a molecular weight of 31,278 (Lin, 1976; Yang and Larson, 1998; Zeng et al., 1996). The biochemical functions of GlpE and GlpG are unknown. The *glpR* gene encodes the GlpR repressor protein. These five operons are located at three different positions on the chromosome. Transcription of these operons is subject to multiple controls, including catabolite repression mediated by cAMP-CRP and respiratory control mediated by the FNR and ArcA/ArcB systems. Moreover, each of the operons is negatively controlled by a repressor specific for the regulon (Lin, 1987; Zeng et al., 1996).

### 2.3.1 GlpR and other regulatory proteins

The GlpR repressor protein, predicted to contain 252 amino acids with a molecular weight of 28,048, belongs to the DeoR bacterial regulatory factor super family. **Although not studied extensively, GlpR of *K. pneumonia* is expected to have the same biochemical properties and regulatory functions as that of *E. coli*. GlpR negatively controls all the *glp* operons by binding to operators that overlap or are close to the *glp* promoters. In case of *E. coli*, glycerol-3-phosphate acts as an inducer and has a high affinity for GlpR ( $K_d = 31 \mu\text{M}$ ) (Lin, 1987; Weissenborn et al., 1992; Zeng et al., 1996). When complexed with glycerol-3-phosphate, GlpR cannot bind to the operator sites, and repression is relieved. Glycerol-3-phosphate is a product of GlpK and a substrate of GlpD/GlpABC; thus, the absence of GlpD causes a substantial increase in the level of glycerol-3-phosphate, resulting in induction of the *glp* regulon (Fig. 2). The GlpR protein exhibits differential binding affinity for the operators**

of corresponding operons, as follows: *glpFK* > *glpD* > *glpTQ* > *glpACB*. The *glpFK* promoter appears to be the strongest among the *glp* promoters. Therefore, the maximum strength of the *glpFK* promoter combined with the high affinity of GlpK for glycerol ( $K_m = 10 \mu\text{M}$ ) should result in sufficient accumulation of the inducer, glycerol-3-phosphate, in the presence of glycerol to relieve repression by GlpR. The relatively lower strength of the *glpD* promoter along with the low affinity of the GlpD enzyme for glycerol-3-phosphate ( $K_m = 1 \text{ mM}$ ) prevents excessive degradation of glycerol-3-phosphate that is needed for phospholipid synthesis when the exogenous source of glycerol-3-phosphate becomes limiting, until after accumulation of the inducer (Lin, 1987; Weissenborn et al., 1992). Recently, Jung et al. (2014) investigated the role of GlpR in the aerobic production of 3-HP from glycerol in *E. coli*. They found that elimination of this regulatory factor caused increased assimilation of glycerol and higher production of 3-HP and suggested that the deletion of *glpR* led to the upregulation of genes involved in glycerol transport and assimilation. *K. pneumoniae* has several other regulatory genes in addition to *glpR*: the global regulatory genes *hdeB*, *hdeD*, and *yfdX*, with unknown functions and *kvgS* and *kvgA*, encoding two proteins comprising a two-component signal transduction system (Sun et al., 2003).

### 2.3.2 Glycerol uptake/transport facilitator and other components

The glycerol facilitator GlpF is an integral membrane protein that forms aqueous pores. The GlpF pores selectively allow passive transport of glycerol and other molecules such as DHA and urea across the cytoplasmic membrane, but not water or ions (Heller et al., 1980; Reizer et al., 1993). The GlpF protein contains six membrane-spanning helices that are unique among prokaryotic transport proteins. These transmembrane helices are arranged in two bundles. The GlpF protein may function as a homodimer with the two six-member domains arranged in the membrane to form a channel (Weissenborn et al., 1992). *K. pneumoniae* has two glycerol transporter genes, one next to *orfZ/dhaF* in the fermentative

*dha* regulon (see Fig. 4A) and the other in the respiratory *glpFK* operon. Both *glpF* genes have a high similarity (85 %) to each other. When one but not both of the *glpF* genes is disrupted, the *glpF* deletion mutants of *K. pneumoniae* can consume glycerol and convert it to PDO at a high rate, similar to the wild type (unpublished data; Supplementary Fig. S1). This observation indicates that both *glpF* genes encode active glycerol transporter proteins. However, their presence is not essential for glycerol utilization in *K. pneumoniae* because the double-deletion mutant for both facilitators can still uptake glycerol and convert it to PDO. In *E. coli*, only one *glpF* gene is present in the *glpFK* operon; the GlpF protein shows a high similarity (80.9%) to that (encoded from *glpFK*) of *K. pneumoniae* (Sun et al., 2003). The importance of GlpF in *E. coli* also seems to be limited only at low levels of glycerol (<8 mM), as glycerol itself may pass through the membrane by diffusion at high concentrations (Richey and Lin, 1972).

### 2.3.3 Simultaneous operation of fermentative and respiratory pathways

The presence of the *dha* and *glp* regulons affords *K. pneumoniae* metabolic flexibility. The *dha* regulon enables fermentative utilization of glycerol, while the *glp* regulon enables respiratory metabolism of glycerol. During fermentative metabolism of glycerol, redox constraints force the production of PDO from glycerol, with a low ratio of carbon conversion yield to cell biomass. Moreover, ATP is formed through substrate-level phosphorylation, and NADH oxidation occurs through the formation of reduced metabolites. In the presence of oxygen, the redox constraints forcing the production of PDO are relieved; therefore, the carbon loss required for regeneration of NAD<sup>+</sup> is greatly reduced. As a consequence, glycerol utilization, biomass synthesis, and even the formation of many metabolites increase (Durnin et al., 2008). Interestingly, the fermentative route (*dha* regulon) is in operation under a wide range of oxygen levels, from microaerobic to highly aerobic conditions, as evidenced by PDO formation. For example, a high titer PDO (102.7 g/L) was produced under the condition

of 2.0 vvm aeration and even at 3.5 vvm, although the production was a little lowered in the latter case (Oh et al., 2012a). These results suggest that the fermentative and respiratory pathways might have evolved to complement each other for optimal control of cell growth under a wide range of aeration conditions except the strictly anaerobic one.

## 2.4 Carbon catabolite repression

Carbon catabolite repression (CCR) refers to the suppression of the use of less-preferred carbon sources, when a more preferred carbon, most prominently glucose, is present. Cyclic AMP, cAMP receptor protein (CRP), and the enzyme EIIA<sup>Glc</sup>, an intermediate in the phosphorylation cascade of the PTS, are key players in the CCR of enterobacteria. The EIIA<sup>Glc</sup>, a cytosolic protein, exists in phosphorylated and unphosphorylated forms. The phosphorylated form stimulates membrane-bound adenylate cyclase, which catalyzes the formation of cAMP; the cAMP-CRP complex then induces expression of the catabolic genes of less-preferred carbon sources. The unphosphorylated form of EIIA<sup>Glc</sup> is also responsible for inhibiting the transport of less-preferred carbon sources by inhibitory binding to permeases, causing so-called inducer exclusion (Eppler and Boos, 1999; Eppler et al., 2002). Glycerol assimilation in *K. pneumoniae* is greatly inhibited by CCR in the presence of glucose. This repression is related to not only low cAMP and cAMP-CRP but also inhibition of GlpK (which functions mainly under aerobic conditions) by fructose 1,6-bisphosphate, a metabolite produced from glucose. The unphosphorylated EIIA<sup>Glc</sup> is also known to allosterically inhibit GlpK (Holtman et al., 2001). Anaerobic glycerol metabolism mediated by the *dha* operons is also suppressed by the presence of glucose. Enzyme-level inhibition similar to that of the aerobic GlpK has not been reported, but we have observed that, in *K. pneumoniae*, transcription of the *dha* operons is greatly reduced in the presence of glucose (Suman et al., 2017).

Interestingly, glycerol exerts catabolite repression on the assimilation of maltose in *E. coli*. To exert catabolite repression, glycerol must be phosphorylated to glycerol-3-phosphate (Hogema et al., 1998; Eppler et al. 2002). However, unlike other carbon sources, further metabolism of glycerol is not required. According to Eppler and Boos (1999), glycerol-3-phosphate reduces expression of MalT, a positive activator of all *mal* genes. The key players in this repression are adenylate cyclase, EIIA<sup>Glc</sup>, and CRP. The growth on maltose of the mutants lacking EIIA<sup>Glc</sup> or containing truncated adenylate cyclase was no longer repressed by glycerol, and CRP-independent transcription of *malT* was also not influenced by glycerol either. In addition, stimulation of adenylate cyclase by phosphorylated EIIA<sup>Glc</sup> was controlled by glycerol-3-phosphate. The mutation in adenylate cyclase relieved the repression caused by glycerol-3-phosphate (Eppler et al., 2002). Although not investigated, we assume that the glycerol repression on maltose catabolism also hold true in *K. pneumoniae*.

Glycerol also exerts catabolite repression on the assimilation of citrate in *K. pneumoniae*. Citrate can be used as the sole carbon and energy source during anaerobic growth of *K. pneumoniae*, but the presence of glycerol suppresses the expression of the citrate fermentation genes. A reduced concentration of the cAMP–CRP complex has been postulated to be the cause of this repression (Meyer et al., 2001). Further research is needed to test the hypothesis.

### **3. *K. pneumoniae* as a microbial cell factory for the production of commodity chemicals**

During glycerol fermentation, *K. pneumoniae* produces many metabolites, including PDO, BDO, and lactic acid. This section describes the biotechnological production of bulk chemicals from glycerol in *K. pneumoniae*. 3-HP is not a natural product during the growth of *K. pneumoniae* on glycerol. However, its production is possible using a minor modification of the PDO synthetic pathway and is included here. The biochemical pathways

used for the synthesis of these chemicals are explained, and the metabolic engineering used to improve their production is discussed.

### 3.1 1,3-Propanediol

PDO, also known as trimethylene glycol, is a promising platform chemical that has two hydroxyl groups. With terephthalic acid, it can be co-polymerized to the novel co-polymer polytrimethylene terephthalate (PTT). PDO also has applications in the food and cosmetic industries. Currently, commercial PDO is mainly produced by recombinant *E. coli* (Maervoet et al., 2011; Saxena et al., 2009). The global demand for PDO was 60.2 kilotons in 2012 and is predicted to reach approximately 150 kilotons by 2019. The market for PDO is growing rapidly, from \$157 million in 2012 with projections up to \$560 million in 2019 (MarketsANDMarkets, 2012; Lee et al., 2015).

*K. pneumoniae* is one of the best native producers of PDO from glycerol. GDHt and PDOR are two important enzymes for the conversion of glycerol to PDO (Fig. 3). Physiologically, the synthesis of PDO from glycerol is essential for the anaerobic growth of *K. pneumoniae*, as described previously. The production of PDO requires two cofactors, coenzyme B<sub>12</sub> and NAD(P)H. *K. pneumoniae* has a *de novo* pathway for the biosynthesis of coenzyme B<sub>12</sub> that comprises more than 20 genes. NAD(P)H is generated through the oxidative metabolism of glycerol, during the production of cell biomass and/or oxidized (pyruvate and acetate) or partially reduced (BDO, lactic acid) metabolites. The production yield of PDO from glycerol varies depending on the cost of carbon for NAD(P)H production; the lower the cost, the higher the yield of PDO from glycerol. NADPH can also be the electron donor for the conversion of 3-HPA to PDO when YqhD is present. However, the contribution of NADPH to PDO production in native *K. pneumoniae* seems to be limited because the PP pathway, which is the main source of NADPH, is not active when glycerol is

the sole carbon source. Furthermore, *K. pneumoniae* has very low transhydrogenase activity for the conversion of NADH to NADPH (Zhang and Xiu, 2009; Chen et al., 2011).

Several metabolic engineering strategies have been employed to improve PDO production by *K. pneumoniae* (Table 2). Amplification of the reductive pathway by overexpressing DhaB, PDOR, or both has been attempted by many research groups. As indicated previously, 3-HPA is highly toxic, and its accumulation is detrimental to cell viability and toxic to PDO production. To reduce 3-HPA accumulation, overexpression of *dhaT* has been attempted. In one study, 3-HPA accumulation was reduced by *dhaT* overexpression, but no improvement in PDO production resulted (Hao et al., 2008). Overexpression of both DhaT and DhaD increased PDO production (56.3%) in batch culture but no increase in fed-batch culture (Chen et al., 2009). With another *K. pneumoniae* strain, DSM 2026, the same experiments were repeated but also with no increase in PDO production (Zheng et al., 2006). These results suggest that the reductive pathway in *K. pneumoniae* is evolutionarily well established (perhaps in concert with the oxidative pathway) and that PDO production is not limited by expression of the *dhaB* and *dhaT* genes of the reductive pathway. However, in one study where the *E. coli yqhD* gene was highly overexpressed to give ~10-fold higher PDOR activity (110 IU/mg total protein), a significant increase in PDO titer (25%; 67.6 g/L) and yield (from 0.53 to 0.62 mol/mol), along with less 3-HPA accumulation, was achieved (Zhu et al., 2009). This observation suggests that in the recombinant strain, NADPH can play an important role as a cofactor and that PDO production can be increased by modifying the reductive pathway. It is also probable that the effect of DhaB and PDOR on PDO production is highly dependent on the strain and/or culture conditions adopted.

Lactic acid, BDO, ethanol, and acetic acid are the major byproducts of PDO synthesis in *K. pneumoniae*. Inactivation of the metabolic pathways that give rise to these products has resulted in increased PDO synthesis. In particular, deletion of the *ldhA* gene encoding lactate



dehydrogenase has been highly beneficial. For example, by disrupting the *ldhA* gene, Xu et al. (2009b) significantly increased PDO production without lactic acid production. They observed increases in the PDO titer (95.4–102.1 g/L), conversion yield (0.48–0.52 mol/mol), and productivity (1.98–2.13 g/L h; Table 2). Similarly, Oh et al. (2012a) achieved the high PDO titer of 102.7 g/L by deleting *ldhA* in their own *K. pneumoniae* isolate. When lactic acid production was eliminated, more ethanol and succinic acid were generated from the pyruvate node. Reduction of these compounds has also been attempted by deleting *adhE* (encoding alcohol dehydrogenase) and *frdA* (encoding fumarate reductase). Although the production of ethanol and succinate was substantially reduced in the triple mutant ( $\Delta ldhA \Delta adhE \Delta frdA$ ), PDO production was only marginally increased compared with that of the single  $\Delta ldhA$  mutant. The failure of the triple mutant ( $\Delta ldhA \Delta adhE \Delta frdA$ ) to increase PDO production was mainly due to an increase in BDO production. Disruption of the BDO pathway encoded by the *bud* operon (*budA*, *budB*, and *budC*) has also been studied (Fig. 5). Although partially effective, individual inactivation of each of the three genes did not successfully decrease BDO production (Oh et al., 2012a; Wu et al., 2013). Deletion of the entire *bud* operon completely eliminated BDO synthesis (Kumar et al., 2016), but seriously hampered cell growth and glycerol consumption. Consequently, no increase in PDO production was resulted. Wu et al. (2013) attempted to decrease BDO production while increasing PDO production by inserting the *fdh* gene from *Candida boidinii* (NADH-forming formate dehydrogenase) into the *budC* locus. This protocol increased the PDO titer (62.3–72.2 g/L) and yield (0.47–0.57 mol/mol) and decreased the production of BDO by 52.2% and formic acid by 73.4%.

Acetic acid is one of the most toxic metabolites, accumulating in large quantities during glycerol fermentation by *K. pneumoniae* (Celińska, 2012). The toxic effects of acetate is realized even at the low concentration of 20 mM. Physiologically, acetate is a preferred metabolite for many *Enterobacter* sp. because its production is accompanied by ATP

generation. In a typical bioreactor run for PDO production by *K. pneumoniae*, the accumulation of acetic acid to >300 mM by the end is not uncommon, which often completely stops glycerol fermentation. Acetate is produced by Pta-Ack and PoxB when pyruvate formation is faster than its consumption, which is so-called ‘overflow metabolism’ (Fig. 6) (De Mey et al., 2007). In *K. pneumoniae*, the contribution of the *poxB* gene is not significant and its deletion does not much affect acetate production. In comparison, deletion of the *pta-ack* genes greatly reduces cell growth and results in high accumulation of pyruvate and pyruvate-based metabolites (Ko et al., 2017). Extensive studies to reduce or eliminate acetate production have been conducted in *E. coli*. Compared to *poxB*, *ackA* and/or *pta* were much more significant in reducing acetate production. However, deletion of the latter genes was not beneficial because the glycolytic flux and cell yield were seriously reduced and the accumulation of other byproducts such as pyruvate, lactate, and formate was greatly increased (Causey et al., 2004; De Mey et al., 2007). Several indirect approaches to decreasing pyruvate/acetate accumulation were investigated: stimulation of the anaplerotic pathway by overexpression of pyruvate carboxylase, PEP carboxylase, or PEP carboxykinase (Abdel-Hamid et al., 2001; Causey et al., 2004; Chao and Liao, 1993; Sanchez et al., 2005); overexpression of acetyl-CoA synthetase (ACS); increase of the TCA cycle throughput by deleting the transcriptional repressor *arcA* gene; and stimulation of the glyoxylate shunt by deleting the transcriptional repressor *iclR* gene (De Mey et al., 2007; Jeong et al., 2004; Lin et al., 2006). Moreover, the use of a mutant pyruvate dehydrogenase which is less sensitive to inhibition by NADH has also been examined (Kim et al., 2008). In *E. coli*, these approaches proved to be effective to varying extents under properly selected culture conditions. However, they have not yet been fully investigated in *K. pneumoniae*.

To eliminate byproduct formation, reduction of glycerol flux through the oxidative pathway at the glycerol node has also been attempted. The deletion of glycerol

dehydrogenase (*dhaD*) and/or dihydroxyacetone kinase (*dhaK*) resulted in decrease of the oxidative flux and increase of the conversion yield of glycerol to PDO. In addition, production of the byproducts lactate, ethanol, and succinate but not acetate decreased significantly, even without blocking the pathways leading to these byproducts (Seo et al. 2009; Horng et al. 2010). However, cell growth and glycerol consumption rate were also reduced. Interestingly, in these mutant strains, deletion of *dhaT* has almost no effect on PDO production. In the absence of *dhaT*, expression of an HOR, highly homologous to YqhD, was upregulated (see section 2.2.4).

The theoretical maximum yields of PDO on glycerol are 0.875 and 0.844 mol/mol under aerobic and anaerobic conditions, respectively (Celińska, 2012). However, experimentally, the yields are reported between 0.35 and 0.65 mol/mol (Saxena et al., 2009; Sun et al., 2008). The low yield is mainly attributed to the accumulation of unnecessary byproducts and inefficient metabolism for generating reducing equivalents (NADH) and ATP. If acetyl-CoA is fully oxidized through the TCA cycle (without byproducts formation) and the activity of the electron transport chain (ETC) oxidizing NADH is properly controlled, a significant increase in PDO yield is expected. Attaining this goal requires extensive pathway engineering of cellular metabolism. In bioreactor operation, the aeration rate should be optimized. Oxygen regulates gene expression and the enzymatic reactions of glycerol metabolism and determines the diversion of DHA to the PP pathway. Regeneration of NADH by the TCA cycle and its oxidation by ETC are also controlled by oxygen levels.

### 3.2 3-Hydroxypropionic acid

As with PDO, 3-HP is an important platform chemical and has been selected by the US Department of Energy (DOE) as one of the top 12 chemicals obtainable from biomass. 3-HP is a C3 bifunctional molecule and a structural isomer of lactic acid (Werpy and Petersen, 2004). 3-HP can be converted into a variety of useful chemicals such as acrylic acid and acryl

576 amide (Pina et al., 2011; Kumar et al., 2013a). 3-HP can be synthesized by chemical routes  
577 from acrylic acid, 3-propiolactone, 3-hydroxypropionitrile, allyl alcohol, vinyl acetate, and  
578 PDO. However, none of the chemical processes is commercially feasible at present due to the  
579 high cost of the starting materials, toxicity of intermediates, and/or the environmental  
580 incompatibility of the processes (Jiang et al., 2009; Pina et al., 2011; Kumar et al., 2013a).  
581 Many microorganisms, including both prokaryotes and eukaryotes, can naturally synthesize  
582 3-HP as either an intermediate or end product through a range of metabolic pathways.  
583 However, the production of 3-HP by these native microorganisms is too low to be  
584 commercially meaningful (Jiang et al., 2009; Kumar et al., 2013a).

585         In studies pursuing commercial production of 3-HP, glycerol is first converted to 3-  
586 HPA by GDHt. 3-HPA then is converted to 3-HP via the coenzyme A (CoA)-dependent or  
587 CoA-independent pathway (Fig. 3). In the CoA-independent pathway, 3-HPA is oxidized to  
588 3-HP by aldehyde dehydrogenase (ALDH); in the CoA-dependent pathway, 3-HPA is  
589 converted to 3-HP via 3-HP-CoA and 3-HP-phosphate. In *K. pneumoniae*, both the CoA-  
590 dependent and CoA-independent pathways are present. However, the titer of 3-HP produced  
591 by the CoA-dependent pathway is low as <5.0 g/L (Luo et al., 2011 and 2012). High  
592 production of 3-HP requires the disruption of oxidoreductases (e.g., DhaT, YqhD) and the  
593 overexpression of a proper aldehyde dehydrogenase (ALDH). Ashok et al. (2013a) developed  
594 a recombinant *K. pneumoniae* in which *dhaT* and *yqhD* were deleted, and the homologous  
595 *puuC* gene encoding the NAD<sup>+</sup>-dependent  $\gamma$ -glutamyl- $\gamma$ -aminobutyraldehyde dehydrogenase  
596 was overexpressed. The recombinant *K. pneumoniae* produced 3.8 g/L 3-HP in 12 h of flask  
597 culture, but only under appropriate aeration (i.e., microaerobic) conditions. Under anaerobic  
598 conditions, PDO (instead of 3-HP) was obtained as the main product, even though two  
599 oxidoreductases were disrupted and *puuC* was highly overexpressed. In contrast, highly  
600 aerobic conditions produced a high cell mass without much accumulation of either 3-HP or

1,3-PDO. In a glycerol-fed batch bioreactor experiment under a constant dissolved oxygen (DO) concentration of 5% (considered “proper” microaerobic condition), the recombinant *K. pneumoniae*  $\Delta dhaT \Delta yqhD$  overexpressing both PuuC and DhaB produced >28 g/L 3-HP in 48 h, with a yield of >40%. In contrast, the same strain produced only 10.5 g/L 3-HP when cultivated under the higher 10% DO conditions. Several important conclusions regarding 3-HP production by *K. pneumoniae* were made, as follows. First, in addition to DhaT and YqhD, *K. pneumoniae* has more unidentified oxidoreductases that can produce PDO from 3-HPA. Second, to produce 3-HP rather than PDO, proper aeration to regenerate  $NAD^+$  is essential. Third, excessively high aeration decreases the expression of the Dha regulon and synthesis of coenzyme B<sub>12</sub>, an essential cofactor for GDHt catalysis.

To eliminate PDO production even under limited aeration conditions, Ko et al. attempted to identify and disrupt other potential PDORs from *K. pneumoniae* (Ko et al., 2015). A mutant strain devoid of *dhaT*, *yqhD*, *ahpF*, and *adhE* genes was developed; however, the mutant neither eliminated 1,3-PDO production nor increased 3-HP production. Again, this result suggests that *K. pneumoniae* has more unidentified oxidoreductases, and thus, the complete elimination of 1,3-PDO production during 3-HP production is highly challenging. To address the important problem, of  $NAD^+$  regeneration and coenzyme B<sub>12</sub> production, Ashok et al. (2013b) studied anaerobic respiration using nitrate as an external electron acceptor. They attempted to regenerate  $NAD^+$  from NADH by nitrate reduction while maintain anaerobic conditions to synthesize and stabilize the oxygen-sensitive coenzyme B<sub>12</sub>. Disruption of the *glpK* gene (encoding for glycerol kinase) was also necessary because with *glpK* intact, the rate of anaerobic respiration was too fast, and most of the glycerol was metabolized through GlpK to produce biomass, as in the case of highly aerobic cultivation. The anaerobic glycerol fermentation in the presence of nitrate was successful: 1.74 g/L 3-HP was obtained from a 12-h flask culture and 22 g/L 3-HP was produced in a 48-

h fed-batch bioreactor culture (Ashok et al., 2013b) (Table 3). However, disadvantages were noted, including the requirement for large amounts of nitrate and the toxic effect of nitrite generated as a reduced intermediate of nitrate. In a recent study, Li et al. (2016) optimized a promoter for ALDH expression, culture conditions, and metabolic flux to achieve high-level production of 3-HP in their *K. pneumoniae* isolate. One recombinant strain, *K. pneumoniae* (pTAC-*puuC*) expressing *puuC* under the IPTG-inducible *tac* promoter, produced 73.4 g/L 3-HP in a bioreactor with a molar yield of glycerol of 0.52 and productivity of 1.53 g/L h. Further, elimination of the *ldh1*, *ldh2*, and *pta* genes elevated the titer and molar yield to 83.8 g/L 3-HP and 0.54 mol/mol, respectively, with decreased cell growth and productivity. This titer of 3-HP is the highest produced by *K. pneumoniae* to date.

As another approach to addressing the problem associated with NAD<sup>+</sup> regeneration and coenzyme B<sub>12</sub> synthesis in 3-HP production, co-production of 3-HP and PDO has been investigated. The simultaneous production of 3-HP and PDO can eliminate the dependency on oxidative metabolism of glycerol (and electron transport chain), as the cofactor required for 3-HP production is regenerated by PDO production or vice versa. Furthermore, the problems associated with the expression of genes for the assimilation of vitamin B<sub>12</sub> and glycerol can be alleviated substantially because co-production can be performed under anaerobic or microaerobic conditions (Kumar et al., 2012, 2013b). Ashok et al. (2011) developed a recombinant strain of *K. pneumoniae* DSMZ by overexpressing ALDH and deleting *dhaT*. With *dhaT* intact, PDOR activity was too high compared with that of ALDH, even with overexpression of the latter enzyme from a multicopy plasmid. The recombinant *K. pneumoniae* DSMZ accumulated 16.0 g/L 3-HP and 16.8 g/L PDO in 24 h, and the cumulative yield of these two metabolites on glycerol was 51% (Table 3). Huang et al. (2012) studied the same co-production using their *K. pneumoniae* isolate overexpressing ALDH and obtained 24.4 3-HP and 49.9 g/L PDO, respectively, after 24 h under anaerobic conditions.

The cumulative molar yield of the two metabolites reached 0.61 (0.18 for 3-HP and 0.43 for PDO). In a following study, they showed that the maximum titer of 48.9 g/L 3-HP (along with 25.3 g/L PDO) was achieved at 1.5 vvm, while the highest concentration of 38.4 g/L PDO (along with 16.6 g/L 3-HP) was obtained at 0.1 vvm (Huang et al., 2013a).

During 3-HP production from glycerol by *K. pneumoniae*, the accumulation of such byproducts as lactic acid, ethanol, acetic acid, formic acid, and 2,3-BDO is a serious consideration, as in the case of 1,3-PDO production (Ashok et al., 2013a,b; Li et al., 2016). Deletion of *ldhA* reduces lactic acid formation and increases the 3-HP yield. However, deletion of other genes such as *adhE*, *pfl*, *budBAC*, and *pta-ack* had little desired effect and seriously hampered cell growth and glycerol assimilation (unpublished data). Even with the co-production of 3-HP and 1,3-PDO, where oxidative glycerol assimilation was greatly reduced, accumulation of these byproducts was a serious drawback (Ko et al., 2017). To prevent the formation of lactic acid along with other byproducts, Kumar et al. (2013b) employed resting cells of recombinant *K. pneumoniae* J2B overexpressing ALDH and devoid of lactate dehydrogenase (*ldhA*). For this strain, final titers of 3-HP and PDO of 22.7 and 23.5 g/L, respectively, were obtained without lactic acid accumulation. The cumulative product yield increased to 0.77.

In addition to *K. pneumoniae*, *E. coli* has been extensively studied for 3-HP production from glycerol. Several successful results have been reported by Samsung Ltd. Co. Korea. One study (Chu et al., 2015) used an engineered strain (*E. coli* W3110  $\Delta$ *ackA-pta*  $\Delta$ *yqhD\_dhaB\_mutant gabD4*) harboring an active ALDH mutant (designated as GabD4) from *Cupriavidus necator*, producing 71.9 g/L 3-HP with a productivity of 1.8 g/L h. However, unlike *K. pneumoniae*, *E. coli* does not synthesize vitamin B<sub>12</sub> naturally, and it was necessary to add the coenzyme externally. Furthermore, a significant amount of glucose for cell growth was required because for *E. coli*, glycerol is a less-preferred carbon source than glucose.

Consequently, the researchers at Samsung initially grew cells to a high density (40–300 OD<sub>600</sub>) on glucose and then produced 3-HP by feeding glycerol (Chu et al., 2015; Kim et al., 2014a). In comparison, *K. pneumoniae* can produce a high concentration of 3-HP growing purely on glycerol at a much lower cell concentration (OD<sub>600</sub>, <20) (Huang et al., 2013a,b; Li et al., 2016). Together, the use of glucose for cell proliferation, exogenous addition of expensive cofactor vitamin B<sub>12</sub>, and low cumulative yield of 3-HP (on glucose plus glycerol) increase the production cost. Thus, *K. pneumoniae* can be considered a better biocatalyst than *E. coli*. However, for the commercial production of 3-HP, many other factors are important, including biosafety, process stability, and downstream processing. Further studies to assess the potential of these strains as hosts for 3-HP production are needed.

### 3.3 2,3-Butanediol

BDO has many applications in the pharmaceutical, biomedical, and other chemical industries for the production of printing inks, perfumes, fumigants, spandex, moistening and softening agents, and plasticizers (e.g., cellulose nitrate, polyvinyl chloride, polyacrylates) (Celińska and Grajek, 2009; Ji et al., 2011). BDO can also be used as an antifreezing agent and octane booster for petrol as is or can be converted to useful derivatives such as 1,3-butadiene, and diacetyl and methyl ethyl ketone.

Unlike 3-HP and PDO, BDO is a product of oxidative metabolism. Of the three stereoisomers of BDO, *K. pneumoniae* mainly produces the *meso* form (Ji et al., 2011). BDO is synthesized from pyruvate via  $\alpha$ -acetolactate and acetoin. The C5 intermediate,  $\alpha$ -acetolactate, is formed from two molecules of pyruvate by self-condensation catalyzed by  $\alpha$ -acetolactate synthase (ALS; *budB*).  $\alpha$ -Acetolactate is decarboxylated to **acetoin by  $\alpha$ -acetolactate decarboxylase (*budA*)**, and acetoin is reduced to BDO by 2,3-butanediol dehydrogenase/acetoin reductase (*budC*) using NADH as a reductant. In the presence of oxygen,  $\alpha$ -acetolactate is spontaneously decarboxylated to diacetyl, further reduced to acetoin



by action of diacetyl reductase, and then reduced to BDO.  $\alpha$ -Acetolactate is also the precursor of branched-chain amino acids (valine, leucine, and isoleucine); thus, elimination of the *bud* operon can arrest *K. pneumoniae* growth when cultured on glycerol minimal medium (Kumar et al., 2016). In the anaerobic process, one NADH molecule is required for the conversion of two pyruvates to one BDO, while the aerobic process requires two NADH (Fig. 5). The enzyme DhaD may also contribute to BDO formation (see Section 2.2.1). In the BDO pathway itself, the conversion of two pyruvates to BDO regulates or is regulated by the intracellular NADH/NAD<sup>+</sup> ratio, similar to other fermentative pathways. Another important role of the BDO pathway is to reduce intracellular acidification by converting acids to the neutral metabolite, BDO (Celińska and Grajek, 2009; Ji et al., 2011).

Although BDO is one of the main byproducts of glycerol fermentation by *K. pneumoniae* (Petrov and Petrova, 2009; Durgapal et al. 2014), its production from glycerol has not been studied extensively. In few studies using glycerol as the carbon source, Petrov and Petrova (2009) focused on the effects of pH and aeration with their *K. pneumoniae* isolate G31. BDO production was greatest at alkaline initial pH when tested at a pH range of 5–8. Without pH control a sharp pH drop occurred initially due to the production of organic acids, and this pH drop triggered the synthesis of BDO, which is considered the only non-inhibitory compound produced by the oxidative metabolism of glycerol. Once neutral pH is restored with the rising concentration of BDO, catabolism shifted toward the production of ATP-generating acetic acid. The amount of BDO generated in each cycle (pH drop and neutralization) was dependent on the extent of the pH drop. In one study, a BDO concentration of 49.8 g/L was reached when the initial pH was set at 8.0 and no pH control was given during the rest of fermentation. PDO was the major byproduct (11.6 g/L) (Table 4) and its production decreased by exclusion of Co<sup>2+</sup>, an essential component of coenzyme B<sub>12</sub>. They also found that intensive aeration led to a significant increase in BDO production; an

increase in aeration rate from 1.1 to 2.2 vvm enhanced the yield from 0.38 to 0.46 mol/mol and productivity from 0.24 to 0.36 g/L h. BDO production was also increased by forced pH fluctuations of the culture medium with discrete  $\Delta\text{pH}$  values (1.0, 2.0, and 3.0) at predetermined time intervals (6, 12, and 24 h). The highest BDO concentration (70.0 g/L) was produced by increasing the pH by one unit ( $\Delta\text{pH} = 1.0$ ) every 12 h (Petrov and Petrova, 2010). The mechanism behind the effect of this peculiar pH change on BDO production has not been elucidated. Although BDO is readily produced from glycerol by *K. pneumoniae*, the volumetric productivity is low due to the long fermentation time (generally >150 h). Huang et al. (2013b) isolated two *Klebsiella* strains that produce BDO plus acetoin with a total yield of 0.44–0.45 (g/g). One isolate utilized 64.5 g/L glycerol in 78 h and produced 24.9 g/L BDO, 3.9 g/L acetoin, and 3.2 g/L ethanol. Again, the volumetric productivity was not high.

Metabolic engineering efforts with *K. pneumoniae* to improve BDO production from glycerol are limited. Using glucose as carbon source, Kim et al. (2014b) reported that overexpression of the BDO pathway (*budA*, *budB*) in the *ldhA* deficient *K. pneumoniae* strain KCTC2242 improved BDO production and resulted in 90 g/L BDO, with a productivity of 2.75 g/L. h. Because both glucose and glycerol are converted through the common intermediate pyruvate, BDO production from glycerol is also expected to increase by amplification of the *bud* operon. Further work with glycerol as carbon source is required to prove this effect. Recently, co-production of BDO and PDO from glycerol has also been attempted (Park et al., 2017). In glycerol fermentation, both alcohols always appear simultaneously and co-production seems to be beneficial for cells to maintain redox balance during anaerobic growth. Park et al. (2017) found that by using the *K. pneumoniae* mutant deficient of *ldhA* and *mdh*, 50.1 g/L BDO was produced along with 63.8 g/L PDO in 75 h. The ratio of the two alcohols varied according to aeration; with increasing aeration, the ratio of [BDO]/[PDO] increased as observed by Petrov and Petrova (2010). In the 5,000 L pilot-

scale fed-batch fermentation using crude glycerol, they could produce 114 g/L diols (70 g/L PDO and 44 g/L BDO) with the yield of 0.60 diols/glycerol (g/g) and a productivity of 2.2 g/L. h of diols. If separation is not an issue, co-production of the two diols can be a good option.

### 3.4 Lactic acid

Lactic acid has applications in the food, chemical, cosmetic, and pharmaceutical industries. Lactic acid is a monomer that can be polymerized to yield the biodegradable plastics polylactic acid (PLA) and poly(3-hydroxybutyrate-co-lactate). Two optical isomers of lactic acid, D- and L-lactic acid, are produced in racemic mixtures depending on the chemical route used. Microbial fermentation can yield optically pure isomers. Currently, nearly all commercial lactic acid comes from microbial fermentation (Abdel-Rahman et al., 2013; Abdel-Rahman and Sonomoto, 2016).

Because of the availability of highly efficient lactic-acid-producing microbes derived from *E. coli*, lactic acid bacteria, and even acid-tolerant yeasts (Abdel-Rahman et al., 2013), *K. pneumoniae* has received little attention as a lactic acid producer. However, as previously described, lactic acid is a major byproduct of glycerol fermentation by *K. pneumoniae*, and its potential as a producer is high. This bacterium contains metabolic pathways for the production of both L- and D-isomers (Fig. 7). They are synthesized through the pyruvate and/or methylglyoxal pathways in *K. pneumoniae* (Ashok et al., 2011). When produced from pyruvate, lactate dehydrogenase (*ldhA*) reduces pyruvate using NADH as reductant. In the methylglyoxal route, DHAP is converted to methylglyoxal, which can be transformed to both D- and L-lactic acid. For the production of L-lactic acid, methylglyoxal is reduced to L-lactaldehyde, which is subsequently oxidized to L-lactic acid. Using glutathione, D-lactic acid is obtained through simultaneous aldehyde group oxidation and keto group reduction of methylglyoxal (Booth et al., 2003; Booth, 2005). The methylglyoxal route has several

disadvantages. Methylglyoxal is highly toxic, and its accumulation can severely impair other metabolisms and even lead to cell death. Second, the route is energetically inefficient, with an ATP yield of  $-1$ . The L- or D-lactic acid product can be converted back into pyruvate by respiratory lactate dehydrogenase (Mazumdar et al., 2010, 2013).

Several studies showing the potential of *K. pneumoniae* as a lactic acid producer are available (Table 4). For example, *K. pneumoniae* DSMZ is reported to accumulate 22.7 g/L lactic acid in 24 h during co-production of 3-HP and PDO from glycerol. The titer of lactic acid was more than that of any of the targeted products, and the yield of lactic acid on glycerol alone was 0.31 (Ashok et al., 2011). Similarly, fed-batch cultivation of *K. pneumoniae* BLh-1 under oxygen-limited conditions gave lactic acid as the main product, outcompeting PDO; 59 g/L lactic acid was produced in 40 h when crude glycerol from biodiesel synthesis without any purification was used (Rossi et al., 2013). Durgapal et al. (2014) also found that *K. pneumoniae* J2B and DSMZ strains produced lactic acid at 37.0 and 47.8 g/L, respectively (Table 4); in both strains, lactic acid levels surpassed those of the target product PDO. Recently, Feng et al. (2014) engineered *K. pneumoniae* ATCC25955 by overexpressing *ldhA* and deleting *dhaT* and *yqhD* (recall that these two genes are mainly responsible for PDO production). They found that the recombinant produced 142.1 g/L of optically pure D-lactic acid from glycerol in fed-batch cultivation under microaerobic conditions (2.5 vvm and 400 rpm). This result is the highest lactic acid concentration produced from glycerol reported to date. *E. coli* has also been engineered for the production of D- and L-lactate from glycerol (Mazumdar et al., 2010, 2013). For diverting glycerol flux toward L-lactic acid in *E. coli*, the chromosomal copy of D-lactate dehydrogenase was replaced with *Streptococcus bovis* L-lactate dehydrogenase (Mazumdar et al., 2013). Moreover, gene encoding *lldD* (respiratory L-lactate dehydrogenase) was inactivated to prevent the consumption of L-lactic acid, and the methylglyoxal route ( $\Delta mgsA$ ) was blocked

to prevent production of toxic methylglyoxal and formation of racemic mixtures of D- and L-lactic acid (Fig. 7). Furthermore, the respiratory route for glycerol assimilation was overexpressed (*glpK* and *glpD*) to improve the ATP yield of the metabolic pathway through transfer of electrons from glycerol-3-phosphate to oxygen (i.e., oxidative phosphorylation). The engineered *E. coli* strain ( $\Delta pflB \Delta pta \Delta adhE \Delta frdA \Delta mgsA \Delta lldD \Delta ldhA::ldhA\_glpK\_glpD$ ) produced 50 g/L of L-lactate from 56 g/L of crude glycerol, at a yield 93% of the theoretical maximum and with high optical (99.9%) and chemical (97%) purity (Mazumdar et al., 2013). The approaches proven successful in *E. coli* should be applicable to *K. pneumoniae*. These studies, along with the results by Feng et al. (2014), suggest that *K. pneumoniae* is a good host for the production of D- and L-isomers of lactic acid, with high optical purity, from glycerol.

### 3.5 Ethanol and other metabolites

*K. pneumoniae* has also been studied for the production of ethanol, succinic acid, 2-ketogluconic acid, catechol, *cis,cis*-muconic acid and 2-butanol although its potential for such production has not been fully explored (Cheng et al., 2013; Jung et al., 2015; Oh et al., 2014; Wang et al. 2015; Wei et al., 2013). Of these chemicals, ethanol production has been relatively well studied. Ethanol is a renewable energy source and is widely used as a fuel additive for partial gasoline replacement. Currently, commercial ethanol is produced from the fermentation of sugar and starch. However, glycerol is one of the cheapest substrates for ethanol production, estimated as nearly 40% cheaper than sugar-based production (Yazdani and Gonzalez, 2007). This factor has encouraged the development of microbial strains for fermentative conversion of glycerol to ethanol. The metabolic route by which ethanol is synthesized from glycerol is shown in Fig. 6. Oh et al. (2011, 2012b) constructed a mutant strain of *K. pneumoniae* (termed GEM167) through  $\gamma$ -irradiation. PDO synthesis in this mutant strain was lower than that of wild type (0.2 vs. 7.93 g/L, respectively), while ethanol

accumulation was higher (8.6 vs. 1.1 g/L, respectively). The mechanism underlying these changes remains unclear. Introduction of the pyruvate decarboxylase (*pdc*) and aldehyde dehydrogenase (*adhII*) genes into mutant strains lacking lactate dehydrogenase (*ldhA*) resulted in the production of 31.0 g/L (673.9 mM) ethanol, with a yield of 0.89 (mol/mol) and productivity of 1.2 g/L h (Oh et al., 2012b), the highest level of ethanol production from glycerol reported to date. More extensive studies including estimation of the maximum achievable titer and microbial tolerance to high-concentration ethanol, both of which are key factors for commercialization, are required to assess the potential to produce ethanol from glycerol by the use of *K. pneumoniae*.

#### **4. Challenges to bioconversion of glycerol using *K. pneumoniae* as a biocatalyst**

The design of microbial cell factories is gaining unprecedented momentum as metabolic engineering is progressively aided by advances in synthetic biology and multi-omics analyses. Several bio-based chemicals have already been launched successfully in the marketplace, and others are in the pipeline. For a bulk chemical to be considered for commercial production, the production process should achieve a product concentration of 100 g/L, carbon yield of 0.50 (g/g), and volumetric productivity of 2.0 g/L/h. The aforementioned products produced by *K. pneumoniae* from glycerol are no exception. Challenges to the use of *K. pneumoniae* as a microbial cell factory using glycerol as a carbon source are described.

##### **4.1 Pathogenicity**

The pathogenicity of *K. pneumoniae* is one of the major obstacles to its commercial application. *K. pneumoniae* is a Bio-safety Level 2 microorganism and causes nosocomial and urinary tract infections. Moreover, capsulated *K. pneumoniae* cells are difficult to separate from culture broth, complicating the downstream processing. The biotechnological application of this bacterium will require attenuation of its pathogenicity and reduction of biosafety concerns. *K. pneumoniae* produces a number of virulence factors that contribute to

its pathogenicity, including lipopolysaccharide (LPS), capsular antigens, fimbrial adhesins, siderophores, and O antigens (Albertí et al., 1993; Fang et al., 2004; Podschun and Ullmann 1998). The mechanism by which this bacterium causes disease is still obscure because most studies have been conducted on a limited number of virulence factors (El Fertat-Aissani et al., 2013).

In the last decade, efforts have been made to overcome the pathogenicity of *K. pneumoniae* (Shrivastav et al., 2013), suggesting hope for the generation of nonpathogenic *K. pneumoniae*. The two major pathogenic determinants in *K. pneumoniae* are the capsular polysaccharides (CPS), which cover the entire bacterial surface, and LPS, expressed on the outer bacterial membrane. LPS consists of lipid A, a core oligosaccharide, and an O-antigenic polysaccharide. In a highly virulent strain of *K. pneumoniae*, mutations in genes involved in LPS biosynthesis (*waaC*, *waaF*, and *wabG*) resulted in a decrease in bacterial colonization and virulence (Izquierdo et al., 2003; Jung et al., 2013). Mutant strains devoid of the outer core LPS were also found to be deficient in the “cell-attached” capsular polysaccharides covering the bacterial surface. These mutations abolished the highly virulent characteristics of pathogenic *K. pneumoniae* when tested in different animal models. Furthermore, these mutants were more sensitive to several hydrophobic compounds than were wild-type strains. Reintroduction of the *waaC*, *waaF*, and *wabG* genes into *K. pneumoniae* rescued the pathogenic properties, confirming the role of these genes in pathogenicity.

Another important study was conducted by Lin et al. (2012), in which eight highly conserved residues of the MagA protein (G308, G310, G334, G337, R290, P305, H323, and N324) encoded by the mucoviscosity-associated gene A (*magA*) of *K. pneumoniae* were subjected to site-directed mutagenesis. The *magA* gene contributes to the biosynthesis of K1 capsular polysaccharide (CPS), which correlates with pathogenic phenotypes including mucoviscosity, serum and phagocytosis resistance, and virulence. Alanine substitutions at

R290A or H323A abolished MagA function, with annihilation of CPS production, serum resistance, and virulence. Some *K. pneumoniae* strains showing less pathogenicity were also isolated. For example, *K. pneumoniae* strain J2B produced less LPS and demonstrated a high sensitivity to many antibiotics and lower pathogenicity. Moreover, its biomass was readily separated from fermentation broth by centrifugation (Arasu et al., 2011). Future studies should include further engineering of these less pathogenic strains to make them completely nonpathogenic, followed by careful evaluation of their toxicity.

#### **4.2 Complex glycerol metabolism**

Although glycerol metabolism has been extensively studied, the mechanism and regulation of its gene expression in *K. pneumoniae* have not been fully elucidated. The diversity of glycerol metabolism should be a great advantage for *K. pneumoniae* survival in different environments and for its biotechnological utilization as well. However, a high level of complexity poses difficulty in the engineering of a strain to suit our purposes. Metabolic complexity is likely responsible in part for the fact that, despite serious attempts, the molar yield of many important products such as PDO and 3-HP is still far below the theoretical maximum. Efforts based on our current knowledge, such as the overexpression of DhaB, DhaT, YqhD and AldH and inactivation/deletion of GlpK, DhaD, DhaK and DhaKLM, were not satisfactory (Maervoet et al. 2011; Kumar et al. 2013a; Kumar et al., 2016). Another challenge is to understand how the ratio of glycerol distribution between respiratory and fermentative routes is controlled. Because the  $K_M$  of glycerol kinase toward glycerol is low, it has been speculated that a major fraction of glycerol flows through the respiratory route when oxygen is present. However, some experimental results do not support this hypothesis. For example, deletion of *glpK* does not decrease the glycerol flow through the oxidative pathway in the presence of oxygen (Ashok et al. 2013a). In fact, wild-type and mutant ( $\Delta glpK$ ) strains of *K. pneumoniae* demonstrate similar cell growth, glycerol consumption, and PDO



production profiles under (micro)aerobic conditions. These observations suggest the presence and/or involvement of other kinases.

Another challenging issue for the use of *K. pneumoniae* is its inefficient TCA cycle (Cabelli, 1955). Several genes, including isocitrate dehydrogenase (*icd*), fumarase (*fumA*), and malate dehydrogenase (*mdh*), in *K. pneumoniae* are significantly less transcribed than those in *E. coli* (unpublished data). Furthermore, the activity of isocitrate dehydrogenase was more than sevenfold lower in *K. pneumoniae* than in *E. coli*. If the TCA cycle is inefficient, achieving a high cell density, which is essential for improving the productivity of target metabolites, is difficult. Moreover, achieving high-yield generation of NADH with little consumption of glycerol carbon is difficult. Furthermore, the overflow metabolism that produces highly toxic acetate becomes more serious. These problems have been well documented in PDO production. In a recent study, Kumar et al. (2016) attempted to increase PDO production yield by blocking the lactate and BDO pathways, while diverting carbon flux through the TCA cycle for efficient regeneration of NADH. However, this effort resulted in the detrimental accumulation of pyruvate, acetate, and ethanol, causing low NADH regeneration and low PDO production. Neither increased aeration nor the addition of good nitrogen sources (complex nitrogen sources) alleviated the accumulation of these intermediates or activated the TCA cycle. Furthermore, the accumulation of pyruvate quickly terminated glycerol assimilation. *K. pneumoniae* has a well-established BDO production pathway that likely compensates for its inefficient TCA cycle, avoiding the overflow metabolism and carbon traffic at the pyruvate node. If this is the case, co-production of BDO along with PDO or 3-HP, rather than single production of PDO or 3-HP, should be more appropriate, which suits the nature of *K. pneumoniae*. In fact, when BDO and PDO were co-produced, more PDO was generated than during PDO production alone. More studies are required to gain a better understanding of glycerol metabolism, the slow operation of the

TCA cycle in *K. pneumoniae*, and its effect on the selection of target products and/or development of suitable strains and processes. Otherwise, it will be difficult to exploit this potential biocatalyst for commercial manufacturing of valuable chemicals.

#### **4.3 Toxicity of intermediate metabolites and end-products**

A general but major challenge to the use of microbial cell factories for producing bulk chemicals is the toxic effect of target products and metabolic intermediates at high concentrations. In particular, during the production of PDO or 3-HP from glycerol, the highly toxic intermediate 3-HPA is inevitably generated. Aldehydes generally damage DNA, inactivate enzymes, and inhibit DNA synthesis. 3-HPA causes irreversible cessation of metabolic activity and cellular growth, even at very low concentrations of 15–30 mM (Hao et al., 2008; Zheng et al., 2008). Native *K. pneumoniae* does not accumulate 3-HPA at toxic levels during PDO production, but in engineered strains that overexpress DhaB, high 3-HPA accumulation is often experienced. To avoid 3-HPA accumulation, enzymes that consume 3-HPA, ALDH (for 3-HP production), and PDOR (for PDO production) are expressed at sufficiently (sometimes excessively) high levels compared with that of DhaB (Lim et al., 2016). Intermediates such as methylglyoxal, DHA, DHAP, and glyceraldehyde-3-phosphate are also known to be toxic.

Organic acids and alcohols are toxic at high concentrations. Generally, organic acids are more toxic than their corresponding alcohols because they disturb the intracellular pH and have anion-specific effects on metabolism (Chun et al., 2014; Warnecke and Gill, 2005). When accumulated in cell membranes, organic compounds damage membrane integrity and inhibit the activity of membrane-bound enzymes. In *K. pneumoniae*, cell growth and glycerol assimilation are severely inhibited by high concentrations of PDO or 3-HP. Furthermore, enzymes that play major roles in 3-HP and PDO production, such as DhaB, DhaT, and AldH, are inhibited by 3-HP and PDO (Kumar et al. 2013b). Furthermore, *K. pneumoniae* is

sensitive to 24 g/L acetic acid, 26 g/L lactic acid, and 17 g/L ethanol under aerobic conditions and 15, 19, and 26 g/L, respectively, under anaerobic condition (Celińska, 2012; Cheng et al., 2005). The inhibition of pathways by target product(s) and/or unwanted byproducts often places a limit on the maximum attainable titer of bulk chemicals.

Several studies have reported engineered improvements in microbial tolerance against biochemicals and biofuels, especially in *E. coli*. Modifications of exporter proteins, heat shock proteins, membrane composition, and stress responses have been attempted. Moreover, *in situ* recovery of target products has been attempted for hydrophobic targets. Some tolerant strains have been developed, but unfortunately, without appreciable increases in productivity (Dunlop, 2011; Dunlop et al., 2011; Jarboe et al., 2011). Such efforts to improve tolerance to chemicals have not been reported for *K. pneumoniae*.

#### **4.4 Availability of glycerol**

The global market for glycerol is known to be unpredictable and complex. Glycerol in the current market mainly is from the biodiesel industry, and its supply is directly affected by the use of biodiesel. In contrast, glycerol demand depends on many industries that use it as feedstock. Between 2001 and 2011, the price varied significantly, from \$0.70/kg to \$1.70/kg for pure glycerol and from \$0.04/kg to \$0.33/kg for crude glycerol (Ayoub and Abdullah, 2012; Ciriminna et al., 2014; Kong et al., 2016; Quispe et al., 2013). Nevertheless, glycerol prices, for pure as well as crude, have come down significantly since 2004, when many biodiesel production plants initiated operation. The countries producing biodiesel in large quantities (in million liters) include Germany (2900), Brazil (2300), Argentina (2100), France (2000), USA (1200), Spain (1100), Italy (800), Indonesia (700), and Thailand (600) (Kong et al., 2016). The Asia-Pacific region overtook Europe as the largest glycerol market in the world in 2009 and has remained so ever since. Market growth is driven by increases in the

uses for glycerol in such sectors as pharmaceuticals, personal hygiene, and food and beverage production (Quispe et al., 2013).

The production of crude glycerol is expected to reach 6 million tons by 2025 (Ciriminna et al., 2014). At present, the surplus of glycerol from biodiesel and oleochemicals is assumed to be sufficient to meet these new demands. However, the scenario may change in the future with rapid growth of glycerol-based industries and a limited supply of crude glycerol. For example, if 3-HP produced from glycerol were used as a substitute for the chemical acrylic acid (global production, 5.85 million tons as of 2014), approximately 7 million tons of glycerol/year would be needed. If we include other platform chemicals such as PDO, BDO, and ethanol, the demand will be even higher, much more than the expected supply. The growth of glycerol-based biotechnology will eventually be constrained by the supply and price of crude glycerol as a carbon source.

## **5. Concluding remarks**

Glycerol is an excellent substrate for the production of biochemicals and biofuels. Although the biodiesel industry has grown rather slowly in recent years, crude glycerol is still cheap and generated in large quantities. *K. pneumoniae* has already been successfully employed for the production of PDO, 3-HP, BDO, lactic acid, and ethanol from glycerol at industrially attractive levels. Other metabolites such as acetic acid, pyruvic acid, malic acid, fumaric acid, alanine, *n*-butanol, 1,2-propanediol, propionic acid, formic acid, and hydrogen can also be produced from glycerol by *K. pneumoniae*, although these have not yet been extensively explored. At present, sugar-based bioprocesses are prevalent, and no bioprocesses employing *K. pneumoniae* and glycerol are commercially available. The pathogenicity of *K. pneumoniae* and other technical issues are existing challenges to its use. However, advancements in the fields of metabolic engineering, synthetic biology, systems biology, and evolutionary engineering will enable us to further exploit the advantageous nature of *K.*

1000 *pneumoniae* as a biocatalyst and glycerol as a carbon source. A better understanding of  
1001 glycerol metabolism in *K. pneumoniae* and improvements in its performance as a microbial  
1002 cell factory should make this organism an attractive alternative to current sugar-based  
1003 methods for the production of biochemicals and biofuels.

#### 1004 **Acknowledgement**

1005 This study was supported financially by the Advanced Biomass R&D Center (ABC) of  
1006 Global Frontier Project funded by the Korean Ministry of Science, ICT and Future planning  
1007 (ABC-2011-0031361).

## References

- Abdel-Hamid, A., Attwood, M., Guest, J., 2001. Pyruvate oxidase contributes to the aerobic growth efficiency of *Escherichia coli*. *Microbiology* 147, 1483–1498.
- Abdel-Rahman, M.A., Tashiro, Y., Sonomoto, K., 2013. Recent advances in lactic acid production by microbial fermentation processes. *Biotechnol. Adv.* 31, 877–902.
- Abdel-Rahman, M.A., Sonomoto, K., 2016. Opportunities to overcome the current limitations and challenges for efficient microbial production of optically pure lactic acid. *J. Biotech.* 236, 176–192.
- Albertí, S., Marqués, G., Camprubi, S., Merino, S., Tomás, J.M., Vivanco, F., Benedí, V.J., 1993. C1q binding and activation of the complement classical pathway by *Klebsiella pneumoniae* outer membrane proteins. *Infect. Immun.* 61, 852–860.
- Arasu, M.V., Kumar, V., Ashok, S., Song, H., Rathnasingh, C., Lee, H.J., Seung, D., Park, S., 2011. Isolation and characterization of the new *Klebsiella pneumoniae* J2B strain showing improved growth characteristics with reduced lipopolysaccharide formation. *Biotechnol. Bioprocess Eng.* 16, 1134–1143.
- Årsköld, E., Lohmeier-Vogel, E., Cao, R., Roos, S., Rådström, P., van Niel, E.W.J., 2008. Phosphoketolase Pathway Dominates in *Lactobacillus reuteri* ATCC 55730 Containing Dual Pathways for Glycolysis. *J. Bacteriol.* 190:206-212.
- Ashok, S., Raj, S.M., Rathnasingh, C., Park, S., 2011. Development of recombinant *Klebsiella pneumoniae*  $\Delta dh a T$  strain for the co-production of 3-hydroxypropionic acid and 1,3-propanediol from glycerol. *Appl. Microbiol. Biotechnol.* 90, 1253–1265.
- Ashok, S., Sankaranarayanan, M., Ko, Y., Jae, K.-E., Ainala, S.K., Kumar, V., Park, S., 2013a. Production of 3-hydroxypropionic acid from glycerol by recombinant *Klebsiella pneumoniae*  $\Delta dh a T \Delta y q h D$  which can produce vitamin B<sub>12</sub> naturally. *Biotechnol. Bioeng.* 110(2), 511–524.
- Ashok, S., Raj, S.M., Ko, Y., Sankaranarayanan, M., Zhou, S., Kumar, V., Park, S., 2013b. Effect of *puuC* overexpression and nitrate addition on glycerol metabolism and anaerobic 3-

1038 hydroxypropionic acid production in recombinant *Klebsiella pneumoniae*  $\Delta glpK\Delta dhaT$ .  
1039 Metab. Eng. 15, 10–24.

1040 Ayoub, M., Abdullah, A.Z., 2012. Critical review on the current scenario and sig- nificance  
1041 of crude glycerol resulting from biodiesel industry towards more sustainable renewable  
1042 energy industry. Renew. Sustain. Energy Rev. 16, 2671–2686.

1043 Bächler, C., Schneider, P., Bähler, P., Lustig, A., Erni, B., 2005. *Escherichia*  
1044 *coli* dihydroxyacetone kinase controls gene expression by binding to transcription factor  
1045 DhaR. EMBO J. 24, 283-293.

1046 Booth, I.R., Ferguson, G.P., Miller, S., Li, C., Gunasekera, B., Kinghorn, S., 2003. Bacterial  
1047 production of methylglyoxal: a survival strategy or death by misadventure? Biochem Soc  
1048 Trans. 31, 1406–1408.

1049 Booth, I.R., 2005. Glycerol and methylglyoxal metabolism. In: (Ed.) Bock, A., Curtiss III, R.,  
1050 Kaper, J. B., Neidhardt, F. C., Nystrom, T., Rudd, K. E., Squires, C. L., EcoSal—*Escherichia*  
1051 *coli* and *Salmonella*: cellular and molecular biology. ASM press, Washington, D.C.

1052 Branduardi, P., de Ferra, F., Longo, V., Porro, D., 2014. Microbial *n*-butanol production from  
1053 Clostridia to non-Clostridial hosts. Eng. Life Sci. 14, 16–26.

1054 Buck, M., Gallegos, M.T., Studholme, D.J., Guo, Y., Gralla, J.D., 2000. The bacterial  
1055 enhancer-dependent  $\sigma^{54}$  ( $\sigma^N$ ) transcription factor. J. Bacteriol. 182, 4129-4136.

1056 Cabelli, V.J., 1955. The tricarboxylic acid cycle in the oxidative and synthetic metabolism of  
1057 *Klebsiella pneumoniae*. J. Bacteriol. 70, 23-29.  
1058

1059 Causey, T.B., Shanmugam, K.T., Yomano, L.P., Ingram, L.O., 2004. Engineering  
1060 *Escherichia coli* for efficient conversion of glucose to pyruvate. Proc Natl Acad Sci USA  
1061 101, 2235–2240.  
1062

1063 Celińska, E., Grajek, W., 2009. Biotechnological production of 2,3-butanediol—current state  
1064 and prospects. Biotechnol. Adv. 27, 715–725.

1065 Celińska, E., 2012. *Klebsiella* spp. as a 1, 3-propanediol producer—the metabolic  
1066 engineering approach. Crit. Rev. Biotechnol. 32, 274–288.

1067 Chao, Y.-P., Liao, J.C., 1993. Alteration of growth yield by overexpression of  
 1068 phosphoenolpyruvate carboxylase and phosphoenolpyruvate carboxykinase in *Escherichia*  
 1069 *coli*. Appl. Environ. Microbiol. 59, 4261–4265.

1070

1071 Chen, Z., Liu, H.J., Liu, D.H., 2009. Regulation of 3-hydroxypropionaldehyde accumulation  
 1072 in *Klebsiella pneumoniae* by overexpression of *dhaT* and *dhaD* genes. Enzyme Microb.  
 1073 Technol. 45 (4), 305–309.

1074 Chen, Z., Liu, H., Liu, D., 2011. Metabolic pathway analysis of 1,3-propanediol production  
 1075 with a genetically modified *Klebsiella pneumoniae* by overexpressing an endogenous  
 1076 NADPH-dependent alcohol dehydrogenase. Biochemical Eng. J. 54, 151–157.

1077 Cheng, K.K., Liu, H.J., Liu, D.H., 2005. Multiple growth inhibition of *Klebsiella pneumoniae*  
 1078 in 1,3-propanediol fermentation. Biotechnol. Lett. 27, 19–22.

1079 Cheng, K.K., Zhang, J.A., Liu, D.H., Sun, Y., Liu, H.-J., Yang, M.-D., Xu, J.M., 2007. Pilot-  
 1080 scale production of 1,3- propanediol using *Klebsiella pneumoniae*. Process Biochem. 2007;  
 1081 42, 740–744.

1082 Cheng, K.K., Wu, J., Wang, G.Y., Li, W.Y., Feng, J., Zhang, J.A., 2013. Effects of pH and  
 1083 dissolved CO<sub>2</sub> level on simultaneous production of 2,3-butanediol and succinate using  
 1084 *Klebsiella pneumoniae*. Bioresour. Technol. 135:500–503.

1085 Chu, H.S., Kim, Y.S., Lee, C.M., Lee, J.H., Jung, W.S., Ahn, J.-H., Song, S.H., Choi, I.S.,  
 1086 Cho, K.M., 2015. Metabolic engineering of 3-hydroxypropionic acid biosynthesis in  
 1087 *Escherichia coli*. Biotechnol. Bioeng. 112(2), 356-364.

1088 Chun, A., Yunxiao, L., Ashok, S., Seol, E., Park, S., 2014. Elucidation of toxicity of organic  
 1089 acids inhibiting growth of *Escherichia coli* W. Biotechnol. Bioprocess Eng. 19, 858–865.

1090 Ciriminna, R., Pina C.D., Rossi, M., Pagliaro, M., 2014. Understanding the glycerol market.  
 1091 Eur. J. Lipid Sci. Technol. 116, 1432–1439.

1092 Clomburg, J.M., Gonzalez, R., 2013. Anaerobic fermentation of glycerol: a platform for  
 1093 renewable fuels and chemicals. Trends Biotechnol. 31(1), 20-28.

1094 da Silva, G.P., Mack, M., Contiero, J., 2009. Glycerol: A promising and abundant carbon  
 1095 source for industrial microbiology. Biotechnol. Adv. 27 (1), 30–39.



1096 De Mey, M., De Maeseneire, S., Soetaert, W., Vandamme, E., 2007. Minimizing acetate  
1097 formation in *E. coli* fermentations. J. Ind. Microbiol. Biotechnol. 34, 689-700.  
1098

1099 Dharmadi, Y., Murarka, A., Gonzales, R., 2006. Anaerobic fermentation of glycerol by  
1100 *Escherichia coli*: a new platform for metabolic engineering. Biotechnol. Bioeng. 94, 821–829.

1101 Dugar, D., Stephanopoulos, G., 2011. Relative potential of biosynthetic pathways for biofuels and  
1102 bio-based products. Nature Biotechnol. 29, 1074–1078.

1103 Dunlop, M.J., 2011. Engineering microbes for tolerance to next generation biofuel.  
1104 Biotechnol Biofuels 4:32.

1105 Dunlop, M.J., Dossani, Z.Y., Szmidt, H.L., Chu, H.C., Lee, T.S., Keasling, J.D., Hadi, M.Z.,  
1106 Mukhopadhyay, A., 2011. Engineering microbial biofuel tolerance and export using efflux  
1107 pumps. Mol. Syst. Biol. 7:487.

1108 Durgapal, M., Kumar, V., Yang, T.H., Lee, H.J., Seung, D., Park S., 2014. Production of 1,3-  
1109 propanediol from glycerol using the newly isolated *Klebsiella pneumoniae* J2B. Bioresour.  
1110 Technol. 159, 223–231.

1111 Durnin, G., Clomburg, J., Yeates, Z., Alvarez, P.J., Zygorakis, K., Campbell, P., Gonzalez,  
1112 R., 2008. Understanding and harnessing the microaerobic metabolism of glycerol  
1113 in *Escherichia coli*. Biotechnol. Bioeng. 103, 148–161.

1114 El Fertas-Aissani, R., Messai, Y., Alouache, S., Bakour, R., 2013. Virulence profiles and  
1115 antibiotic susceptibility patterns of *Klebsiella pneumoniae* strains isolated from different  
1116 clinical specimens. Pathol. Biol. 61(5), 209-216.

1117 Eppler, T., Boos, W., 1999. Glycerol-3-phosphate-mediated repression of *maltT* in  
1118 *Escherichia coli* does not require metabolism, depends on enzyme IIAGlc and is mediated by  
1119 cAMP levels. Mol. Microbiol. 33, 1221–1231.  
1120

1121 Eppler, T., Postma, P., Schutz, A., Volker, U., Boos, W., 2002. Glycerol-3-phosphate-  
1122 induced catabolite repression in *Escherichia coli*. J. Bacteriol., 184, 3044-3052.

1123 Fang, C.T., Chuang, Y.P., Shun, C.T., Chang, C.S., Wang, J.T., 2004. A novel virulence gene  
1124 in *Klebsiella pneumoniae* strains causing primary liver abscess and septic metastatic  
1125 complications. J. Exp. Med. 199, 697–705.

1126 Feng, X., Ding, Y., Xian, M., Xu, X., Zhang, R., Zhao, G., 2014. Production of optically pure  
 1127 D-lactate from glycerol by engineered *Klebsiella pneumoniae* strain. *Bioresour. Technol.* 172,  
 1128 269–275.

1129 Forage, R.G., Foster, M.A., 1982. Glycerol fermentation in *Klebsiella pneumoniae*: functions  
 1130 of the coenzyme B12-dependent glycerol and diol dehydratases. *J. Bacteriol.* 149, 413–419.

1131 Garcia-Alles, L.F., Siebold, C., Nyffeler, T.L., Flukiger-Bruhwiller, K., Schneider, P., Burgi,  
 1132 H.-B., Baumann, U., Erni, B., 2004. Phosphoenolpyruvate- and ATP-dependent  
 1133 dihydroxyacetone kinases: covalent substrate-binding and kinetic mechanism. *Biochemistry*  
 1134 43, 13037–13045.

1135 Gutknecht, R., Beutler, R., Garcia-Alles, L., Baumann, U., Erni, B., 2001. The dihydro-  
 1136 xyacetone kinase of *Escherichia coli* utilizes a phosphoprotein instead of ATP as phosphoryl  
 1137 donor. *EMBO J.* 20, 2480–2486.

1138 Hao, J., Wang, W., Tian, J., Li, J., Liu, D., 2008. Decrease of 3-hydroxypropionaldehyde  
 1139 accumulation in 1,3-propanediol production by over-expressing *dhaT* gene in *Klebsiella*  
 1140 *pneumoniae* TUAC01. *J. Ind. Microbiol. Biotechnol.* 35, 735–741.

1141

1142 Heller, K.B., Lin, E.C., Wilson, T.H., 1980. Substrate specificity and transport properties of  
 1143 the glycerol facilitator of *Escherichia coli*. *J. Bacteriol.* 144, 274–278.

1144

1145 Hogema, B. M., Arents, J. C., Bader, R., Eijkemans, K., Yoshida, H., Takahashi, H., Aiba,  
 1146 H., Postma, P. W., 1998. Inducer exclusion in *Escherichia coli* by non-PTS substrates: the  
 1147 role of the PEP to pyruvate ratio in determining the phosphorylation state of enzyme IIAGlc.  
 1148 *Mol. Microbiol.* 30, 487–498.

1149

1150 Holtman, C. K., Pawlyk, A. C., Meadow, N. D., Pettigrew, D. W., 2001. Reverse genetics of  
 1151 *Escherichia coli* glycerol kinase allosteric regulation and glucose control of glycerol  
 1152 utilization in vivo. *J. Bacteriol.* 183, 3336–3344.

1153

1154 Horng, Y.T., Chang, K.C., Chou, T.C., Yu, C.J., Chien, C.C., Wei, Y.H., Soo, P.C., 2010.  
 1155 Inactivation of *dhaD* and *dhaK* abolishes by-product accumulation during 1,3-propanediol  
 1156 production in *Klebsiella pneumoniae*. *J. Ind. Microbiol. Biotechnol.* 37(7), 707–716.

1157

1158 Huang, Y., Li, Z., Shimizu, K., Ye, Q., 2012. Simultaneous production of 3-  
1159 hydroxypropionic acid and 1,3-propanediol from glycerol by a recombinant strain of  
1160 *Klebsiella pneumoniae*. *Bioresour. Technol.* 103, 351–359.

1161

1162 Huang, Y., Li, Z., Shimizu, K., Ye, Q., 2013a. Co-production of 3-hydroxypropionic acid and  
1163 1,3-propanediol by *Klebsiella pneumoniae* expressing *aldH* under Microaerobic conditions.  
1164 *Bioresour. Technol.* 128, 505–512.

1165

1166 Huang, C.-F., Jiang, Y.-F., Guo, G.-L., Hwang, W.-S., 2013b. Method of 2,3-butanediol  
1167 production from glycerol and acid-pretreated rice straw hydrolysate by newly isolated strains:  
1168 Pre-evaluation as an integrated biorefinery process. *Bioresour. Technol.* 135, 446–453.

1169

1170 Izquierdo, L., Coderch, N., Pique, N., Bedini, E., Corsaro, M.M., Merino, S., Fresno, S.,  
1171 Tomás, J.M., Regué, M., 2003. The *Klebsiella pneumoniae wabG* gene: role in biosynthesis  
1172 of the core lipopolysaccharide and virulence. *J. Bacteriol.* 185, 7213–7221.

1173 Jarboe, L.R. 2011. YqhD: a broad-substrate range aldehyde reductase with various  
1174 applications in production of biorenewable fuels and chemicals. *Appl. Microbiol.*  
1175 *Biotechnol.* 89, 249–257.

1176 Jarboe, L.R., Liu, P., Royce, L.A., 2011. Engineering inhibitor tolerance for the production of  
1177 biorenewable fuels and chemicals. *Curr. Opin. Chem. Eng.* 1, 38–42.

1178 Jeong, J.-Y., Kim, Y.-J., Cho, N., Shin, D., Nam, T.-W., Ryu, S., Seok YJ., 2004. Expression  
1179 of *ptsG* encoding the major glucose transporter is regulated by *arcA* in *Escherichia coli*. *J.*  
1180 *Biol. Chem.* 279, 38513–38518.

1181

1182 Ji, X.-J., Huang, H., Ouyang, P.-K., 2011. Microbial 2,3-butanediol production: A state-of-  
1183 the-art review. *Biotechnol. Adv.* 29, 351–364.

1184 Jiang, X., Meng, X., Xian, M., 2009. Biosynthetic pathways for 3-hydroxypropionic acid  
1185 production. *Appl. Microbiol. Biotechnol.* 82, 995–1003.

1186 Jung, S.G., Jang, J.H., Kim, A.Y., Lim, M.C., Kim, B., Lee, J., Kim, Y.-R., 2013. Removal of  
1187 pathogenic factors from 2,3-butanediol producing *Klebsiella* species by inactivating virulence  
1188 related *wabG* gene. *Appl. Microbiol. Biotechnol.* 97, 1997–2007.

1189

1190 Jung, W.S., Kang, J.H., Chu, H.S., Choi, I.S., Cho, K.M., 2014. Elevated production of 3-

1191 hydroxypropionic acid by metabolic engineering of the glycerol metabolism in *Escherichia*

1192 *coli*. Metab. Eng. 23, 116–122.

1193 Jung, H.-W., Jung, M.-W., Oh, M.-K., 2015. Metabolic engineering of *Klebsiella*

1194 *pneumoniae* for the production of cis,cis-muconic acid. Appl. Microbiol. Biotechnol. 99,

1195 5217–5225.

1196 Katryniok, B., Paul, S., F. Dumeignil, F., 2013. Recent Developments in the Field of Catalytic

1197 Dehydration of Glycerol to Acrolein. ACS Catal. 3, 1819-1834.

1198 Kim, Y., Ingram, L.O., Shanmugam, K.T., 2008. Dihydrolipoamide dehydrogenase mutation

1199 alters the NADH sensitivity of pyruvate dehydrogenase complex of *Escherichia coli* K-12. J.

1200 Bacteriol. 190, 3851–3858.

1201

1202 Kim, K., Kim, S.K., Park, Y.C., Seo, J.H., 2014a. Enhanced production of 3-

1203 hydroxypropionic acid from glycerol by modulation of glycerol metabolism in

1204 recombinant *Escherichia coli*. Bioresour. Technol. 156, 170-175.

1205 Kim, B., Lee, S., Jeong, D., Yang, J., Oh, M.-K., Lee, J., 2014b. Redistribution of Carbon

1206 Flux toward 2,3-Butanediol Production in *Klebsiella pneumoniae* by Metabolic Engineering.

1207 PLoS One 9(10), e105322.

1208 Ko, Y., Ashok, S., Zhou, S., Kumar, V., Park, S., 2012. Aldehyde dehydrogenase activity is

1209 important to the production of 3-hydroxypropionic acid from glycerol by recombinant

1210 *Klebsiella pneumoniae*. Process Biochem. 47, 1135–1143.

1211

1212 Ko, Y., Ashok, S., Seol, E., Ainala, S., Park, S., 2015. Deletion of putative oxidoreductases

1213 from *Klebsiella pneumoniae* J2B could reduce 1,3-propanediol during the production of 3-

1214 hydroxypropionic acid from glycerol. Biotechnol. Bioprocess Eng. 20, 834–843.

1215

1216 Ko, Y., Seol, E., Sekar, B.S., Kwon, S., Lee, J., Park, S., 2017. Metabolic engineering of

1217 *Klebsiella pneumoniae* J2B for co-production of 3- hydroxypropionic acid and 1,3-

1218 propanediol from glycerol: Reduction of acetate and other by-products. Bioresour. Technol.

1219 244, 1096–1103.

1220 Kong P.S., Aroua M.K., Daud, W.M.A.D. 2016. Conversion of crude and pure glycerol into  
1221 derivatives: A feasibility evaluation. *Renew. Sustain. Energy Rev.* 63, 533-555.

1222 Kraus, G.A., 2008. Synthetic methods for the preparation of 1, 3-propanediol. *Clean* 36 (8),  
1223 648–651.

1224 Kumar, V., Mugesh, S., Jae, K., Durgapal, M., Ashok, S., Ko, Y., Sarkar, R., Park, S., 2012.  
1225 Co-production of 3-hydroxypropionic acid and 1,3-propanediol from glycerol using resting  
1226 cells of *Klebsiella pneumoniae* J2B strain with overexpression of KGSADH. *Appl. Microbiol.*  
1227 *Biotechnol.* 96, 373–383.

1228 Kumar, V., Ashok, S., Park, S., 2013a. Recent advances in biological production of 3-  
1229 hydroxypropionic acid. *Biotechnol. Adv.* 31(6), 945–961

1230 Kumar, V., Mugesh, S., Durgapal, M., Zhou, S., Ko, Y., Ashok, S., Sarkar, R., Park, S.,  
1231 2013b. Simultaneous production of 3-hydroxypropionic acid and 1,3-propanediol from  
1232 glycerol using resting cell system of D-lactic acid deficient mutant of *Klebsiella pneumoniae*  
1233 with overexpression of KGSADH. *Bioresour. Technol.* 135, 555-563.

1234 Kumar, V., Durgapal, M., Sankaranarayanan, M., Somasundar, A., Rathnasingh, C., Song,  
1235 H., Seung, D., Park, S., 2016. Effects of mutation of 2,3-butanediol formation pathway on  
1236 glycerol metabolism and 1,3-propanediol production by *Klebsiella pneumoniae* J2B.  
1237 *Bioresour. Technol.* 214, 432–440.

1238 Lama, S., Ro, S. M., Seol, E., Sekar, B.S., Ainala, S.K., Thangappan, J., Song, H., Seung,  
1239 D., Park, S., 2015. Characterization of 1,3-propanediol oxidoreductase (DhaT)  
1240 from *Klebsiella pneumoniae* J2B. *Biotechnol. Bioproc. Eng.* 20: 971–979.

1241 Lama, S. , Seol, E., Park, S., 2017. Metabolic engineering of *Klebsiella pneumoniae* J2B for  
1242 the production of 1,3-propanediol from glucose. *Bioresour. Technol.* doi:  
1243 10.1016/j.biortech.2017.05.052.

1244 Lee, C.S., Aroua M.K. Aroua, Daud, W.M.A.W., Cognet, P., Pérès-Lucchese, Y., Fabre, P.-  
1245 L., Reynes, O., Latapie, L., 2015. A review: Conversion of bioglycerol into 1,3-propanediol  
1246 via biological and chemical method. *Renew. Sustain. Energy Rev.* 42, 963-972.

1247 Li, Y., Wang, X., Ge, X. & Tian, P., 2016. High production of 3-hydroxypropionic acid  
1248 in *Klebsiella pneumoniae* by systematic optimization of glycerol metabolism. *Sci. Rep.* 6.  
1249 26932.

1250 Liao, D.-I., Reiss, L., Turner, Jr. I., Dotson, G., 2003. Structure of Glycerol Dehydratase  
1251 Reactivase: A New Type of Molecular Chaperone. *Structure* 11, 109–119.

1252 Lim, H.G., Noh, M.H., Jeong, J.H., Park, S., Jung, G.Y., 2016. Optimum rebalancing of the  
1253 3-hydroxypropionic acid production pathway from glycerol in *Escherichia coli*. *ACS Synth.*  
1254 *Biol.* 5 (11), 1247–1255.

1255

1256 Lin, E. C., 1976. Glycerol dissimilation and its regulation in bacteria. *Annu. Rev. Microbiol.*  
1257 30, 535-578.

1258

1259 Lin, E. C. C., 1987. *Escherichia coli* and *Salmonella typhimurium*: Cellular and Molecular  
1260 Biology. In: Neidhardt, F. C. (Ed.), Vol. 1. American Society for Microbiology, Washington,  
1261 D. C, pp. 244-284.

1262

1263 Lin, H., Castro, N.M., Bennett, G.N., San, K.-Y., 2006. Acetyl-CoA synthetase  
1264 overexpression in *Escherichia coli* demonstrates more efficient acetate assimilation and lower  
1265 acetate accumulation: a potential tool in metabolic engineering. *Appl. Microbiol. Biotechnol.*  
1266 71(6), 870–874.

1267

1268 Lin, T.-L., Yang, F.-L., Yang, A.-S., Peng, H.-P., Li, T.-L., Tsai, M.-D., Wu, S.-H., Wang, J.-  
1269 T., 2012. Amino acid substitutions of *MagA* in *Klebsiella pneumoniae* affect the biosynthesis  
1270 of the capsular polysaccharide. *PLoS One* 7:e46783.

1271 Liu, H.J., Zhang, D.J., Xu, Y.H., Mu, Y., Sun, Y.Q., Xiu, Z.L., 2007. Microbial production of  
1272 1, 3-propanediol from glycerol by *Klebsiella pneumoniae* under micro-aerobic conditions up  
1273 to a pilot scale. *Biotechnol. Lett.* 29:1281–1285.

1274

1275 Liu, Y., Gallo, A.A., Bajpai, R.K., Chistoserdov, A., Nelson, A., Segura, L., Xu W., 2010.  
1276 The diversity and molecular modelling analysis of B-12 and B-12-independent glycerol  
1277 dehydratases. *Int. J. Bioinform. Res. Appl.* 6(5), 484–507.

1278

1279 Luo, L.H., Seo, J.W., Baek, J.O., Oh, B.R., Heo, S.Y., Hong, W.K., Kim, D.H., Kim, C.H.,  
1280 2011. Identification and characterization of the propanediol utilization protein PduP of  
1281 *Lactobacillus reuteri* for 3-hydroxypropionic acid production from glycerol. Appl. Microbiol.  
1282 Biotechnol. 89, 697–703.

1283 Luo, L.H., Kim, C.H., Heo, S.Y., Oh, B.R., Hong, W.K., Kim, S., Kim, D.H., Seo, J.W.,  
1284 2012. Production of 3-hydroxypropionic acid through propionaldehyde dehydrogenase PduP  
1285 mediated biosynthetic pathway in *Klebsiella pneumoniae*. Bioresour. Technol. 103, 1–6.

1286 Lütke-Eversloh, T., Bahl, H., 2011. Metabolic engineering of *Clostridium acetobutylicum*:  
1287 Recent advances to improve butanol production. Curr. Opin. Biotechnol. 22, 634–647.

1288 Ma, B.B., Xu, X.L., Zhang, G.L., Wang, L.W., Wu, M., Li, C., 2009. Microbial production of  
1289 1,3-propanediol by *Klebsiella pneumoniae* XJPD-Li under different aeration strategies. Appl.  
1290 Biochem. Biotechnol. 152(1), 127–134.

1291 Ma, C., Zhang, L., Dai, J., Xiu, Z., 2010. Relaxing the coenzyme specificity of 1,3-  
1292 propanediol oxidoreductase from *Klebsiella pneumoniae* by rational design. J. Biotechnol.  
1293 146(4), 173–178.

1294 Maervoet, V.E.T., Mey, M.D., Beauprez, J., Maeseneire, S.D., Soetaert, W.K., 2011.  
1295 Enhancing the microbial conversion of glycerol to 1,3-propanediol using metabolic  
1296 engineering. Org. Process Res. Dev. 15, 189–202.

1297 MarketsANDMarkets, 2012. Global 1,3-propanediol (1,3-PDO) market worth \$560 million  
1298 by 2019.

1299 Martins-Pinheiro, M., Lima, W.C., Asif, H., Oller, C.A., Menck, C.F.M., 2016. Evolutionary  
1300 and Functional Relationships of the dha Regulon by Genomic Context Analysis. PLOS  
1301 ONE 11, e0150772.

1302 Mattam, A.J., James, M., Clomburg, J.M., Gonzalez, R., Yazdani, S.S., 2013. Fermentation  
1303 of glycerol and production of valuable chemical and biofuel molecules. Biotechnol. Lett. 35,  
1304 831–842.

1305 Mazumdar, S., Clomburg, J.M., Gonzalez, R., 2010. *Escherichia coli* strains engineered for  
1306 homofermentative production of D-lactic acid from glycerol. Appl. Environ. Microbiol. 76,  
1307 4327–4336.

1308 Mazumdar, S., Blankschien, M.D., Clomburg, J.M., Gonzalez R., 2013. Efficient synthesis of  
 1309 L-lactic acid from glycerol by metabolically engineered *Escherichia coli*. Microb. Cell Fact.  
 1310 12:7.

1311 Meyer, M., Dimroth, P., Bott., M., 2001. Catabolite repression of the citrate fermentation  
 1312 genes in *Klebsiella pneumoniae*: evidence for involvement of the cyclic AMP receptor  
 1313 protein. J. Bacteriol. 183, 5248–5256.

1314

1315 Organisation for Economic Co-operation and Development (OECD), Food and Agriculture  
 1316 Organization of the United Nations (FAO), Biofuels, in OECD-FAO Agricultural Outlook  
 1317 2011–2020, OECD, Paris, 2011, pp. 77–94.

1318

1319 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Joe, M., Park, D.H., Kim, C.H., 2011.  
 1320 Efficient production of ethanol from crude glycerol by a *Klebsiella pneumonia* mutant strain.  
 1321 Bioresour. Technol. 102, 3918-3922.

1322 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Kim, S.H., Park, D.H., Kim, C.H.,  
 1323 2012a. Optimization of culture conditions for 1,3-propanediol production from glycerol using  
 1324 a mutant strain of *Klebsiella pneumoniae*. Appl. Biochem. Biotechnol. 166, 127–137.

1325 Oh, B.R., Seo, J.W., Heo, S.Y., Hong, W.K., Luo, L.H., Kim, S., Kwon, O., Sohn, J.-H., Joe,  
 1326 M.-H., Park, D.-H., Kim, C.H., 2012b. Enhancement of ethanol production from glycerol in a  
 1327 *Klebsiella pneumoniae* mutant strain by the inactivation of lactate dehydrogenase. Process  
 1328 Biochem. 47, 156–159.

1329 Oh, B.R., Heo, S.Y., Lee, S.M., Hong, W.K., Park, J.M., Jung, Y.R., Kim, D.H., Sohn, J.H.,  
 1330 Seo, J.W., Kim, C.H., 2014. Production of 2- butanol from crude glycerol by a genetically-  
 1331 engineered *Klebsiella pneumoniae* strain. Biotechnol. Lett. 36 (1), 57–62.

1332 Park, J.M., Rathnasingh, C., Song H., 2017. Metabolic engineering of *Klebsiella pneumoniae*  
 1333 based on in silico analysis and its pilot-scale application for 1,3-propanediol and  
 1334 2,3-butanediol co-production Jong. J. Ind. Microbiol. Biotechnol. 44, 431–441.

1335 Petrov, K., Petrova, P., 2009. High production of 2,3-butanediol from glycerol by *Klebsiella*  
 1336 *pneumoniae* G31. Appl. Microbiol. Biotechnol. 84, 659–665.



1337 Petrov, K., Petrova, P., 2010. Enhanced production of 2,3-butanediol from glycerol by forced  
1338 pH fluctuations. *Appl. Microbiol. Biotechnol.* 87, 943–949.

1339 Pina, C.D., Falletta, E., Rossi, M., 2011. A green approach to chemical building blocks. The  
1340 case of 3-hydroxypropanoic acid. *Green Chem.* 13, 1624–1632.

1341 Podschun, R., Ullmann, U., 1998. *Klebsiella* spp. as nosocomial pathogens: epidemiology,  
1342 taxonomy, typing methods, and pathogenicity factors. *Clin. Microbiol. Rev.* 11, 589–603.

1343 Quispe C. A. G., Coronado, C.J.R., Carvalho, Jr.J.A., Glycerol: Production, consumption,  
1344 prices, characterization and new trends in combustion. *Renew. Sustain. Energy Rev.* 27, 475–  
1345 493.

1346 Raynaud C., Sarcabal P., Meynial-Salles I., Croux C., Soucaille P. 2003. Molecular  
1347 characterization of the 1,3-propanediol (1,3-PD) operon of *Clostridium butyricum*. *Proc. Natl.*  
1348 *Acad. Sci. USA* 100, 5010–5015.

1349 Raynaud, C., Lee, J., Sarcabal, P., Croux, C., Meynial-Salles, I., Soucaille, P., 2011.  
1350 Molecular characterization of the glycerol-oxidative pathway of *Clostridium butyricum* VPI  
1351 1718. *J. Bacteriol.* 193, 3127–3134.

1352 Reizer, J., Reizer, A., Saier, M.H. Jr., 1993. The MIP family of integral membrane channel  
1353 proteins: sequence comparisons, evolutionary relationships, reconstructed pathway of  
1354 evolution, and proposed functional differentiation of the two repeated halves of the proteins.  
1355 *Crit Rev Biochem. Mol. Biol.* 28:235-257.

1356 Richey, D.P., Lin, E.C., 1972. Importance of facilitated diffusion for effective utilization of  
1357 glycerol by *Escherichia coli*. *J. Bacteriol.* 112, 784-790.

1358 Rossi, D.M., de Souza, E.A., Ayub, M.A., 2013. Biodiesel residual glycerol metabolism by  
1359 *Klebsiella pneumoniae*: pool of metabolites under anaerobiosis and oxygen limitation as a  
1360 function of feeding rates. *Appl. Biochem. Biotechnol.* 169, 1952–1964.

1361 Sanchez, A.M., Bennett, G.N., San, K.Y., 2005. Efficient succinic acid production from  
1362 glucose through overexpression of pyruvate carboxylase in an *Escherichia coli* alcohol  
1363 dehydrogenase and lactate dehydrogenase mutant. *Biotechnol. Prog.* 21, 358–365.

1364

1365 Saxena, R.K., Anand, P., Saran, S., Isar, J., 2009. Microbial production of 1,3-propanediol:  
 1366 Recent developments and emerging opportunities. *Biotechnol. Adv.* 27, 895–913.  
 1367

1368 Seo, M.Y., Seo, J.W., Heo, S.Y., Baek, J.O., Rairakhwada, D., Oh, B.R., Seo, P.S., Choi,  
 1369 M.H., Kim, C.H., 2009. Elimination of by-product formation during production of 1,3-  
 1370 propanediol in *Klebsiella pneumoniae* by inactivation of glycerol oxidative pathway. *Appl.*  
 1371 *Microbiol. Biotechnol.* 84, 527–534.  
 1372

1373 Shrivastav, A., Lee, J., Kim, H.-Y., Kim, Y.-R., 2013. Recent insights in the removal  
 1374 of *Klebsiella* pathogenicity factors for the industrial production of 2,3-butanediol. *J.*  
 1375 *Microbiol. Biotechnol.* 23, 885–896.  
 1376

1377 Stieb, M., Schink, B., 1984. A new 3-hydroxybutyrate fermenting anaerobe, *Ilyobacter*  
 1378 *polytropus*, gen nov, sp nov, possessing various fermentation pathways. *Arch Microbiol.*  
 1379 140(2), 139–146.  
 1380

1381 Sun, J., van den Heuvel, J., Soucaille, P., Qu, Y., Zeng, A., 2003. Comparative genomic  
 1382 analysis of *dha* regulon and related genes for anaerobic glycerol metabolism in bacteria.  
 1383 *Biotechnol. Progr.* 19, 263–272.  
 1384

1385 Sun, Y.Q., Qi, W.T., Teng, H., Xiu, Z.L., Zeng, A.P., 2008. Mathematical modelling of  
 1386 glycerol fermentation by *Klebsiella pneumoniae*: Concerning enzyme-catalytic reductive  
 1387 pathway and transport of glycerol and 1,3-propanediol across cell membrane. *Biochem. Eng.*  
 1388 *J.* 38, 22–32.  
 1389

1390 Sun, D., Yamada, Y., Sato, S., Uedab, W., 2017. Glycerol as a potential renewable raw  
 1391 material for acrylic acid production. *Green Chem.* DOI: 10.1039/C7GC00358G.  
 1392

1393 Taylor, B.L., Zhulin, I.B., 1999. PAS domains: internal sensors of oxygen, redox potential,  
 1394 and light. *Microbiol. Mol. Biol. Rev.* 63, 479-506.  
 1395

1396 Trinh, C.T., Sreenc, F., 2009. Metabolic Engineering of *Escherichia coli* for efficient  
 1397 conversion of glycerol to ethanol. *Appl. Environ. Microbiol.* 75, 6696–6705.  
 1398

1399 Wang, W., Sun, J., Hartlep, M., Deckwer, W.D., Zeng, A.P., 2003. Combined use of  
1400 proteomic analysis and enzyme activity assays for metabolic pathway analysis of glycerol  
1401 fermentation by *Klebsiella pneumoniae*. Biotechnol. Bioeng. 83, 525–536.  
1402

1403 Wang, Y., Tao, F., Xu P., 2014. Glycerol dehydrogenase plays a dual role in glycerol  
1404 metabolism and 2,3-butanediol formation in *Klebsiella pneumoniae*. J. Biol. Chem. 113, 525–  
1405 535.  
1406

1407 Wang, M., Hu, L., Fan, L., Tan, T., 2015. Enhanced 1-Butanol Production in Engineered  
1408 *Klebsiella pneumoniae* by NADH Regeneration. Energ. Fuel 29, 1823–1829.  
1409

1410 Warnecke, T., Gill, R.T., 2005. Organic acid toxicity, tolerance, and production in  
1411 *Escherichia coli* biorefining applications. Microb. Cell Fact. 4, 425.  
1412

1413 Wei, D., Xu, J., Sun, J., Shi, J., Hao, J., 2013. 2-Ketogluconic acid production by *Klebsiella*  
1414 *pneumoniae* CGMCC 1.6366. J. Ind. Microbiol. Biotechnol. 40, 561–570.

1415 Wei., D., Wang, M., Jiang, B., Shi, J., Hao, J., 2014. Role of dihydroxyacetone kinases I and  
1416 II in the *dha* regulon of *Klebsiella pneumoniae*. J. Biotechnol. 177, 13-19.

1417 Weissenborn, D. L., Wittekindt, N., Larson, T.J., 1992. Structure and regulation of the *glpFK*  
1418 operon encoding glycerol diffusion facilitator and glycerol kinase of *Escherichia coli* K-12. J.  
1419 Biol. Chem. 267, 6122–6131.  
1420

1421 Werpy, T, Petersen, G., 2004. Top value added chemicals from biomass, vol 1: results of  
1422 screening for potential candidates from sugars and synthesis gas. US Department of Energy  
1423 [<http://www.osti.gov/bridge>].

1424 Wu, Z., Wang, Z., Wang, G., Tan, T., 2013. Improved 1,3-propanediol production by  
1425 engineering the 2,3-butanediol and formic acid pathways in integrative recombinant  
1426 *Klebsiella pneumoniae*. J. Biotechnol. 168, 194–200.

1427 Xu, X.L., Zhang, G.L., Wang, L.W., Ma, B.B., Li C., 2009a. Quantitative analysis on  
1428 inactivation and reactivation of recombinant glycerol dehydratase from *Klebsiella*  
1429 *pneumoniae* XJPD-Li. J Mol. Catal. B Enzym. 56, 108–114.

1430 Xu, Y.Z., Guo, N.N., Zheng, Z.M., Ou, X.J., Liu, H.J., Liu, D.H., 2009b. Metabolism in 1,3-  
1431 Propanediol Fed-Batch Fermentation by a D-Lactate Deficient Mutant of *Klebsiella*  
1432 *pneumoniae*. Biotechnol. Bioeng. 104, 965–972.

1433 Yang, B., Larson, T.J., 1998. Multiple promoters are responsible for transcription of the  
1434 *glpEGR* operon of *Escherichia coli*. Biochim. Biophys. Acta 1396, 114–126.

1435 Yazdani, S.S., Gonzalez, R., 2007. Anaerobic fermentation of glycerol: a path to economic  
1436 viability for the biofuels industry. Curr. Opin. Biotechnol. 18(3), 213–219.

1437 Zeng, G., Ye, S., Larson, T.J., 1996. Repressor for the sn-glycerol 3-phosphate regulon of  
1438 *Escherichia coli* K-12: primary structure and identification of the DNA-binding domain. J.  
1439 Bacteriol. 178, 7080–7089.

1440 Zhang, Y., Li, Y., Du, C., Liu, M., Cao, Z., 2006. Inactivation of aldehyde dehydrogenase: a  
1441 key factor for engineering 1,3-propanediol production by *Klebsiella pneumoniae*. Metab. Eng.  
1442 8, 578–586.

1443 Zhang, Q.R., Xiu, Z.L., 2009. Metabolic pathway analysis of glycerol metabolism in  
1444 *Klebsilla pneumonia* incorporating oxygen regulatory system. Biotechnol. Prog. 25, 103–115.

1445 Zhao, L., Zheng, Y., Ma, X., Wei, D., 2009. Effects of over-expression of glycerol  
1446 dehydrogenase and 1,3-propanediol oxidoreductase on bioconversion of glycerol into 1,3-  
1447 propandediol by *Klebsiella pneumoniae* under micro-aerobic conditions. Bioprocess Biosyst.  
1448 En. 32(3), 313–320.

1449 Zheng, P., Wereath, K., Sun, J., Van den Heuvel, J., Zeng AP., 2006. Overexpression of  
1450 genes of the *dha* regulon and its effects on cell growth, glycerol fermentation to 1,3-  
1451 propanediol and plasmid stability in *Klebsiella pneumoniae*. Process Biochem. 41, 2160–  
1452 2169.

1453 Zheng, Z.M., Cheng, K.K., Hu, Q.L., Liu, H.J., Guo, N.N., Liu, D., 2008. Effect of culture  
1454 conditions on 3-hydroxypropionaldehyde detoxification in 1,3-propanediol fermentation by  
1455 *Klebsiella pneumoniae*. Biochem. Eng. J. 39(2), 305–310.

1456 Zhu, J.G., Li, S., Ji, X.J., Huang, H., Hu, N., 2009. Enhanced 1,3-propanediol production in  
1457 recombinant *Klebsiella pneumoniae* carrying the gene *yqhD* encoding 1,3-propanediol  
1458 oxidoreductase isoenzyme. W. J. Microbiol. Biotechnol. 25, 1217–1223.

Zhuge, B., Zhang, C., Fang, H., Zhuge, J., Permaul, K., 2010. Expression of 1,3-propanediol oxidoreductase and its isoenzyme in *Klebsiella pneumoniae* for bioconversion of glycerol into 1,3-propanediol. Appl. Microbiol. Biotechnol. 87(6), 2177–2184.

## Figure captions

**Figure 1:** Transformation of glycerol into a number of valuable chemicals. The solid line represent single step while broken one indicate multiple steps.

**Figure 2:** Oxidative metabolism of glycerol in *K. pneumoniae* (Ashok et al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrows. Solid lines indicate single steps; broken lines indicate multiple steps.

**Figure 3:** Metabolic pathways for the production of PDO and 3-HP from glycerol (Ashok et al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrows.

**Figure 4 (A):** Arrangement of genes in the *dha* regulon of *K. pneumoniae*. 1, *dhaK* (ATP-dependent dihydroxyacetone kinase); 2, *dhaK3/dhaM* (large subunit of ATP-dependent dihydroxyacetone kinase); 3, *dhaK2/dhaL* (small subunit of ATP-dependent dihydroxyacetone kinase); 4, *dhaK1/dhaK* (medium subunits of ATP-dependent dihydroxyacetone kinase); 5, *orfW* (hypothetical protein); 6, *dhaD* (glycerol dehydrogenase); 7, *dhaR* (transcription regulator); 8, *orfW/CdAT* [cob(I)yrinic acid a,c-diamide adenosyltransferase]; 9, *orfX/dhaG* (small subunit of reactivating factor of glycerol dehydratase); 10, *dhaT* (1,3-propanediol oxidoreductase); 11, *orfY* (hypothetical protein); 12, *dhaB/B1* (large subunits of glycerol dehydratase); 13, *dhaC/B2* (medium subunit of glycerol dehydratase); 14, *dhaE/B3* (small subunit of glycerol dehydratase); 15, *orfZ/dhaF* (large subunit of reactivating factor of glycerol dehydratase); 16, *glpF* (glycerol uptake facilitator); 17, 22- *hdeB*; 18, *hdeD* (*hdeBD*-genes related to global regulation); 19, *yfdX*; 20, *kvgS*; and 21, *kvgA* ( *kvgS* and *kvgA* encode for a two-component signal transduction system) (Celińska, 2012; Sun et al., 2003; Wei et al., 2014). **(B):** Glycerol dehydratase gene arrangements and reactivation factors in different organisms (Martins-Pinheiro et al., 2016).

**Figure 5:** Metabolic pathway for the synthesis of BDO from glycerol (Ashok et al., 2011; Celińska and Grajek, 2009; Ji et al., 2011). Gene names are shown in italics on arrows. Solid lines indicate single steps; broken lines indicate multiple steps.

1488 **Figure 6:** Biochemical routes for ethanol and acetic acid production from glycerol (Ashok et  
1489 al., 2011; Kumar et al., 2013a). Gene names are shown in italics on arrow. Solid lines  
1490 indicate single steps; broken lines indicate multiple steps.

1491 **Figure 7:** Lactic acid production from glycerol via methylglyoxal and pyruvate routes  
1492 (Ashok et al., 2011; Mazumdar et al., 2010; 2013). Gene names are shown in italics on arrow.  
1493 Solid lines indicate single steps; broken lines indicate multiple steps.

1494 **Fig. S1** Effect of deletion of glycerol transporters, *glpF1* and *glpF2*, on glycerol consumption  
1495 and PDO production in *K. pneumoniae* J2B3.

1496

Figure 1





Figure 2



Key enzymes: 1- Glycerol kinase; 2- Glycerol-3-phosphate dehydrogenase (aerobic); 3- Glycerol-3-phosphate dehydrogenase (anaerobic); 4- Glycerol dehydrogenase; 5- Putative glycerol dehydrogenase; 6- Dihydroxyacetone kinase I; 7- Dihydroxyacetone kinase II; 8- Pyruvate kinase







Figure 3



Key enzymes: 1- Glycerol dehydratase; 2- NADH-dependent PDOR/Propanol dehydrogenase; 3- Hypothetical oxidoreductase; 4- Aldehyde dehydrogenase; 5- Propionaldehyde dehydrogenase; 6- Phosphotransacylase; 7- Propionate kinase

CoA-independent route:  $C_3H_8O_3 + NAD^+ \longrightarrow C_3H_6O_3$  (3-HP) + NADH + H<sup>+</sup>

CoA-dependent route:  $C_3H_8O_3 + NAD^+ + ADP + Pi \longrightarrow C_3H_6O_3$  (3-HP) + NADH + H<sup>+</sup> + ATP

$C_3H_8O_3 + NAD(P)H + H^+ \longrightarrow C_3H_8O_2$  (PDO) + NAD(P)<sup>+</sup> + H<sub>2</sub>O

Figure 4

A)



B)






Figure 5



Key enzymes: 1- α-Acetolactate synthase; 2- α-Acetolactate decarboxylase; 3- Acetoin reductase ; 4-Diacetyl reductase




Figure 6




Key enzymes: 1- Pyruvate oxidase; 2-Pyruvate-formate lyase, putative formate acetyltransferase 3- Pyruvate dehydrogenase complex; 4- Phosphoacetyl transferase  
5- Acetate kinase; 6 and 7- Bifunctional (acetaldehyde/alcohol) dehydrogenase/putative alcohol dehydrogenase; 8-Formate dehydrogenase, part of the formate hydrogenlyase complex





Figure 7



Key enzymes :1-D-Lactate dehydrogenase (fermentative); 2-D-Lactate dehydrogenase (respiratory); 3-Methylglyoxal synthase; 4- Glyoxalase I and II; 5-Glycerol dehydrogenase; 6-Aldehyde dehydrogenase isozymes; 7-L-Lactate dehydrogenase (respiratory).



## Supplementary Figure S1



\**glpF1* designates the gene located at the respiratory *glpFK* operon, while *glpF2* indicates a separate operon in the fermentative *dha* regulon located between *orfZ*/*dhaF* operon and *hdeB* operon (see Fig. 4A). For the growth of *K. pneumoniae* J2B3 which is deleted for *glpK*, *glpD*, *glpA* and *dhaD* (see Fig. 2 for their roles), glucose was added at 100 mM because the strain cannot grow on glycerol as the sole carbon source (Lama et al., 2017)

**Table 1: Comparison of maximum theoretical yield, ATP yield and redox balance during microbiological production of different metabolites from glycerol and glucose\***

<b>Metabolite</b>	<b>Maximum theoretical yield**</b>		<b>ATP generated/Metabolite (mol/mol)</b>		<b>Redox constrained [NAD(P)H generated &amp; consumed/ Metabolite]</b>	
	<b>Glucose (mol/0.5 mol)</b>	<b>Glycerol (mol/mol)</b>	<b>Glucose</b>	<b>Glycerol</b>	<b>Glucose</b>	<b>Glycerol</b>
1,3-propanediol $C_3H_8O_2$	0.75	0.88	-1	0	Yes (0 & 2)	Yes (0 & 1)
3-hydroxypropionic acid $C_3H_6O_3$	1.0	1.17	1/0/-1/-0.33	0/1	No (1 & 1; 2 & 2)	Yes (1 & 0)
2,3-butanediol $C_4H_{10}O_2$	0.55	0.64	2	2	Yes/No (2 & 1/2)	Yes (4 & 1/2)
Ethanol $C_2H_6O$	1.0	1.17	1	1	Yes/No (1/2 & 2)	Yes/No (2/3 & 2)
Succinic acid $C_4H_6O_4$	0.86	1.0	0/1	0/1	Yes (1 & 2)	No (2 & 2)
Lactic acid $C_3H_6O_3$	1.0	1.17	1/-1	1/-1	No (1 & 1; 0 & 0)	Yes (2 & 0; 1 & 0)
Pyruvic acid $C_3H_4O_3$	1.2	1.4	1	1	Yes (1/2 & 0)	Yes (2/3 & 0)
Acetic acid $C_2H_4O_2$	1.5	1.75	1/2	1/2	Yes (1/2 & 0)	Yes (2/3 & 0)
1-Butanol $C_4H_{10}O$	0.5	0.58	2	2	No (4 & 4)	Yes (6 & 4)

\*The variation in ATP yield, NADH generated and consumed is due to formation of metabolite through different routes.

\*\*The theoretical yields of metabolites were calculated on the basis of degree of reduction (Dugar and Stephanopoulos, 2011).

**Table 2: Summary of PDO production from glycerol using different *K. pneumoniae* strains**

Strain	Mode of operation	Carbon source(s)	Aeration condition	Titer (g/L)	Yield (mol/mol)	Productivity (g/L. h)	Reference
<b>Wild type strains</b>							
<i>K. pneumoniae</i> M5a1	Batch	Glycerol	Anaerobic	58.8	0.53	0.92	Cheng et al. 2007
<i>K. pneumoniae</i> XJPD-Li	Fed-batch	Glycerol	Aerobic	65.3	0.56	3.16	Ma et al. 2009
<i>K. pneumoniae</i> AC 15	Fed-batch	Glycerol	Aerobic	74.1	0.62	3.08	Zheng et al. 2008
<i>K. pneumoniae</i> DSM 2026	Fed-batch	Glycerol	Anaerobic	81.0	0.64	3.00	Liu et al. 2007
<i>K. pneumoniae</i> DSM 2026	Fed-batch	Glycerol	Microaerobic	77.0	0.62	2.70	Liu et al. 2007
<i>K. pneumoniae</i> HR526	Fed-batch	Glycerol	Aerobic	95.4	0.48	1.98	Xu et al. 2009b
<i>K. pneumoniae</i> KG1	Fed-batch	Glycerol	Aerobic	98.8	0.51	3.29	Zhao et al. 2009
<b>Mutant strains</b>							
<i>K. pneumoniae</i> HR526 $\Delta$ ldhA	Fed-batch	Glycerol	Aerobic	102.1	0.52	2.13	Xu et al. 2009b
<i>K. pneumoniae</i> Cu $\Delta$ ldhA	Fed-batch	Glycerol	Aerobic	102.7	0.50	1.53	Oh et al. 2012a
<i>K. pneumoniae</i> YMU2 $\Delta$ aldA	Fed-batch	Glycerol	Anaerobic	70.6	0.70	1.1	Zhang et al. 2006
<i>K. pneumoniae</i> YMU2	Fed-batch	Glycerol	Anaerobic	72.2	0.57	2.41	Wu et al. 2013



<i>ΔbudC_fdh</i>							
Recombinant strains (Overexpression of Genes from the Reductive Branch)							
<i>K. pneumoniae</i> TUAC01_ <i>dhaT</i>	Batch	Glycerol	Aerobic	15.0	0.61	1.36	Hao et al. 2008
<i>K. pneumoniae</i> KG1_ <i>dhaT</i>	Fed-batch	Glycerol	Aerobic	90.9	0.64	2.16	Zhao et al. 2009
<i>K. pneumoniae</i> ME- 308_ <i>yqhD</i>	Fed-batch	Glycerol	Microaerobic	67.6	0.62	1.69	Zhu et al. 2009
<i>K. pneumoniae</i> <i>ΔdhaD</i> <i>ΔdhaK ΔdhaT_yqhD</i>	Batch	Glycerol	Aerobic	7.7	0.53	0.26	Seo et al. 2010

**Table 3: Different strategies carried out for 3-HP production through fed-batch mode**

Strain	Carbon source(s)	Aeration condition	Titer* (g/L)	3-HP yield on glycerol (mol/mol)	Productivity (g/L. h)	Reference
<i>K. pneumoniae</i> DSM 2026 $\Delta dhaT\_puuC$	Glycerol	Microaerobic	16.0 (16.8)	0.23	0.67	Ashok et al. 2011
<i>K. pneumoniae</i> WM3 <i>pUC18kan_aldHec</i>	Glycerol	Anaerobic	24.4 (49.3)	0.18	1.02	Huang et al. 2012
<i>K. pneumoniae</i> J2B _ <i>KGSADH</i>	Glycerol	Anaerobic	11.3 (15.9)	0.27	0.94	Kumar et al. 2012
<i>K. pneumoniae</i> $\Delta glpK\Delta dhaT\_puuC$	Glycerol	Anaerobic	22.0 (5.9)	0.30	0.46	Ashok et al. 2013a
<i>K. pneumoniae</i> $\Delta dhaT\Delta yqhD\_dhaB\_puuC$	Glycerol	Aerobic	28.1 (3.3)	0.40	0.58	Ashok et al. 2013b
<i>K. pneumoniae</i> WM3 <i>pUC18kan_aldHec</i>	Glycerol	Microaerobic	48.9 (25.3)	0.41	1.75	Huang et al. 2013a
<i>K. pneumoniae</i> J2B $\Delta ldhA\_KGSADH$	Glycerol	Microaerobic	22.7 (23.4)	0.35	0.38	Kumar et al. 2013b
<i>E. coli</i> W3110 $\Delta ackA\_pta$ $\Delta yqhD\_dhaB\_mutant\ gabD4$	Glucose & glycerol	Aerobic	71.9	-	1.8	Chu et al. 2015
<i>K. pneumoniae</i> DSM 2026 $\Delta ldh1\Delta ldh2\Delta pta\_puuC$ (pTAC)	Glycerol	Microaerobic	83.8 (22.1)	0.54	1.16	Li et al. 2016

\*The values shown in the bracket are the PDO concentration obtained along with 3-HP.

**Table 4: Comparison of BDO, ethanol and lactic acid production by different strains of *K. pneumoniae* and other organisms**

Strain	Mode of operation	Carbon source(s)	Aeration condition	Titer (g/L)	Yield (mol/mol)	Productivity (g/L. h)	Reference
<b>BDO</b>							
<i>K. pneumoniae</i> G31	Fed-batch	Glycerol	Microaerobic	49.2	0.37	0.17	Petrov and Petrova, 2009
<i>K. pneumoniae</i> G31	Fed-batch	Glycerol	Aerobic	70.0	0.40	0.47	Petrov and Petrova, 2010
<i>Klebsiella</i> sp. 2	Fed-batch	Glycerol	Aerobic	24.9	0.40	0.34	Huang et al. 2013b
<i>K. pneumoniae</i> SDM	Fed-batch	Glucose	Aerobic	150.0	0.84	3.93	Ma et al. 2009
<i>K. pneumoniae</i> KCTC2242 $\Delta$ ldhA_budA_budB	Fed-batch	Glucose	Aerobic	90.0	0.76	2.75	Kim et al. 2014b
<b>Ethanol</b>							
<i>K. pneumoniae</i> GEM167 mutant	Fed-batch	Glycerol	Microaerobic	21.5	0.62	0.93	Oh et al. 2011
<i>K. pneumoniae</i> GEM167 mutant $\Delta$ ldhA_pdc_adhII	Fed-batch	Glycerol	Microaerobic	31.0	0.89	1.20	Oh et al. 2012b
<i>E. coli</i> (nine gene knockout strain)	Batch	Glycerol	Microaerobic	18.0	0.90	0.38	Trinh and Srienc, 2009
<b>Lactic acid</b>							
<i>K. pneumoniae</i> DSM 2026 $\Delta$ dhaT_puuC	Fed-batch	Glycerol	Microaerobic	22.7	0.32	0.95	Ashok et al. 2011
<i>K. pneumoniae</i> J2B	Fed-batch	Glycerol	Microaerobic	28.2	0.28	0.39	Kumar et al. 2013b

$\Delta ldhA\_KGSADH$							
<i>K. pneumoniae</i> BLh-1	Fed-batch	Glycerol	Microaerobic	59.0	0.48	1.48	Rossi et al. 2013
<i>K. pneumoniae</i> J2B	Fed-batch	Glycerol	Microaerobic	37.0	0.32	0.77	Durgapal et al. 2014
<i>K. pneumoniae</i> DSMZ	Fed-batch	Glycerol	Microaerobic	47.8	0.39	1.00	Durgapal et al. 2014
<i>K. pneumoniae</i> ATCC25955	Fed-batch	Glycerol	Microaerobic	142.1	0.84	2.96	Feng et al. 2014
$\Delta dhaT\_ \Delta yqhD\_ ldhA$				(D-isomer)			
<i>E. coli</i> MG1655	Fed-batch	Glycerol	Microaerobic	32.0	0.82	0.54	Mazumdar et al. 2010
$\Delta pta\_ \Delta adhE\_ \Delta frdA\_ \Delta dld\_ glpK\_ glpD$				(D-isomer)			
<i>E. coli</i> MG1655	Fed-batch	Glycerol	Microaerobic	50.0	0.92	0.60	Mazumdar et al. 2013
$\Delta pflB\_ \Delta pta\_ \Delta adhE\_ \Delta frdA\_ \Delta mgsA\_ \Delta lldD\_ \Delta ldhA::ldhA\_ glpK\_ glpD$				(L-isomer)			

# Potential and limitations of *Klebsiella pneumoniae* as a microbial cell factory utilizing glycerol as the carbon source

Kumar, Vinod

2017-10-19

Attribution-NonCommercial-NoDerivatives 4.0 International

---

Vinod Kumar, Sunghoon Park, Potential and limitations of *Klebsiella pneumoniae* as a microbial cell factory utilizing glycerol as the carbon source, *Biotechnology Advances*, Volume 36, Issue 1, January–February 2018, Pages 150-167

<https://doi.org/10.1016/j.biotechadv.2017.10.004>

*Downloaded from CERES Research Repository, Cranfield University*