

# Advances in molecular and genomic research to safeguard food and feed supply from aflatoxin contamination

D. Bhatnagar<sup>1\*</sup>, K. Rajasekaran<sup>1</sup>, M. Gilbert<sup>1</sup>, J.W. Cary<sup>1</sup> and N. Magan<sup>2</sup>

<sup>1</sup>US Department of Agriculture, Agricultural Research Service, 1100 Robert E. Lee Boulevard, New Orleans, LA 70124, USA;

<sup>2</sup>Applied Mycology Group, Cranfield University, MK45 4DT, Cranfield, United Kingdom; [deepak.bhatnagar@ars.usda.gov](mailto:deepak.bhatnagar@ars.usda.gov)

Received: 1 December 2017 / Accepted: 26 January 2018

© 2018 Wageningen Academic Publishers

OPEN ACCESS 

REVIEW ARTICLE

## Abstract

Worldwide recognition that aflatoxin contamination of agricultural commodities by the fungus *Aspergillus flavus* is a global problem has significantly benefitted from global collaboration for understanding the contaminating fungus, as well as for developing and implementing solutions against the contamination. The effort to address this serious food and feed safety issue has led to a detailed understanding of the taxonomy, ecology, physiology, genomics and evolution of *A. flavus*, as well as strategies to reduce or control pre-harvest aflatoxin contamination, including (1) biological control, using atoxigenic aspergilli, (2) proteomic and genomic analyses for identifying resistance factors in maize as potential breeding markers to enable development of resistant maize lines, and (3) enhancing host-resistance by bioengineering of susceptible crops, such as cotton, maize, peanut and tree nuts. A post-harvest measure to prevent the occurrence of aflatoxin contamination in storage is also an important component for reducing exposure of populations worldwide to aflatoxins in food and feed supplies. The effect of environmental changes on aflatoxin contamination levels has recently become an important aspect for study to anticipate future contamination levels. The ability of *A. flavus* to produce dozens of secondary metabolites, in addition to aflatoxins, has created a new avenue of research for understanding the role these metabolites play in the survival and biodiversity of this fungus. The understanding of *A. flavus*, the aflatoxin contamination problem, and control measures to prevent the contamination has become a unique example for an integrated approach to safeguard global food and feed safety.

**Keywords:** *Aspergillus flavus*, secondary metabolites, preharvest control, postharvest control, aflatoxin

## 1. Introduction

Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) is a human carcinogen and acutely toxic to humans, pets, livestock and wildlife. Aflatoxins are secondary metabolites produced by the fungus *Aspergillus flavus* in many commodities, both prior to harvest and during storage after harvest, thereby greatly reducing their value and marketability due to the toxic effects of these compounds. Aflatoxins are among the most carcinogenic natural toxins known, and ingestion by animals and human beings can result in diseases, such as liver cancer. Children are particularly affected by aflatoxin exposure, which leads to stunted growth and delayed development (CAST, 2003). In animal models, activation of AFB<sub>1</sub> by mammalian microsomal cytochrome P450 is required for its carcinogenic effect. Cytochrome P450 monooxygenases

in the liver convert AFB<sub>1</sub> to a variety of metabolites of increased polarity, including AFB<sub>1</sub>-8,9-epoxide, the ultimate carcinogen, which binds covalently to guanine residues in DNA resulting in DNA damage, mutations, and ultimately carcinoma (Egner *et al.*, 2003). Aflatoxin contamination is a worldwide problem, especially in the warmer parts of the world. The US Food and Drug Administration prohibits interstate commerce of feed grain containing more than 20 g/kg AFB<sub>1</sub> and prohibits the sale of milk and eggs containing more than 0.5 g/kg aflatoxin M<sub>1</sub>. Researchers from around the world have attempted to measure the impact of this contamination on the economics of agriculture, as well as the food and feed safety issues (reviewed in a special issue of *World Mycotoxin Journal* by Brown *et al.*, 2015). So far, no completely effective control strategies are available to prevent aflatoxin accumulation in

the field when conditions are favourable for the fungus. A complete understanding of the ecology of the fungus, host plant-*A. flavus* interaction, and aflatoxin contamination processes will help in the development of new control measures aimed at interrupting the mechanisms responsible for preharvest aflatoxin contamination with the goal of producing a safer, economically viable food and feed supply. Numerous investigations, as reviewed in this article, have attempted to gain a better understanding of the relationship between the host plant and the invading fungus, *A. flavus*, because this relationship affects the aflatoxin contamination process. It is now hypothesised that several host plant and fungal genes are probably involved in determining the degree of aflatoxin contamination of the crop, as outlined in this review. More recently, use of biotechnology has enabled researchers to decipher and find solutions much more rapidly to the aflatoxin contamination problem. Even the best laid crop production efforts cannot avoid all preharvest aflatoxin contamination, and proper post-harvest management techniques are required to avoid fungal growth and toxin production. Sorting, drying, shelling, aeration in storage, as well as control of temperature, moisture, and pests all contribute to lower aflatoxin levels and improved food quality at the household, village, and industrial levels. Such a concerted effort to minimise or control aflatoxin contamination will economically benefit the agricultural industry in developed countries, where strict regulatory guidelines prevent the sale of contaminated commodities, but also will increase food safety for populations in developing countries where regulations, if they exist, may not be enforced. Recent research advances to understand biodiversity and ecology of the fungus, host plant-pathogen interaction, genomic and environmental aspects of production of various secondary metabolites, including aflatoxins, and finally the importance of continued postharvest efforts to minimise the aflatoxin contamination to keep the food and feed supply safe for consumption are summarised in this review.

## 2. Biodiversity of *Aspergillus flavus*

*A. flavus* is a soil fungus found in temperate parts of the world. *A. flavus* produces aflatoxins B<sub>1</sub> and B<sub>2</sub> and cyclopiazonic acid (CPA), but is incapable of producing aflatoxins G<sub>1</sub> and G<sub>2</sub>, which are produced by another related *Aspergillus* species, *A. parasiticus*. Only half of the *A. flavus* isolates found in the field are toxigenic. The populations of this fungus are very diverse and are comprised of many vegetative compatibility groups (VCGs), which prevent hyphal fusion across VCGs, but allow genetic exchange within the same VCG. New VCGs are created by chance mutations or through the genetic recombination. As a consequence, isolates of *A. flavus* from different VCGs differ from each other in many ways including enzyme production, virulence characteristics, and aflatoxin-producing ability. *A. flavus* was thought to

exclusively propagate vegetatively through asexual spores (conidia). However, recent studies (Horn *et al.*, 2009) have demonstrated that another type of spore, the ascospore, is produced by the sexual teleomorph (*Petromyces flavus*) of *A. flavus*. Long history of genetic recombination in this fungus is being attributed to sexual recombination in laboratory and limited field studies (Moore *et al.*, 2013). The diversity of this fungus is also described in terms of the sclerotia size these strains produce; the two morphotypes being S (small sclerotia) and L (large sclerotia). Apart from sclerotia size, the S-group isolates produce more sclerotia and less conidia than the L-group; the two types of isolates respond differently to pH and grow differently in light and dark conditions. An unnamed taxon isolated from West Africa called S<sub>BG</sub>, which is phylogenetically divergent from, but morphologically similar to the S-type *A. flavus*, that produces small sclerotia and large amounts of both B- and G-aflatoxins has been identified (Atehnkeng *et al.*, 2008). However, further genomic analysis of S<sub>BG</sub> strains should clarify their exact taxonomic or possible hybrid status.

## 3. Ecology of *Aspergillus flavus*

The ecology of *A. flavus* is directly related to the life cycle of the fungus and the influence that the environment has on the saprophytic and pathogenic phases. The most important factors for the former phase are the soil water potential and temperature, the relative organic matter content and the presence of crop residue. These all govern the inoculum potential of this soil-borne fungus. *A. flavus* survives as sclerotia and mycelium on crop debris, and under conducive conditions this facilitates mycelial growth and sporulation from either sclerotia or asexual conidial formation. The conidia from these sources are spread by insects and wind to ripening crops, e.g. maize, where pathogenic infection phase is initiated on ripening maize cobs during silking. These phases are all influenced by the prevailing water availability (water potential in soil or water activity [a<sub>w</sub>] in crop debris and ripening crops) and temperature. pH optimum for colonisation of crop debris is in the range of 5 to 6.5 (Griffin *et al.*, 2001). In addition, soil nutritional content will interact with environmental conditions. Thus, a significant positive correlation between organic matter content of soil and the isolation of *A. flavus* populations was demonstrated (Zablotowicz *et al.*, 2007). In the life cycle of *A. flavus*, the optimum and range of interacting environmental conditions necessary for sporulation, germination, and fungal growth are thus important. This needs to be compared with those which allow aflatoxin production.

Germination of conidia of *A. flavus* in terms of germination rate and percentage viable germinating spores is significantly influenced by a<sub>w</sub> × temperature conditions. Marín *et al.* (1998) showed that *A. flavus* conidia can germinate rapidly and linearly at between 0.995-0.90 a<sub>w</sub> at 25-30 °C. At ≤0.85 a<sub>w</sub> this is much slower. Overall, germination of conidia was

maximum at 35–40 °C with practically freely available water, but this was slower at 0.95 and 0.90  $a_w$  (25% germination) and <20% at 0.85  $a_w$  based on a 40 day incubation period on a maize-meal based medium.

In terms of mycelial colonisation, studies by Nesci *et al.* (2004) and Giorni *et al.* (2008a) showed that growth of strains of *A. flavus* was more sensitive to matric water stress where soil conditions were simulated than to solute stress. Thus, growth was optimum at -2.8 to -7.0 MPa water potential (= 0.98 to 0.95  $a_w$  at 25/30 °C) in ionic and non-ionic solute potential, while this was optimum at -1.4 to -2.9 MPa (= 0.99–0.98  $a_w$ ) matric potential. *A. flavus* was significantly more sensitive to matric stress with no growth at -14.0 MPa (= 0.90  $a_w$ ) while in solute stress this was at -21.0 MPa (= 0.85  $a_w$ ; Giorni *et al.*, 2008b). The sporulation of *A. flavus* from sclerotia under different  $a_w$  conditions *in vitro* and on maize stalks has been quantified (Giorni *et al.*, 2012). Comparison of artificial (Czapek Dox Agar) and natural (maize stalks) substrates under different temperatures (5–45 °C) and  $a_w$  levels (0.50–0.99) showed that on maize stalks the number of spores/sclerotium was log 4.62 with production initiated after 24 h. The optimal temperature was 20–25 °C on maize stalks (log 4.79 spores/sclerotium). Water stress imposition only reduced sporulation at  $\leq 0.90 a_w$ . With more available water, no significant differences were found in the range 0.90 to 0.99  $a_w$  (Figure 1). Asexual conidial production on maize-based medium suggested that optimum  $a_w \times$  temperature conditions were 30–35 °C at 0.95  $a_w$  (log 7.05 to 7.16 spores/cm<sup>2</sup> colony) where glycerol was used to amend the  $a_w$  levels. At 20 °C there was no sporulation at 0.90  $a_w$  although there was >log 5.75 spores/cm<sup>2</sup> colony at >0.95  $a_w$  (Abdel-Hadi, 2011).

The effect of interacting conditions of  $a_w \times$  temperature is critical in determining the effects on colonisation and AFB<sub>1</sub>

contamination of food matrices such as ripening maize, peanut, tree nuts or spices (Sanchis and Magan, 2004). Recent studies have examined the effect of  $a_w \times$  temperature conditions on growth and AFB<sub>1</sub> production by a type strain of *A. flavus* (NRRL 3357 strain; Abdel-Hadi *et al.*, 2012) (Figure 2). The boundary conditions of germination, growth and toxin differed with the widest/broadest  $a_w \times$  temperature ranges; being wider for germination, than growth and toxin production. Such data were used to develop contour maps of the profiles which indicate the optimum and marginal interacting conditions for growth and for AFB<sub>1</sub> production. This clearly shows that the optimum conditions for growth and for AFB<sub>1</sub> production may be slightly different. At marginal conditions of <20 °C and <0.90  $a_w$  minimal growth and toxin are produced. Of course the interface with the food plant species is critical. For example, in maize there is a period during silking when the  $a_w$  of the ripening maize cobs are ideal for colonisation by *A. flavus*. This is particularly relevant during the dough or R4 stage (0.976  $a_w$ ), and the dent or R5 stage (0.958  $a_w$ ) of maize kernels. Thus any damage to maturing maize cobs during these periods via insects, carrying *A. flavus* spores would lead to rapid colonisation and AFB<sub>1</sub> contamination. Similarly, drought stress in peanut ripening represents conditions where ecologically *A. flavus* is at a competitive advantage, which facilitates colonisation and contamination with the toxin. These types of data sets on the ecology of the life cycle of *A. flavus* can be used to develop models to predict the relative potential risk from toxin contamination based on prevailing weather conditions (Battilani *et al.*, 2013). *A. flavus* is very resilient ecologically and this suggests that under future changes in climate conditions it will perhaps cause more problems in staple food production systems in areas where it was not considered important previously (Medina *et al.*, 2017a).





Figure 1. Sporulation rate of *A. flavus* (MPVP 2062) grown on Czapek agar (CZ) (▲) and maize stalks (□) at (A) different temperatures and (B) at different  $a_w$  levels. The dotted line is the best fit line (from Giorni *et al.*, 2012).





**Figure 2. (A) Contour plot for temperature versus water activity on the growth rate (in mm/d). Numbers on the isopleth lines join conditions of the same growth rate. (B) Contour plot for effect of temperature versus water activity conditions on aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) production (in µg/g). The isopleth lines join conditions at which similar quantities of AFB<sub>1</sub> were produced (from Abdel-Hadi et al. 2012).**

#### 4. Genomics of *Aspergillus flavus* and aflatoxin production

##### Genome sequences and comparative genomics

Comparative genomics, comparing multiple genomes, combined with phenotypic or transcriptomic data has been used in *A. flavus* isolates to elucidate gene function, define evolutionary relationships, secondary metabolite gene function, biotechnology applications, and factors related to pathogenicity (Ehrlich and Mack, 2014; Horn et al., 1996; Moore et al., 2016; Payne et al., 2006). Strain NRRL 3357 (ATCC 200026, GenBank assembly accession: GCA\_000006275.2) was the first fully sequenced *A. flavus* genome (Nierman et al., 2015; Payne et al., 2006) and has been developed as a model organism from studies examining secondary metabolite production (Ehrlich and Mack, 2014; Kim et al., 2008; Rank et al., 2012), proteomics (Pechanova et al., 2013), pathogenesis (Chitarrini et al., 2014; Mylroie et al., 2016), and biological control (Donner et al., 2010). The second major strain sequenced, AF70, was originally isolated from soil obtained from a cotton

field in Yuma Valley, AZ, USA; and due to high AFB<sub>1</sub> levels produced during infection of maize kernels (Cotty 1989) has been used in characterising secondary metabolite production (Ehrlich and Mack, 2014), fungal development (Cary et al., 2015a; Gilbert et al., 2016), pathogen-host interaction (Hruska et al., 2014), genome profiling (Moore et al., 2015), and oxidative stress (Baidya et al., 2014). Recently 14 additional isolates have been sequenced and made publicly available on the National Center for Biotechnology Information website (Faustinelli et al., 2016; Weaver et al., 2017). Nine of these isolates were obtained from peanut seeds in Georgia, USA, three of which do not produce aflatoxin. Eight of these isolates were sequenced as part of a larger project that demonstrated how fingerprinting by genomic insertion/deletion events (indels) across the aflatoxin cluster was used to cluster isolates, which corresponded with aflatoxin production and potentially the amount of aflatoxin produced (Faustinelli et al., 2017).

Comparative genomics of *A. flavus* has been reviewed and reported previously (Cleveland et al., 2009; Georgianna and Payne, 2009; Payne et al., 2006; Rokas et al., 2007; Yu et al., 2007). These early studies revealed the high genomic identity between *Aspergillus oryzae* and *A. flavus*, and that *A. flavus* is enriched in secondary metabolic biosynthesis genes relative to other sequenced Aspergilli. More recent genomic comparison of these strains indicated that some *A. flavus* strains are even more phenotypically and genetically similar to *A. oryzae* than other *A. flavus* strains, considering factors such as amylase production, secondary metabolic profiles, and some secondary metabolic gene cluster sequences (Gibbons et al., 2012). A highly comprehensive comparative genome analysis of 34 ascomycete strains covering 19 Aspergilli strains (De Vries et al., 2017) was recently conducted, which also incorporated extensive phenotypic and expression analysis. These analyses showed that *A. flavus* has distinctly low levels of cellulose-degrading enzymes, and a relatively high number of xylan-degrading enzymes produced, although the presence of regulatory sequences between species is nearly constant. In addition, *A. flavus* exhibits differences in the molecular regulation of conidiation than some other Aspergilli, and the authors report an absence or negligible effect on *A. flavus* spore production under illumination; however, this particular observation is not consistent with previous reports (Calvo et al., 1999).

##### Transcriptomics

Early functional genomics technologies characterising *A. flavus* included microarray analysis of natural antisense transcripts (Smith et al., 2008), transcriptional regulation (Cary et al., 2007), and aflatoxin production (Guo et al., 2003; O'Brian et al., 2003; Wilkinson et al., 2007; Yu et al., 2006, 2007). Later, studies looking at additional secondary metabolic clusters were conducted

that illustrated the transcriptomic response to a range of media and environmental conditions (Georgianna *et al.*, 2010; Reverberi *et al.*, 2013) and the influence of the global regulatory factor *veA* (Cary *et al.*, 2014, 2015a). These studies aided in the identification or further characterisation of gene clusters for piperazines, aflatrem, leporins, aflavarin, asparosone, and aflatoxin, and several of these gene cluster products have yet to be determined (discussed in detail later).


Given the comprehensive and increasingly affordable nature of RNA-sequencing (RNA-seq) technologies, it is not surprising that RNA-seq has replaced many previous genome-wide transcriptomic profiling strategies (Goodwin *et al.*, 2016; 't Hoen *et al.*, 2008; Zhao *et al.*, 2014). A relatively recent report (Ehrlich and Mack, 2014) was among early attempts to utilise RNA-seq to characterise *A. flavus*, where several putative gene clusters were identified through transcriptomic comparison. The use of knockout technology combined with RNA-seq was conducted on the large sclerotia producer *A. flavus* strain AF70 to examine the effects of knocking out the global regulatory zinc-finger transcription factor *nsdC*, which resulted in decreased aflatoxin and aflatrem production, and altered expression of the genes responsible for penicillin, asparosone, and ustiloxin B biosynthesis (Gilbert *et al.*, 2016). Clusters 5, 20, 41, 44 (Georgianna *et al.*, 2010) and others were also affected in the *nsdC* knockout strain. Recently, treatment with chemicals in conjunction with RNA-seq technology has successfully characterised various gene clusters and developmental processes in *A. flavus*, including the aflatoxin cluster. Treatment of *A. flavus* in culture by the naturally occurring phytoalexin and antifungal compound resveratrol resulted in decreased aflatoxin production, and of the 30 genes the authors identify as aflatoxin cluster genes, *aflA* and *aflB* were the only significantly downregulated genes. The regulatory gene *aflR* was not significantly affected, suggesting individual cluster genes could be subjected to independent regulation (Wang *et al.*, 2015). Treatment of *A. flavus* with decanal, another plant volatile, caused 26 of the 55 gene clusters to be affected, a decrease in growth, conidia and sclerotia production, and increase in expression of most aflatoxin cluster genes (Chang *et al.*, 2014). It has been suggested the DNA methyltransferase inhibitor 5-azacytidine interrupts toxosome formation

(Wilkinson *et al.*, 2011), and treatment of *A. flavus* with such an inhibitor resulted in down regulation of genes in the aflatoxin cluster (Lin *et al.*, 2013). However, it has been shown that *A. flavus* does not undergo DNA methylation (Liu *et al.*, 2012), indicating the potential role of a feedback regulatory mechanism in aflatoxin production.

Finally, a 2016 RNA-seq study characterising oxidative stress in *A. flavus* was conducted whereby different strains were subjected to H<sub>2</sub>O<sub>2</sub> exposure. A subset of genes within the aflatoxin cluster (*aflG*, *I*, *K*, *M*, *Ma*, *O*, *P*, *Q*, *V*, *W*, *X*, and *Y*; Figure 3) responded in a dosage dependent manner in two *A. flavus* strains, AF13 and Strain 3357. In addition, some genes in the aflatrem and kojic acid (KA) clusters were affected (Fountain *et al.*, 2016). This again suggests cluster genes can be subject to individual regulation, as *aflR* remained highly expressed in all samples.

### Bioinformatics and aflatoxin production

Analysing RNA-seq data to characterise the regulation of metabolite production, their interaction with the host transcriptome, and broader impacts on fungal gene expression, an array of bioinformatics tools have been applied in functional genomics analysis of *A. flavus* (Cary *et al.*, 2014; Gilbert *et al.*, 2016; Lin *et al.*, 2013; Wang *et al.*, 2015), the most common of which include KEGG (Kyoto Encyclopedia of Genes and Genomes) (Aoki-Kinoshita and Kanehisa, 2007), gene co-expression network analysis (Langfelder and Horvath, 2008), and gene ontology (GO) enrichment analysis (Young *et al.*, 2010). Examples of these strategies applied to *A. flavus* and aflatoxin production include Fountain *et al.* (2016) whereby *A. flavus* was treated with H<sub>2</sub>O<sub>2</sub> to induce oxidative stress and a weighted co-expression analysis identified a network of oxidative stress, fungal growth, and secondary metabolic genes, including aflatoxin production, describing a series of coordinated processes facilitating pathogenesis. In another study, Musungu *et al.* (2016) also applied co-expression analysis, this time to maize kernels infected with *A. flavus*, which were subjected to a dual-genome RNA-seq. They found *aflS* was co-regulated with multiple ROS-related maize genes, suggesting its functioning as a possible 'monitor' for host ROS levels. Finally, the emergence of secondary metabolite and genomic databases, such as ClusterMine360, IMG-



**Figure 3. Schematic of the aflatoxin gene cluster.** Genomic nomenclature at top of aflatoxin cluster represents functionality of genes, while the single letter *afl* nomenclature is also presented (below the horizontal line). The *nadA* gene (grey arrow) is part of the adjacent sugar cluster. Arrows denote direction of transcription.

ABC, MIBiG, and FungiDB, combined with computational strategies like those described here using comparative and computational genomics, transcriptomics and secondary metabolic gene cluster prediction is resulting in more complex data sets providing meaningful insights into fungal and crop biology (Andersen *et al.*, 2013; Nagano *et al.*, 2016; Vesth *et al.*, 2016; Weber and Kim 2016). Tools, such as these applied specifically to *A. flavus*, will continue to aid in understanding aflatoxin regulation, with a goal towards identifying novel targets and strategies for mitigation.

## 5. *Aspergillus flavus* secondary metabolite gene clusters

This section briefly summarises the genetics and biochemistry of secondary metabolite (SM) gene clusters in *A. flavus* with particular emphasis on the aflatoxin biosynthetic gene cluster and those regulatory factors known to be involved in aflatoxin production. In addition, information will be provided on other SM gene clusters and their associated metabolites that have been experimentally identified and characterised in *A. flavus* and their potential biological functions.

Secondary metabolites play many important roles in fungal ecology, such as virulence factors, chemical defence agents, developmental regulators and as compounds that are toxic to humans and livestock. Identification of the form and function of the products of these SM gene clusters will undoubtedly provide invaluable insights into the pathobiology of *A. flavus* as well as other fungal plant pathogens harbouring similar gene clusters. Web-based tools such as the SMURF (Secondary Metabolite Unique Regions Finder) (Georgianna *et al.*, 2010) and antiSMASH (antibiotics and Secondary Metabolite Analysis Shell) (Medema *et al.*, 2011) algorithms have shown that *A. flavus* has the potential to produce numerous other SMs, many whose biological activities are unknown. SMURF analysis predicted that the *A. flavus* genome harbours up to 55 SM gene clusters [56 when including the KA gene cluster identified originally in *A. oryzae* by Marui *et al.* (2011)]. Identification of these clusters is based on the presence of cluster 'backbone' genes, such as non-ribosomal peptide synthases (NRPSs), polyketide synthases (PKSs), PKS-NRPS hybrids, geranylgeranyl pyrophosphate synthases (GGPSs) and dimethylallyltryptophan synthases (DMATs), as well as flanking genes encoding decorating enzymes, transcription factors, and transporters. Although about 40 SMs have been identified from *A. flavus* cultures (Cole *et al.*, 2003; Turner and Aldridge, 1983), metabolites have only been experimentally assigned to 12 *A. flavus* gene clusters. Brief descriptions of the physical, genetic, and biological properties of these twelve SM gene clusters and their associated metabolites are provided below and in Table 1.

## Aflatoxin cluster

The aflatoxin (AF) gene cluster is probably the most extensively studied fungal SM gene cluster and has been the subject of numerous comprehensive reviews (Bhatnagar *et al.*, 2003; Kumar *et al.*, 2016; Roze *et al.*, 2013; Yu, 2012). The AF gene cluster (~70 kb) is located near the telomere on the right arm of chromosome 3 and is composed of 25 genes (Yu *et al.*, 2004; Figure 3). Significant contributions to the elucidation of aflatoxin biosynthesis and its regulation can be attributed to studies in *Aspergillus nidulans*, a model fungus that produces the aflatoxin precursor sterigmatocystin (STE) (Brown *et al.*, 1996; Calvo *et al.*, 2002). Many of the ground-breaking enzymatic and genetic studies on aflatoxin biosynthesis were also performed in the closely related aflatoxin producer, *Aspergillus parasiticus*. Prior to elucidation of the physical and functional genetics of the AF biosynthetic gene cluster, many of the enzymes involved in aflatoxin biosynthesis were identified and characterised (reviewed in Yu, 2012). The genetic makeup of the AF gene cluster was fully characterised before sequencing and annotation of the *A. flavus* genome in large part through genetic complementation of AF biosynthetic mutants using *A. flavus* and *A. parasiticus* cosmid libraries (Yu *et al.*, 1995).

Regulation of aflatoxin biosynthesis is controlled by pathway-specific and globally-acting regulatory factors as well as epigenetic factors (reviewed in Amare and Keller, 2014; Brakhage, 2013; Macheleidt *et al.*, 2016). Annotation of the *A. flavus* genome indicates that about 50% of SM gene clusters do not encode a cluster-specific transcriptional activator suggesting that regulation of these clusters occurs at the global level (Ehrlich and Mack, 2014). A genetic linkage governing both development and secondary metabolism was demonstrated in *A. nidulans* (Adams and Yu, 1998; Hicks *et al.*, 1997) and this is also true for regulation of aflatoxin biosynthesis in *A. flavus*. Discussion of only a select few regulators of aflatoxin biosynthesis will be presented here with emphasis on those not presented in the latest review (Macheleidt *et al.*, 2016).

## Pathway-specific regulatory factors

A Gal4-type binuclear cluster Zn(II)<sub>2</sub>-Cys<sub>6</sub> transcriptional activator encoded by *afIR* regulates expression of most AF/STE cluster genes (Flaherty *et al.*, 1995; Woloshuk *et al.*, 1994; Yu *et al.*, 1996). *A. flavus afIR* deletion mutants do not make aflatoxins and demonstrate loss of AF cluster gene expression while *afIR* overexpressing strains demonstrated increased levels of AF gene cluster transcripts and aflatoxin production. AfIR binds to the palindromic sequence 5'-TCGN5CGA-3' present in the promoter region of most AF biosynthetic genes (Ehrlich *et al.*, 1999; Fernandes *et al.*, 1998). Interestingly, AfIR was also shown to control expression of the *nadA*, *hlyC*, and



Table 1. Features of experimentally characterised *Aspergillus flavus* secondary metabolite gene clusters.

Cluster metabolite <sup>1</sup>	Cluster size (kb)/ chromosome	Backbone gene classification <sup>2</sup>	Cluster-specific transcriptional regulator	Number of genes in cluster <sup>3</sup>	Putative biological function	References
Aflatoxin	66.7/3	PKS	yes	25	antiinsectan/oxidative stress resistance?	Kumar <i>et al.</i> (2017); Roze <i>et al.</i> (2013)
Aflatrem	9.7/7 and 6.9/5	DMATs/GGPPs	no	5 and 3	antiinsectan/antifeedant	Nicholson <i>et al.</i> (2009); Tang <i>et al.</i> (2015)
Aflavarin	15/3	PKS	no	5	antiinsectan/sclerotial development	Cary <i>et al.</i> (2015a)
Asparasone A	15.8/1	PKS	yes	4	antiinsectan/sclerotial pigment/abiotic stress resistance	Cary <i>et al.</i> (2014)
Cyclopiazonic acid	22.9/3	PKS-NRPS	no <sup>4</sup>	5	mammalian antifeedant?	Chang <i>et al.</i> (2009a); Kato <i>et al.</i> (2011)
Ditryptophenaline	14.3/4	NRPS	no	3	?	Saruwatari <i>et al.</i> (2014)
Kojic acid	5.8/5	oxidoreductase	yes	3	insect antifeedant/antioxidant?	Terabayashi <i>et al.</i> (2010)
Leporin	31.6/6	PKS-NRPS	yes	9	insect antifeedant/sclerotial development	Cary <i>et al.</i> (2015b)
Piperazine	15.1/6 and 14.0/8	NRPS-like	no	6 and 6	sclerotial development	Forseth <i>et al.</i> (2013)
Ustiloxin B	27/5	RiPS	yes	16	?	Umemura <i>et al.</i> (2014); Nagano <i>et al.</i> (2016)

<sup>1</sup> Some clusters produce more than one metabolite.

<sup>2</sup> NRPS = non-ribosomal peptide synthetases; PKS = polyketide synthetases; GGPPs = geranylgeranyl pyrophosphate synthetases; DMAT = dimethylallyltryptophan synthetases; RiPS = ribosomal peptide synthetic pathway.

<sup>3</sup> As defined experimentally or as defined in Georgianna *et al.* (2010).

<sup>4</sup> The cyclopiazonic acid (CPA) cluster does contain a gene encoding a putative C6 transcription factor, *cpaR*, but the role of this gene in regulation of CPA cluster gene expression has not been experimentally determined.

*niiA* genes located outside of the AF gene cluster (Price *et al.*, 2006). Aflatoxin biosynthesis is also regulated by *AflS* (*aflI*) located adjacent to and divergently transcribed from *aflR* (Meyers *et al.*, 1998). *AflS* does not present similarity to other proteins of known function and the exact role of *AflS* in aflatoxin biosynthesis is not clear despite intense investigation. Deletion of *aflS* in *A. flavus* did not appear to affect expression of AF cluster genes, however, *aflS* deletion mutants did not produce aflatoxins (Meyers *et al.*, 1998). An *A. flavus* strain overexpressing *aflS* did not elevate transcription of late AF pathway genes *aflM* (*ver-1*) and *aflP* (*omtA*) nor did it increase expression of *aflR* (Du *et al.*, 2007). However, an increase in expression of early and mid-AF pathway genes was observed when overexpressing *aflR* and *aflS*, even more so than when overexpressing *aflR* alone. This suggested that *AflS* modulates the expression of AF early and mid-pathway genes in conjunction with *aflR* (Chang *et al.*, 2002). This is in agreement with earlier yeast two-hybrid assays that showed *AflS* interacts with *AflR* suggesting that an intimate association of *AflS* and

*AflR* is required to fully activate expression of some AF structural genes (Chang, 2003).

### Globally-acting regulatory factors

A heterotrimeric, nuclear-localised regulatory complex consisting of the proteins VeA, LaeA and VelB has been shown to coordinate fungal development and secondary metabolism in a light-dependent manner (Bayram *et al.*, 2008). VeA, was originally characterised as a positive regulator of sexual development and STE production in *A. nidulans* (Kim *et al.*, 2002). Studies in *A. flavus* demonstrated that VeA is required for production of both sclerotia and aflatoxin as well as numerous other secondary metabolites including CPA and aflatrem (Duran *et al.*, 2007). LaeA, is a putative chromatin-modifying protein containing a methyltransferase domain (Reyes-Dominguez *et al.*, 2010). Similar to *veA*, deletion of *laeA* results in abrogation of sclerotia and aflatoxin production and transcriptomic analysis has indicated regulation by LaeA of as many as 24 SM gene clusters (Kale *et al.*, 2008). The

velvet-like protein B (VelB) forms a heterodimer with VeA in the cytosol, migrates into the nucleus in the absence of light, and then interacts with LaeA (Bayram and Braus, 2012). Deletion of *velB* results in reduced conidiation and loss of sclerotia and aflatoxin production (Chang *et al.*, 2013). In *A. flavus*, VeA appears to take a dominant role to velB in control of conidiation. Interestingly, yeast two-hybrid assays showed that both LaeA and VelB interact with FluG, a positive regulator of conidiogenesis, but negative regulator of sclerotial formation in *A. flavus* (Chang *et al.*, 2012). This study showed that FluG does not control secondary metabolism in *A. flavus* though it was shown to be required for STE production in *A. nidulans* (Hicks *et al.*, 1997).

The global transcription factor genes *nsdC* (C<sub>2</sub>H<sub>2</sub>-type) and *nsdD* (GATA-type) control both sclerotial, conidial and aflatoxin production in *A. flavus* (Cary *et al.*, 2012). Inactivation of *nsdC* or *nsdD* in *A. flavus* resulted in loss of sclerotial production while aflatoxin biosynthesis was severely reduced or totally lost in  $\Delta nsdD$  and  $\Delta nsdC$  mutants, respectively. A subsequent RNA-seq study comparing gene expression in an *A. flavus*  $\Delta nsdC$  mutant compared to a control revealed differential expression of genes associated with SM production including down-regulation of genes involved in aflatoxin biosynthesis but upregulation of penicillin biosynthetic cluster genes (Gilbert *et al.*, 2016). A recently identified global regulator, *hbx1*, encoding a homeobox transcription factor was shown to be required for production of aflatoxins, CPA and aflatoxin in *A. flavus* (Cary *et al.*, 2017). Hbx1 was also shown to be essential for conidial and sclerotial production making it the first transcription factor identified in *A. flavus* required for production of conidia, sclerotia and aflatoxins.

Lipid metabolism especially that of fatty acid-derived oxylipins, has been shown to play a role in *A. flavus* virulence, development and aflatoxin production (Amaike and Keller, 2009; Burow *et al.*, 1997). Recently, Luo *et al.* (2016) identified *far* transcription factor genes that regulate fatty acid beta-oxidation required for aflatoxin biosynthesis. Aflatoxin production was significantly decreased in a  $\Delta farB$  mutant. A putative RNA-Pol II transcription elongation factor gene, *rtfA*, was shown to regulate development and SM production in *A. flavus* possibly through an epigenetic mechanism (Lohmar *et al.*, 2016). A  $\Delta rtfA$  mutant exhibited a significant decrease in sclerotial and aflatoxin production as well as other uncharacterised SMs. Another epigenetic factor controlling development and aflatoxin production in *A. flavus* is the histone-modifying methyltransferase, RmtA (Satterlee *et al.*, 2016). RmtA was shown to be a positive regulator of sclerotial production while suppressing conidiation and aflatoxin production. Upon forced overexpression, a globally-acting C<sub>2</sub>H<sub>2</sub> zinc-finger transcription factor, MtfA, was shown to be required

for normal development and aflatoxin production in *A. flavus* (Zhuang *et al.*, 2016).

## 6. Other *Aspergillus flavus* secondary metabolite gene clusters

Of the 56 predicted SM gene clusters in the *A. flavus* genome, only twelve clusters have been associated with a specific metabolite (Figure 4). A number of metabolites from uncharacterised gene clusters have been identified utilising comparative metabolomics of cluster gene knockout and overexpressing mutants to wild-type strains (Cary *et al.*, 2014; Forseth *et al.*, 2013; Saruwatari *et al.*, 2014). However, elucidation of the products of some SM gene clusters is confounded by the fact that many are transcriptionally silent, inactive due to gene mutations, or expressed at extremely low levels thus making identification of their respective products problematic under laboratory growth conditions. These types of gene clusters are termed cryptic or orphan and identification of their metabolite(s) cannot be achieved using classical deletion or overexpression of a cluster gene followed by comparative metabolite analysis by LC-MS. Analysis of RNA-Seq data from a number of *A. flavus* strains growing on artificial media indicates that approximately half of the SM gene clusters are silent or expressed at very low levels (Ehrlich and Mack, 2014). A number of methods have been used to overcome silent gene clusters thereby enabling identification of some previously unknown cluster metabolites (Brakhage, 2013; Brakhage and Schroeckh, 2011; Strauss and Reyes-Dominguez, 2011). In addition to the AF gene cluster, the metabolites produced by the following clusters have been experimentally identified in *A. flavus*.

### Cyclopiazonic acid cluster

CPA is an indole-tetramic acid produced by the activity of a hybrid PKS-NRPS in *A. flavus*. The toxicology and biosynthesis of CPA has been reviewed by Chang *et al.* (2009a). In addition to being an inhibitor of calcium-dependent ATPases present in the sarcoplasmic reticulum of animals (Riley *et al.*, 1992), CPA has also been suggested to function as a pathogenicity factor that can induce cell death in plants (Chalivendra *et al.*, 2017). The CPA cluster is located in the subtelomeric region adjacent to the AF gene cluster and can be absent in some *A. flavus* strains (Chang *et al.*, 2009b). Five genes are thought to comprise the *A. flavus* CPA cluster; a monoamine oxidase, *maoA*; dimethylallyl tryptophan synthase, *dmtA*; hybrid PKS-NRPS, *pks/nrps*; MFS transporter, *mfs1*; and a putative C6-type transcription factor, *ctfR1*. Disruption of any of these genes except the *ctrR1* and *mfs1* resulted in loss of CPA production. A CPA-producing *A. oryzae* strain was shown to have two additional genes, *cpaH* and *cpaM*, located between *maoA* (*cpaO*) and *mfs1* (*cpaT*) (Kato *et al.*, 2011). While no function was attributed to *cpaM*, *cpaH* encodes a






Figure 4. Representative structures of secondary metabolites that have been experimentally linked to a gene cluster in *Aspergillus flavus*. Cluster designations are based on the numbering system of Georgianna *et al.* (2010).

<sup>a</sup> The kojic acid cluster was characterised in *Aspergillus oryzae* and was not identified by SMURF analysis of *A. flavus* so a cluster number was not designated.

<sup>b</sup> Genes required for aflatrem biosynthesis are also present on Cluster 15.

<sup>c</sup> In addition to the Ina gene cluster (#35) a highly homologous Inb piperazine gene cluster (#48) was also present in *A. flavus*.

cytochrome P450 that mediates the conversion of CPA to 2-oxoCPA, the less toxic end product of the CPA pathway in *A. oryzae*. The absence of homologs of *cpaH* and *cpaM* in *A. flavus* 3357 suggests that they were lost during the evolution of *A. flavus* and may reflect the need for the more toxic form of CPA in the field.

### Aflatrem clusters

Aflatrem and its isomer,  $\beta$ -aflatrem, are examples of indole-diterpenes (IDTs), a class of structurally-diverse secondary metabolites (Saikia *et al.*, 2008). Aflatrem was shown to be a potent mammalian tremorgen (Gallagher and Wilson, 1979). Identification of the genes required for aflatrem biosynthesis (*atm*) in *A. flavus* was made possible through comparative genomics with *Penicillium paxilli* genes (*pax*) required for the production of the IDT, paxilline (Saikia *et al.*, 2006; Zhang *et al.*, 2004). Unlike most SM gene clusters, the genes required for the synthesis of aflatrem are located on two different chromosomes (5 and 7) in *A. flavus* (Nicholson *et al.*, 2009). In addition to aflatrem, *A. flavus* also produces a similar class of IDTs termed aflavinines that demonstrate anti-insectan properties. Both aflatrem and aflavinines are derived from a common core structure consisting of a tetracyclic diterpene originating from the activities of a geranylgeranyl diphosphate synthase (*AtmG*) and geranylgeranyl transferase (*AtmC*). The diterpene is subsequently converted to either aflatrem or aflavinine by the action of unique IDT cyclases (IDTCs) (Tang *et al.*, 2015). Interestingly, Tang *et al.* (2015) identified a standalone IDTC (not associated with the aflatrem/aflavinine gene cluster) that was responsible for production of aflavinine but not aflatrem. It is unclear how *A. flavus* coordinates the expression of the IDTCs responsible for production of aflatrem or aflavinines (mainly present in sclerotia) but it may depend on the type of biotic pressure present in the particular ecological niche inhabited by the fungus.

### Piperazine clusters

Similar to aflatrem, two separate gene clusters were identified termed *lna* (chromosome 6) and *lnb* (chromosome 8) that are involved in the biosynthesis of piperazines and associated derivatives in *A. flavus* (Forseth *et al.*, 2013). Both clusters contain genes encoding putative NRPS-like proteins (*lnaA/lnbA*), as well as decorating enzymes, such as an NMR-like protein (*lnaB/lnbB*), cytochrome P450s (*lnaC/lnbC*), alcohol dehydrogenase/NADH:flavin oxidoreductase (*lnaE/lnbE*), and a transporter (*lnaF/lnbF*). The *lna* cluster also contains a second putative P450 enzyme gene (*lnaD*) while the *lnb* cluster contains a gene encoding a hypothetical protein not present in the *lna* cluster.

Comparative metabolomics of *A. flavus* wild-type, deletion, and overexpression (OE) strains allowed identification

of two major piperazine metabolites and eight other *lna* cluster-derived metabolites that had not been previously reported in aspergilli or other fungi (Forseth *et al.*, 2013). Both *lnaA* and *lnbA* proteins activate tyrosine, but because they lacked condensation domains, there was no peptide bond formation. It was postulated that *lnaA* and *lnbA* C-terminal reductase domains reduce L-Tyr to its aldehyde suggesting that these enzymes more closely resemble amino acid reductases characteristic of Lys2-type reductases involved in fungal L-lysine biosynthesis (Ehmann *et al.*, 1999). The resulting L-Tyr aldehyde subsequently dimerizes and is further reduced to piperazine and morpholine-like compounds. Of particular interest was the finding that *lnaB* can participate in the reduction of *lnaA*-derived intermediates indicating that the *lna* and *lnb* clusters encode partially redundant biosynthetic pathways capable of sharing biosynthetic intermediates. Down-regulation of both of *lnaA* and *lnbA* expression in *A. flavus* resulted in significant reduction in sclerotial formation compared to the wild type suggesting a role for the secondary metabolite production of these clusters in fungal development (Forseth *et al.*, 2013).

### Ditryptophenaline cluster

The ditryptophenaline (DTP) cluster is responsible for the production of the dimeric diketopiperazine alkaloid, (-)-ditryptophenaline, that is derived from two molecules of tryptophan and phenylalanine (Saruwatari *et al.*, 2014). Comparative metabolomics of putative *A. flavus* DTP cluster gene (*dtp*) mutants with a control strain enabled determination of the function of the cluster genes. The cluster was found to be composed of three genes encoding a NRPS (*dtpA*) that condenses L-tryptophan and L-phenylalanine together to form the diketopiperazine core; an N-methyltransferase (*dtpB*) that methylates the core structure; and a cytochrome P450 (*dtpC*) that is likely responsible for cyclization and dimerization of the methylated diketopiperazine to form DTP. Though DTP has been shown to have weak activity as an inhibitor of the neurotransmitter, substance P, in animal models, its function in the biology of *A. flavus* is not known (Barrow and Sedlock, 1994).

### Ustiloxin B cluster

Ustiloxins are a diverse class of tetrapeptides with ustiloxin B consisting of the amino acids Tyr-Ala-Ile-Gly (YAIG) (Koiso *et al.*, 1994). Ustiloxin B is derived from enzymes encoded by a cluster of 18 genes needed to cyclise the tetrapeptide at the side chains of Tyr and Ile and catalyse side chain modifications and link a non-proteinogenic amino acid, norvaline, to the Tyr moiety (Umemura *et al.*, 2014). Ustiloxin B was initially thought to be the product of a NRPS-like enzyme, but no NRPS-like genes were present in or near the gene cluster. The presence of a gene (*ustA*)

in the cluster encoding a precursor protein harbouring 16 short YAIG repeat peptide regions and an N-terminal signal peptide suggested that ustiloxin B is biosynthesised through a ribosomal peptide synthetic pathway (RiPS) not previously described in *A. flavus* or other ascomycetes (Umemura *et al.*, 2014).

Screening of *Aspergillus* genome sequences by Nagano *et al.* (2016) for the presence of the *ustYa/ustYb* and *ustA* gene homologs indicated that there are as many as 94 ribosomal peptide precursor candidate genes displaying a variety of core peptide sequences in addition to YAIG. From this computational analysis a novel bicyclic peptide designated asperipin-2a was experimentally identified in *A. flavus*. Ustiloxin B was shown to inhibit polymerisation of purified porcine microtubule proteins as well as mitosis in human cell lines, however the role of ustiloxin B in the biology of *A. flavus* remains to be determined (Koiso *et al.*, 1994).

### Kojic acid cluster

Though not originally confirmed in *A. flavus*, the KA gene cluster was experimentally identified in its close relative, *A. oryzae* (Terabayashi *et al.*, 2010). Comparative transcriptomics of *A. oryzae* cultures grown under KA-conducive and non-conducive conditions, allowed identification of two genes responsible for KA biosynthesis. It was determined from analysis of the *A. oryzae* genome sequence that these two genes mapped within 3 kb of one another. One of the genes designated *kojA*, encodes a predicted FAD-dependent oxidoreductase while the other, *kojT* encodes a major facilitator superfamily transporter. *KojA* is believed to oxidise glucose to KA, although other as of yet unidentified gene products may be required for complete conversion of glucose to KA (Terabayashi *et al.*, 2010). Located between *kojA* and *kojT* is a gene designated *kojR*, encoding a Zn(II)-Cys6 transcription factor that when deleted caused loss of KA production while its overexpression led to significantly higher levels of KA (Marui *et al.*, 2011).

Kojic acid may function as an insect anti-feedant and it has also been suggested that it functions as an antioxidant that inhibits the ability of insect oxidative enzymes to detoxify AFB<sub>1</sub> (Dowd, 1988). While KA was shown to be relatively non-toxic in mouse assays, it appeared to act synergistically with AFB<sub>1</sub> in insect feeding studies causing increased mortality compared to feeding of either metabolite alone (Dowd, 1988).

### Asparasone cluster

Asparasone A, a PKS-derived anthraquinone, is responsible for the formation of the dark pigment associated with sclerotia (Cary *et al.*, 2014). Asparasone A was first described in *A. parasiticus* and demonstrated to possess anti-insectan

properties (Sobolev *et al.*, 1997). The asparasone A gene cluster consists of four genes, a polyketide synthase (*pks27*), a Zn(II)-Cys6-type transcription factor (*znf27*) and putative high-affinity glucose (*mfs1*) and MFS transporters (*mfs2*). Deletion mutants of *A. flavus pks27* produced greyish-yellow sclerotia rather than the wild-type dark brown pigment. Gene expression analysis showed a significant reduction in expression of *pks27*, *mfs1* and *mfs2* in the  $\Delta znf27$  mutant compared to the control indicating that *znf27* functions as a pathway-specific transcription factor. Expression of *pks27* was shown to be dependent on VeA.

Cary *et al.* (2014) revealed that the dark brown pigments present in *A. flavus* sclerotia are the product of Pks27-derived anthraquinones rather than typical tetrahydronaphthalene-based melanins. The asparasone A-based anthraquinone pigment in sclerotia was shown to provide resistance to both biotic and abiotic stressors.

### Leporin cluster

Leporins were originally identified in *Aspergillus leporis* as 2-pyridones that exhibited anti-insectan properties (TePaske *et al.*, 1991). Cary *et al.* (2015b) described a silent gene cluster in *A. flavus* that consisted of a hybrid PKS-NRPS (*lepA*) and six additional genes responsible for production of leporin B and the novel metabolite, leporin C. Overexpression of a putative cluster-specific transcriptional regulatory gene, *lepE*, resulted in transformants producing a distinctive orange-red pigment identified as a highly stable leporin B trimer complexed with an atom of iron (trioxyleporin B). Such a trimeric iron chelate had not been previously described for a 2-pyridone nor had there been any reports of siderophore-like molecules produced by a hybrid PKS-NRPS in fungi.

While leporin A was shown to have anti-insectan properties there were no previous reports on the insecticidal properties of leporins B or C. Insect feeding studies performed with a preparation of the leporin B-iron chelate (LIC) compound showed a significant reduction in growth of first instar larvae of Fall armyworms (*Spodoptera frugiperda*) and Corn earworms (*Helicoverpa zea*) indicating that the LIC was acting as an antifeedant, but not an anti-insectan agent (Cary *et al.*, 2015b). A 10-fold reduction in conidia production was also observed in the *lepE* overexpressing strain (OE::*lepE*) and sclerotial production was either reduced or delayed compared to the control.

### Aflavarin cluster

The anti-insectan secondary metabolite, aflavarin, was first isolated and characterised from sclerotia of *A. flavus* (TePaske *et al.*, 1992). Comparative transcriptomics of an *A. flavus*  $\Delta veA$  mutant and isogenic control strain led to the identification of the aflavarin gene cluster (Cary *et al.*,



2015a). The aflavarin gene cluster is composed of genes encoding a PKS (*afvB*), O-methyl transferase (*afvC*), methyl transferase (*afvD*), cytochrome P450 (*afvE*) and a NADH oxidase (*afvA*). Comparative analysis of the metabolomes of an *A. flavus* control and  $\Delta afvB$  (PKS) mutant identified a series of bicoumarins that were present in the control and absent in the mutant (Cary *et al.*, 2015a). The major metabolite identified had a calculated mass of 455, which was identical to the mass of aflavarin.

In addition to its anti-insectan properties, aflavarin was required for normal levels of sclerotial production in *A. flavus* (Cary *et al.*, 2015a). While no cluster-specific transcriptional activator gene was present in the aflavarin cluster, a number of putative binding sites for stress-responsive DNA binding proteins as well the aflatoxin cluster-specific transcriptional activator, AflR were identified. This suggested that there may be regulatory cross-talk between these two clusters (Cary *et al.*, 2015a).

## 7. Environmental impacts on *Aspergillus flavus* and aflatoxin production

Environmental conditions in agriculturally relevant areas will likely undergo dramatic changes that are expected to affect precipitation, temperature, and the composition of atmospheric gasses within the next 100 years. Data from the authoritative UN Panel on Climate Change indicate that global temperatures will increase approximately 0.2-0.4 °C per decade at increasing rates (IPCC, 2014) depending on mitigation efforts. Forecasts further indicate increases in carbon dioxide (CO<sub>2</sub>), methane, nitrous oxide, aerosol, and other gases, with CO<sub>2</sub> reaching as high as 1000 ppm, nearly tripling current levels. Periods of heavier rainfall and periods of longer drought are also forecast (IPCC, 2013, 2014). Not surprisingly, many of these factors are known to impact fungal growth, secondary metabolite production, and host-pathogen interaction (Giorni *et al.*, 2008a,b; Marín *et al.*, 1998; Medina *et al.*, 2015; Sultan, 2010; Wu *et al.*, 2011). Gaining insight into how the combination of climate change-related weather variables will influence *A. flavus* pathogenicity is vital for the continued development of strategies aimed at preserving global food security.

Both laboratory and farming experiences have provided examples of drought conditions and increased temperature causing increased aflatoxin production (Diener *et al.*, 1987; Giorni *et al.*, 2007; Kos *et al.*, 2013; Payne *et al.*, 1988; Piva *et al.*, 2006). A particularly illustrative example occurred in 2012 in Serbia when daily temperatures (30-40 °C) and the number of hot days increased significantly from previous years and precipitation decreased to drought levels. That year, 68% of maize samples tested were positive for aflatoxin contamination whereas no contamination was reported during the previous years (Kos *et al.*, 2013). A similar incident in 2004-2005 was also documented in

northern Italy (Giorni *et al.*, 2007). While these examples provide adequate evidence for concern, it is essential for researchers to focus on exacerbating circumstances and identify means of mitigation.

The influence of temperature and water on fungal growth and pathogenicity has been studied for decades (Anderson and Smith, 1972; Holmquist *et al.*, 1983; Northolt *et al.*, 1977; Sabburg *et al.*, 2015; Schmidt-Heydt *et al.*, 2010). Data from our laboratory and that of others (Marín *et al.*, 1998; O'Brian *et al.*, 2007; Yu *et al.*, 2011) have demonstrated optimal conditions for the expression of aflatoxin cluster genes and aflatoxin production between 28 °C and 30 °C. Aflatoxin biosynthesis increases as temperatures increase between 20 °C and 30 °C, where it can remain in production until temperatures reach 37 °C (Abdel-Hadi *et al.*, 2012; Schmidt-Heydt *et al.*, 2010). This is significant in the context of crop contamination and especially that of maize, which typically grows when air temperatures are between 10 °C and 30 °C (Cross and Zuber, 1972; Greaves, 1996). Greenhouse and field studies have characterised this interaction in the context of pathogenesis. In one study maize grown at 32 to 38 °C had 73% of kernels infected, whereas at 21-26 °C, only 2.5% of kernels were infected (Jones *et al.*, 1980). Drought stress has also been shown to induce higher levels of aflatoxin contamination (Davis *et al.*, 1986), and the irrigation of maize fields during periods of low precipitation has been shown to reduce fungal infection and aflatoxin contamination (Jones *et al.*, 1981).

Recently, functional genomics tools, i.e. microarray technologies, quantitative proteomics, and RNA-sequencing, have examined the effect of abiotic factors on fungal development and plant-fungus interactions (Abdel-Hadi *et al.*, 2012; Georgianna *et al.*, 2008; Lanubile *et al.*, 2010; Luo *et al.*, 2010; Zhang *et al.*, 2014). Microarray analysis demonstrated a complex regulation of genes within the aflatoxin cluster in response to temperature and water. In general, the aflatoxin cluster genes exhibit lower expression at 37 compared to 28 °C (O'Brian *et al.*, 2007). It was further determined by Schmidt-Heydt *et al.* (2009) that different temperature and water combinations induce increased expression of the regulatory gene *aflS*, which interacts with *aflR*, and is correlated with high aflatoxin production. Further, a high *aflS:aflR* gene expression ratio (above 1) was also highly correlated with aflatoxin production. Subsequent microarray analysis and predictive modelling done by Abdel-Hadi *et al.* (2012) extended this by establishing that *aflD* and *aflM* are also highly subjected to temperature and water availability, and the relative expression of *aflD* or *aflM* to that of *aflR* and *aflS*, are related to temperature, water activity levels and aflatoxin production. An RNA-seq study of *A. flavus* interacting with maize kernels at various temperature and water activity levels corroborated previous observations, and identified

an interactive effect of temperature and water activity on *aflR* and *aflS* expression (Medina *et al.*, 2017b).

Research has recently begun addressing the impact of increased carbon dioxide levels on *A. flavus*, as research in other host-pathogen systems has indicated that CO<sub>2</sub> levels can exacerbate host infection (Vary *et al.*, 2015). In the first attempt we know of to characterise the interaction of three environmental variables (temperature, water and CO<sub>2</sub>) on *A. flavus* growth and aflatoxin production, Medina *et al.* (2015) demonstrated that growth of *A. flavus* was unaffected by CO<sub>2</sub> levels at 1000 ppm. However, there was markedly increased expression of aflatoxin cluster genes and AFB<sub>1</sub> levels in response to high CO<sub>2</sub> levels, especially at high temperature (37 °C) and low water activity levels (0.92-0.95<sub>aw</sub>). Similar but modest effects were observed at the lower temperature tested (34 °C). Ongoing work in our laboratory and from others will address this important research area of interest.

The possible combined effects of drought stress, temperature, and carbon dioxide levels on the fungus-plant interaction and aflatoxin contamination level have not yet been extensively studied. It has been suggested that elevated CO<sub>2</sub> may increase crop metabolism resulting in increased crop yields (Driever and Kromdijk, 2013). However, several researchers have further speculated that higher temperatures and higher CO<sub>2</sub> levels will still likely result in conditions further accommodating fungal growth (DaMatta *et al.*, 2010; Paterson and Lima, 2011; Porter and Semenov, 2005). It has also been shown that drought-stressed maize exhibits suppression of several genes related to pathogen resistance such as pathogenesis-related proteins, heat shock proteins, and cell wall degrading enzymes (Luo *et al.*, 2010; Yu and Setter, 2003). Thus, there remains a pressing need to examine the effects of CO<sub>2</sub> on plant processes during fungal infection.

The effects of temperature and water activity on small non-coding RNA's (Bai *et al.*, 2015), development and the production of other secondary metabolites have also been examined (Medina *et al.*, 2017b; Yu *et al.*, 2011; Zhang *et al.*, 2014). Yu *et al.* (2011) demonstrated that 10 of the 55 gene clusters contained genes with increased expression at 30 °C as compared to 37 °C (cluster 1, 10, 11, 19, 20, 21, 24, 45, 54, and 55) and Cluster 3 decreased at the lower temperature examined. Medina *et al.* (2017b) also examined the secondary metabolic gene cluster response, where changes in water activity or temperature affected genes in the CPA, ustiloxin B, and penicillin clusters. Because secondary metabolites are intricately involved in *A. flavus* growth, development, and have toxigenic potential, characterising their expression levels in addition to aflatoxin is also of importance.

## 8. Genetic improvement of crops to control aflatoxin contamination

Pre-harvest aflatoxin contamination is a very complex problem affected by a multitude of biotic and abiotic factors (Bhatnagar-Mathur *et al.*, 2015; CAST, 2003; Gressel and Polturak, 2017). Therefore, a multi-pronged approach may need to be employed to control aflatoxin contamination when conditions in the field are favourable for fungal infection. Prominent among these approaches is the development of preharvest host plant-resistance through classical or molecular breeding (Bhatnagar *et al.*, 2008a,b). Less contamination during preharvest stage is very beneficial for reduced postharvest contamination. With the knowledge available so far on aflatoxin biosynthesis and natural and synthetic antifungal genes, it is now possible to impart innate host plant resistance using a combination of approaches.

### Classical breeding

Although classical breeding is a proven approach to develop resistance, it is slow and is hampered by the lack of available resistance genes in germplasm in some susceptible crops. Several maize lines have been identified and developed with increased resistance to *A. flavus* infection and aflatoxin contamination and this has enabled the identification of natural resistance traits (Brown *et al.*, 2004, 2006, 2010; Campbell and White, 1995). However, these investigations have indicated that resistance to aflatoxin contamination is polygenic. Therefore, attempts to move resistance from inbred lines into commercial varieties with desirable agronomic characteristics have been a slow process due to the lack of availability of biomarkers to facilitate the transfer of resistance genes (Menkir *et al.*, 2008; Williams *et al.*, 2015). However, the roles of some antifungal proteins in maize have been validated (Chen *et al.*, 2010; Majumdar *et al.*, 2017b) and mapped to the genome (Chen *et al.*, 2016). More research is needed to elucidate the biochemical mechanisms that manifest the resistance phenotype in maize kernels or other sources so that they can be utilised to enhance resistance through marker-assisted breeding in maize or genetic engineering in cotton (Brown *et al.*, 2009, 2010; Cary *et al.*, 2009; Rajasekaran *et al.*, 2006, 2009a,b; Williams *et al.*, 2015).

Recent breeding efforts towards the development of aflatoxin-resistant maize lines have resulted in a number of germplasm release as IITA-SRRC TZAR inbreds. In 2008, TZAR 101-106, derived from a combination of African and southern-adapted US lines have been field-tested in different parts of the Southern USA with promising results (Menkir *et al.*, 2008). Recently, these lines were also determined to be resistant to both aflatoxin and fumonisin contamination (A. Menkir, unpublished data). They also exhibited resistance to lodging and common foliar diseases.

GT-603 was released in 2011, after having been derived from GT-MAS:gk (Guo *et al.*, 2011), while Mp-718 and Mp-719 were released as southern adapted resistant lines which are both shorter and mature earlier than previous Mp lines (Williams and Windham, 2012). These lines are also being tested as inbreds and in hybrid combinations in the Southern USA.

Unlike maize, cotton or peanut have a limited diversity in their germplasm and no defined mechanism or well-characterised resistance proteins/genes have been identified to date for natural resistance to *A. flavus* (Holbrook *et al.*, 2000; Moore *et al.*, 2017). For this reason, it is critical that a seed-based resistance be developed using either genetic engineering or molecular breeding approaches in these crops.

### Molecular breeding

Molecular breeding through biotechnological means is very attractive because it is less time consuming. Only a limited success has been demonstrated in the development of antifungal transgenic crops (Parisi *et al.*, 2016). Development of aflatoxin-resistant crops is further exacerbated by the fact that the causal fungus is an opportunistic saprophyte and does not follow typical host-pathogen interaction. However, success in this area will depend in large part on the following factors:

- Identification of resistance proteins, native or foreign, that express inhibitory activity against the mycotoxigenic fungi. A number of potential maize resistance-associated proteins (RAPs) and the genes encoding them have been identified (Table 2) and some of these have been introduced into other crops for evaluation (Chen *et al.*, 2006, 2007, 2010; Rajasekaran *et al.*, 2005a, 2008). Proteomic analyses of resistant maize kernels revealed existence of many resistance associated proteins and they have been shown to play an active role in resistance to *A. flavus* contamination (Chen *et al.*, 2002, 2015; Majumdar *et al.*, 2017b). Unlike maize, there are no well-characterised resistance proteins/genes available in cotton germplasm that demonstrate enhanced resistance to *A. flavus* infection and aflatoxin contamination. A sample of maize antifungal proteins that are inhibitory to *A. flavus* are given in Table 3.
- Development of efficient transformation methods that can be readily adapted to agronomically superior crop varieties (Cary *et al.*, 2011; Holbrook *et al.*, 2000; Rajasekaran *et al.*, 2005b).
- Selection of gene promoters that will provide the desired spatial and temporal regulation of antifungal transgenes and result in efficacious levels of production of the antifungal protein/peptide in targeted tissue or organelle (DeGray *et al.*, 2001; Rajasekaran *et al.*, 2012).

**Table 2. Resistance-associated proteins identified through comparative proteomics (adapted from Brown *et al.*, 2010).**

<i>Antifungals</i>
Zeamatin
Trypsin inhibitor 14kDa (TI)
Ribosome inactivating protein (RIP)
$\beta$ -1,3-glucanase
Pathogenesis-related protein 10 (PR10)
PR10.1
Trypsin inhibitor 10 kDa (ZmTI)
<i>Stress-related</i>
Aldose reductase (ALD)
Cold-regulated protein (ZmCORp)
Water stress inducible protein (WSI)
Anionic peroxidase
Small heat shock protein 16.9/17.2 kDa
Glyoxalase I (GLX I)
Peroxiredoxin 1 (PER1)
<i>Storage</i>
Globulin I
Globulin II
Cupin domain containing protein (Zmcup)
Late embryogenesis abundant protein (LEA III)
LEA 14
<i>Other</i>
Serine/threonine protein kinase
Translation initiation factor 5A

Molecular breeding has become a useful rapid technology for development of crops resistant to aflatoxin contamination. Prominent among them is heterologous expression of foreign antifungal gene(s), either natural or synthetic, to provide resistance to the fungus and toxin production (Rajasekaran *et al.*, 2005b; Ruhlman *et al.*, 2014; Schubert *et al.*, 2015; Sharma *et al.*, in press). RNA-interference (RNAi)-mediated host induced gene silencing (HIGS) of key fungal genes critical for fungal pathogenesis and aflatoxin production has been successfully employed to incorporate *A. flavus* resistance in susceptible maize or peanut varieties (Arias *et al.*, 2015; Bhatnagar-Mathur *et al.*, 2015; Majumdar *et al.*, 2017a; Masanga *et al.*, 2015; Power *et al.*, 2017; Sharma *et al.*, in press; Thakare *et al.*, 2017). HIGS technology does not require that the host plant express a foreign protein so food and feed produced from resistant lines of transgenic maize should be more acceptable to regulatory agencies and consumers. Another set of novel methods that is available for crop improvement includes several gene-editing methods such as site-specific nucleases (SSNs), such as meganucleases (MNs), also known as homing nucleases (HNs), zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and RNA-guided nuclease (RGN) systems. The most widely used RGN is the clustered regularly interspaced short palindromic repeat (CRISPR)/



**Table 3. Natural and synthetic proteins/peptides with antifungal activity against *Aspergillus flavus*.**

Protein/peptide	Protein family	Source	Mode of action	Reference
Haloperoxidase	peroxidase	<i>Pseudomonas pyrocinia</i>	produce antimicrobial compounds – peracetic acid and hypohalites	Rajasekaran <i>et al.</i> (2000); Jacks <i>et al.</i> (2000)
$\beta$ 1-3 glucanase	glycosyl hydrolase	tobacco	hydrolysis of fungal cell wall components	Sundaresha <i>et al.</i> (2010)
Ib-AMP <sub>3</sub>	defensin	sweet potato	lytic	De Lucca <i>et al.</i> (1999)
AILP	lectin	hyacinth bean	inhibits germination and hyphal growth	Fakhoury and Woloshuk (2001)
Chitinase	glycosyl hydrolase	maize inbred Tex6	hydrolysis of fungal cell wall components	Moore <i>et al.</i> (2004)
ZmCORp	lectin	maize kernels	hemagglutination activity against fungal conidia	Baker <i>et al.</i> (2009)
Mod-1/RIP-1	ribosome-inhibiting protein	maize	inhibits hyphal tip growth	Nielsen <i>et al.</i> (2001); Weissinger <i>et al.</i> (2007)
Zeamatin	PR-5	maize	inhibits hyphal tip growth	Guo <i>et al.</i> (1997)
ZmPR10	PR-10	maize	RNAse activity	Chen <i>et al.</i> (2006)
ZmPRms	PR	maize kernels	defence regulator to modulate resistance	Majumdar <i>et al.</i> (2017b)
Thanatin	antimicrobial	Spined soldier bug	inhibits cellular respiration	Schubert <i>et al.</i> (2015)
Trypsin inhibitor	protease inhibitor	maize	trypsin/amylase inhibition	Chen <i>et al.</i> (1998)
Purothionin/hordothionin	thionin	barley/wheat	lytic	K. Rajasekaran, unpublished data
D4E1	cecropin	synthetic peptide	lytic	Cary <i>et al.</i> (2000); Rajasekaran <i>et al.</i> (2005b)
D5C/D5C1		synthetic peptide	lytic	Weissinger <i>et al.</i> (2000)
AGM	tachyplesin	synthetic peptide	lytic	Rajasekaran <i>et al.</i> (2018)
D2A21		synthetic peptide	lytic	Weissinger <i>et al.</i> (2000)
MSI99	magainin	synthetic peptide	lytic	DeGray <i>et al.</i> (2001)

CRISPR-associated system 9 (CRISPR/Cas9), and DNA-guided nuclease (DGN) system, i.e. NgAgo (an acronym for *Natronobacterium gregoryi* Argonaute) (Miglani, 2017). CRISPR/Cas9/sgRNA system has been applied successfully to improve cultivated crops with agronomic traits. It is an inexpensive, easy, most user friendly, and rapidly adopted genome editing tool that enables precise genomic modifications in many different organisms including higher plants (Khatodia *et al.*, 2016; Wolt *et al.*, 2016; Zhang *et al.*, 2017). Development of genetically edited crops is similar to those developed by conventional or mutation breeding and this potential technique is useful in providing sustainable productive agriculture for better feeding of rapidly growing population in a changing climate. Once the target host plant genes are identified through comparative transcriptomic and gene network analyses in susceptible crops (Bedre *et al.*, 2015; Clevenger *et al.*, 2016; Lee *et al.*, 2012; Musungu *et al.*, 2015) these modern tools can be applied for generating aflatoxin resistant crops.

## 9. Post-harvest aflatoxin control strategies

Control of preharvest aflatoxin contamination has been addressed in the previous sections; however, it is important to consider aspects of postharvest control as well. Harvested produce with minimal or no fungal growth or aflatoxin levels is a good starting point to avoid postharvest aflatoxin

contamination problems. Raw commodities prone to aflatoxin contamination (maize, groundnuts, tree nuts, spices, dried figs) must be effectively dried to facilitate short and medium term storage. Often in tropical countries aflatoxin is formed pre-harvest because of the conducive environmental conditions during ripening. However, to minimise subsequent toxin contamination effective drying and storage regimes are required for food and feed use. For example, maize is often harvested at 17-20% moisture content (m.c.) and the cereal must be dried effectively to around 15% m.c. (= 0.70  $a_w$ ) to avoid continued and increased contamination with aflatoxins. Some of these commodities, e.g. ground nuts and some trees nuts are very hygroscopic and can absorb moisture during storage from the atmosphere. They need to be stored at <7.5% m.c. (= 0.70  $a_w$ ). Otherwise, it can result in water and temperature conditions conducive to growth of xerophilic fungi such as *A. flavus* and aflatoxin contamination. Effective solar drying or in more developed systems heated drying is necessary for reaching the target safe m.c. levels. Table 4 shows the relationship between safe m.c. for these commodities and the equivalent water availability level for these commodities for safe short and medium term storage and those which represent marginal conditions and represent a risk of aflatoxin contamination.

**Table 4. The relationship between storage moisture content (m.c.) for safe storages and those representing a risk from aflatoxin contamination.**

Commodity	Safe m.c. (% wet weight basis) = water activity ( $a_w$ )	At risk m.c. conditions
Maize	15.0% (= 0.70 $a_w$ )	>16.5-17.0%
Peanut	9% (unshelled; = 0.70 $a_w$ ) 7.5% (shelled; = 0.65 $a_w$ )	>10%
Pistachio	6.5-7.0% (unsalted; = 0.65-0.70 $a_w$ )	>8.5-9.0%
Chillies	8-12% (= 0.60-0.70 $a_w$ )	>12%

Molecular approaches have been used to monitor and control aflatoxin contamination post-harvest, especially in stored peanuts. For example, Abdel-Hadi *et al.* (2010) showed that the structural *aflD* gene in the biosynthetic cluster could be effectively used to monitor changes in *A. flavus* populations in peanuts stored under different storage conditions. Passone *et al.* (2010) found a correlation between populations of *Aspergillus* section *Flavi* (colony forming units) and RT-PCR for the *aflD* gene ( $r=0.613$ ;  $P<0.0001$ ). Changes in key structural and regulatory genes (e.g. *aflR*; *aflS*) can also be utilised effectively to monitor relative aflatoxin risk post-harvest.

There have also been studies which suggest that real time monitoring of CO<sub>2</sub> production, temperature and relative humidity could be used as good indicators of whether moulding and mycotoxin contamination may be initiated. Indeed, it has been shown that this approach can give an early warning system of relative risks of mycotoxin contamination. For example, for fumonisin contamination of maize it was shown that <0.8-0.9% dry matter loss would result in maize being contaminated with toxin at above the EU legislative limits (Mylona *et al.*, 2014). Recently, it has also been shown that in brown rice the risk of aflatoxin contamination post-harvest may be much higher than in unhusked paddy rice (Martin Castaño *et al.*, 2017).

A recent study by Waliyar *et al.* (2015) showed that effective drying of peanut pods or heated dryers and shellers can reduce post-harvest losses and reduce aflatoxin contamination of peanut. Storage at <7.5% m.c. at 10 °C can result in safe storage for between 6-12 months. However, at higher temperatures this time frame may be shortened as this may allow some reabsorption of moisture from the atmosphere and allow initiation of spoilage mould activity and an increase in aflatoxin contamination.

An integrated approach in Argentina encompassing pest control with entomogenous fungi + antioxidants in stored peanut and maize has shown effective control of aflatoxin contamination (Barra *et al.*, 2013, 2015a,b). This

approach has optimised the use of entomogenous strains of *Purpureocillium lilacinum* for control of storage pests in silos, as well the use of antioxidants for controlling *A. flavus* colonisation. This has achieved significant reductions in AFB<sub>1</sub> contamination. This post-harvest integrated pest management strategy may be an effective way forward for controlling both pests and *A. flavus*, especially where hygiene practices are not effective. Again the cost-effectiveness of this approach and formulation of this type of application needs to be evaluated.

Gaseous treatment of stored commodities with either sulphur dioxide (SO<sub>2</sub>) or ozone (O<sub>3</sub>) has been examined for control of aflatoxins and other mycotoxins (Magan, 1993; Mylona *et al.*, 2014). Indeed, O<sub>3</sub> has been suggested as a method for control of aflatoxins in Brazil nuts (Giordana *et al.*, 2012). However, the economics of this approach combined with the associated deleterious health and safety issues limited its use. In addition, effects on organoleptic properties may also need to be considered as thresholds to prevent quality deterioration may be very low. The recent report by WHO (2017) also suggests that where pre- and post-harvest treatments are considered the potential for co-occurrence of different mycotoxins needs to be considered. Certain post-harvest control strategies may result in a switch to another mycotoxin, which could subsequently pose problems (e.g. aflatoxins to CPA).

## 10. Conclusions

Based on the severity and the extent of aflatoxin contamination of various commodities prior to harvest, and many more in storage worldwide, the health implications of this contamination as well as the very significant economic impact to many economies (discussed in WHO, 2017), *A. flavus* has probably become one of the most thoroughly studied microorganism, both at the macro level as well as at the molecular level. The purpose of all these studies is to find meaningful and lasting strategies to control this contamination. Biological control has been demonstrated to be a strategy that is consistently functional in various parts of the world to significantly reduce pre-harvest aflatoxin contamination, which may be carried over to storage as well. However, biocontrol has its limitations and will not completely eliminate toxin contamination. Therefore, this strategy must be complemented with enhancement of host-resistance either by transgenic expression of antifungal genes or through host induced silencing of fungal genes. Identification of resistance associated markers, combined with other molecular tools to turn off aflatoxin production during the fungal-host interaction has now become reliable means of developing crops resistant to aflatoxin contamination. Moreover, with significant information being currently developed on the genomics and transcriptomics of various commodities affected by toxin contamination, additional knowledge of host-fungal

interactions at the molecular level will identify control strategies that will be most effective in eliminating aflatoxin contamination worldwide. In this regard, availability of modern targeted gene editing in susceptible plants should pave the way for incorporating resistance to aflatoxin contamination. To improve the overall food and feed safety free of dangerous aflatoxin levels, it is not only necessary to minimise or eliminate the preharvest contamination levels but also extend constant postharvest monitoring to avoid fungal growth and toxin production during storage and handling. Although all the control measures, one at a time, discussed in this review will significantly minimise the occurrence of aflatoxin contamination in susceptible crops, it is unlikely to result in complete elimination of this problem because of (1) the adaptability of the fungus to broad-range of environments during crop production and during postharvest handling and storage, and (2) the complex role of the ubiquitous, opportunistic, saprophytic fungus in host plant-fungal pathogen relationship. Thus, it is essential to adopt multiple, yet coordinated, and complementary processes to control *A. flavus* and aflatoxin contamination using agronomic, genetic, biological, biotechnological, and molecular approaches.

## Acknowledgements

The authors are grateful to Suhad Wojkowski for assistance with the preparation of this manuscript.

## References

- Abdel-Hadi, A., 2011. Molecular ecology of *Aspergillus* section flavi species: approaches to understand the role of aflatoxin genes in aflatoxin biosynthesis. PhD-thesis, Cranfield University, Cranfield, UK.
- Abdel-Hadi, A., Caley, D.P., Carter, D.R.F. and Magan, R., 2011. Control of aflatoxin production of *Aspergillus flavus* and *Aspergillus parasiticus* using RNA silencing technology by targeting *afD* (*nor-1*) gene. *Toxins* 3: 647-659.
- Abdel-Hadi, A., Carter, D. and Magan, N., 2010. Temporal monitoring of the *nor-1* (*afID*) gene of *Aspergillus flavus* in relation to aflatoxin B<sub>1</sub> production during storage of peanuts under different environmental conditions. *Journal of Applied Microbiology* 109: 1914-1922.
- Abdel-Hadi, A., Schmidt-Heydt, M., Parra, R., Geisen, R. and Magan, N., 2012. A systems approach to model the relationship between aflatoxin gene cluster expression, environmental factors, growth and toxin production by *Aspergillus flavus*. *Journal of the Royal Society of London Interface* 9: 757-767.
- Adams, T.H. and Yu, J.H., 1998. Coordinate control of secondary metabolite production and asexual sporulation in *Aspergillus nidulans*. *Current Opinion in Microbiology* 1: 674-677.
- Amaike, S. and Keller, N.P., 2009. Distinct roles for VeA and LaeA in development and pathogenesis of *Aspergillus flavus*. *Eukaryotic Cell* 8: 1051-1060.
- Amare, M.G. and Keller, N.P., 2014. Molecular mechanisms of *Aspergillus flavus* secondary metabolism and development. *Fungal Genetics and Biology* 66: 11-18.
- Andersen, M.R., Nielsen, J.B., Klitgaard, A., Petersen, L.M., Zachariassen, M., Hansen, T.J., Blicher, L.H., Gotfredsen, C.H., Larsen, T.O., Nielsen, K.F. and Mortensen, U.H., 2013. Accurate prediction of secondary metabolite gene clusters in filamentous fungi. *Proceedings of the National Academy of Sciences of the USA* 110: E99-E107.
- Anderson, J.G. and Smith, J.E., 1972. The effects of elevated temperatures on spore swelling and germination in *Aspergillus niger*. *Canadian Journal of Microbiology* 18: 289-297.
- Aoki-Kinoshita, K.F. and Kanehisa, M., 2007. Gene annotation and pathway mapping in KEGG. *Methods in Molecular Biology* 396: 71-91.
- Arias, R.S., Dang, P.M. and Sobolev, V.S., 2015. RNAi-mediated control of aflatoxins in peanut: method to analyze mycotoxin production and transgene expression in the peanut/*Aspergillus* pathosystem. *Journal of Visualized Experiments* 106: e53398.
- Atehkneng, J., Ojiambo, P.S., Donner, M., Ikotun, T., Sikora, R.A., Cotty, P.J. and Bandyopadhyay, R., 2008. Distribution and toxigenicity of *Aspergillus* species isolated from maize kernels from three agro-ecological zones in Nigeria. *International Journal of Food Microbiology* 122: 74-84.
- Bai, Y., Lan, F., Yang, W., Zhang, F., Yang, K., Li, Z., Gao, P. and Wang, S., 2015. sRNA profiling in *Aspergillus flavus* reveals differentially expressed miRNA-like RNAs response to water activity and temperature. *Fungal Genetics and Biology* 81: 113-119.
- Baidya, S., Duran, R.M., Lohmar, J.M., Harris-Coward, P.W., Cary, J.W., Hong, S.Y., Roze, L.V., Linz, J.E. and Calvo, A.M., 2014. VeA is associated with the response to oxidative stress in the aflatoxin producer *Aspergillus flavus*. *Eukaryotic Cell* 13: 1095-1103.
- Baker, R., Brown, R.L., Cleveland, T.E., Chen, Z.-Y. and Fakhoury, A., 2009. A maize lectin-like protein with antifungal activity against *Aspergillus flavus*. *Journal of Food Protection* 72: 120-127.
- Barra, P., Etcheverry, M. and Nesci, A., 2015a. Efficacy of 2, 6-di (t-butyl)-p-cresol (BHT) and the entomopathogenic fungus *Purpureocillium lilacinum*, to control *Tribolium confusum* and to reduce aflatoxin B<sub>1</sub> in stored maize. *Journal of Stored Product Research* 64: 72-79.
- Barra, P., Etcheverry, M. and Nesci, A., 2015b. Improvement of the insecticidal capacity of two *Purpureocillium lilacinum* strains against *Tribolium confusum*. *Insects* 6: 206-223.
- Barra, P., Rosso, L., Nesci, A. and Etcheverry, M., 2013. Isolation and identification of entomopathogenic fungi and their evaluation against *Tribolium confusum*, *Sitophilus zeamais*, and *Rhizopertha dominica* in stored maize. *Journal of Pest Science* 86: 217-226.
- Barrow, C.J. and Sedlock, D.M., 1994. 1'-(2-Phenyl-ethylene)-ditryptophenalanine, a new dimeric diketopiperazine from *Aspergillus flavus*. *Journal of Natural Products* 57: 1239-1244.
- Battilani, P., Camardo Leggieri, M., Rossi, V. and Giorni, P., 2013. AFLA-maize, a mechanistic model for *Aspergillus flavus* infection and aflatoxin B<sub>1</sub> contamination in maize. *Computers and Electronics in Agriculture* 94: 38-46.



- Bayram, O. and Braus, G.H., 2012. Coordination of secondary metabolism and development in fungi: the velvet family of regulatory proteins. *FEMS Microbiology Reviews* 36: 1-24.
- Bayram, O., Krappmann, S., Ni, M., Bok, J.W., Helmstaedt, K., Valerius, O., Braus-Stromeier, S., Kwon, N.J., Keller, N.P., Yu, J.H. and Braus, G.H., 2008. VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science* 320: 1504-1506.
- Bedre, R., Rajasekaran, K., Mangu, V.R., Sanchez Timm, L.E., Bhatnagar, D. and Baisakh, N., 2015. Genome-wide transcriptome analysis of cotton (*Gossypium hirsutum* L.) identifies candidate gene signatures in response to aflatoxin producing fungus *Aspergillus flavus*. *PLoS ONE* 10: e0138025.
- Bhatnagar, D., Ehrlich, K.C. and Cleveland, T.E., 2003. Molecular genetic analysis and regulation of aflatoxin biosynthesis. *Applied Microbiology and Biotechnology* 61: 83-93.
- Bhatnagar, D., Rajasekaran, K., Cary, J.W., Brown, R.L., Yu, J. and Cleveland, T.E., 2008a. Molecular approaches to development of resistance to preharvest aflatoxin contamination. In: Leslie, J.F., Bandyopadhyay, R. and Visconti, A. (eds.) *Mycotoxins: detection methods, management, public health and agricultural trade*. CABI, Cambridge, MA, USA, pp. 257-276.
- Bhatnagar, D., Rajasekaran, K., Payne, G., Brown, R., Yu, J. and Cleveland, T., 2008b. The 'omics' tools: genomics, proteomics, metabolomics and their potential for solving the aflatoxin contamination problem. *World Mycotoxin Journal* 1: 3-12.
- Bhatnagar-Mathur, P., Sunkara, S., Bhatnagar-Panwar, M., Waliyar, F. and Sharma, K.K., 2015. Biotechnological advances for combating *Aspergillus flavus* and aflatoxin contamination in crops. *Plant Science* 234: 119-132.
- Brakhage, A.A. and Schroeckh, V., 2011. Fungal secondary metabolites – strategies to activate silent gene clusters. *Fungal Genetics and Biology* 48: 15-22.
- Brakhage, A.A., 2013. Regulation of fungal secondary metabolism. *Nature Reviews Microbiology* 11: 21-32.
- Brown, D.W., Yu, J.H., Kelkar, H.S., Fernandes, M., Nesbitt, T.C., Keller, N.P., Adams, T.H. and Leonard, T.L., 1996. Twenty-five coregulated transcripts define a sterigmatocystin gene cluster in *Aspergillus nidulans*. *Proceedings of the National Academy of Sciences of the USA* 93: 1418-1422.
- Brown, R.L., Bhatnagar, D., Herrera, E. and Bowman, J.E., 2015. Aflatoxins in maize and other crops. *World Mycotoxin Journal* 8: 122.
- Brown, R.L., Chen, Z., Menkir, A. and Cleveland, T.E., 2006. Proteomics to identify resistance factors in corn – a review. *Mycotoxin Research* 22: 22-26.
- Brown, R.L., Chen, Z.-Y., Cleveland, T.E., Menkir, A. and Fakhoury, A., 2009. Identification of maize breeding markers through investigations of proteins associated with aflatoxin-resistance. In: Appell, M., Kendra, D.F. and Trucksess, M.W. (eds.) *Mycotoxin prevention and control in agriculture*. ACS Symposium Series. Vol. 1031. American Chemical Society, Washington, DC, USA, pp. 157-165.
- Brown, R.L., Chen, Z.Y., Gembeh, S.V., Cleveland, T.E., Bhatnagar, D. and Howard, K., 2004. Identification of natural resistance in corn against mycotoxin-producing fungi. In: Mohan, R.M. (ed.) *Research advances in food science*. Global Research Network, Kerala, India, pp. 85-96.
- Brown, R.L., Chen, Z.Y., Warburton, M., Luo, M., Menkir, A., Fakhoury, A. and Bhatnagar, D., 2010. Discovery and characterization of proteins associated with aflatoxin-resistance: Evaluating their potential as breeding markers. *Toxins* 2: 919-933.
- Burow, G.B., Nesbitt, T.C., Dunlap, J. and Keller, N.P., 1997. Seed lipoxygenase products modulate *Aspergillus* mycotoxin biosynthesis. *Molecular Plant-Microbe Interactions* 10: 380-387.
- Calvo, A.M., Hinze, L.L., Gardner, H.W. and Keller, N.P., 1999. Sporogenic effect of polyunsaturated fatty acids on development of *Aspergillus* spp. *Applied and Environmental Microbiology* 65: 3668-3673.
- Calvo, A.M., Wilson, R.A., Bok, J.W. and Keller, N.P., 2002. Relationship between secondary metabolism and fungal development. *Microbiology and Molecular Biology Reviews* 66: 447-459.
- Campbell, K.W. and White, D.G., 1995. Evaluation of corn genotypes for resistance to *Aspergillus* ear rot, kernel infection, and aflatoxin production. *Plant Disease* 79: 1039-1045.
- Cary, J.W., Harris-Coward, P.Y., Ehrlich, K.C., Di Mavungu, J.D., Malysheva, S.V., De Saeger, S., Dowd, P.F., Shantappa, S., Martens, S.L. and Calvo, A.M., 2014. Functional characterization of a veA-dependent polyketide synthase gene in *Aspergillus flavus* necessary for the synthesis of asparosone, a sclerotium-specific pigment. *Fungal Genetics and Biology* 64: 25-35.
- Cary, J.W., O'Brian, G.R., Nelsen, D.M., Nierman, W., Harris-Coward, P., Yu, J., Bhatnagar, D., Cleveland, T.E., Payne, G.A. and Calvo, A.M., 2007. Elucidation of veA-dependent genes associated with aflatoxin and sclerotial production in *Aspergillus flavus* by functional genomics. *Applied Microbiology and Biotechnology* 76: 1107-1118.
- Cary, J.W., Han, Z., Yin, Y., Lohmar, J.M., Shantappa, S., Harris-Coward, P.Y., Mack, B., Ehrlich, K.C., Wei, Q., Arroyo-Manzanares, N., Uka, V., Vanhaecke, L., Bhatnagar, D., Yu, J., Nierman, W.C., Johns, M.A., Sorensen, D., Shen, H., De Saeger, S., Di Mavungu, J.D. and Calvo, A.M., 2015a. Transcriptome analysis of *Aspergillus flavus* reveals veA-dependent regulation of secondary metabolite gene clusters, including the novel aflavarin cluster. *Eukaryotic Cell* 14: 983-997.
- Cary, J.W., Harris-Coward, P., Scharfenstein, L., Mack, B.M., Chang, P.K., Wei, Q., Lebar, M., Carter-Wientjes, C., Majumdar, R., Mitra, C., Banerjee, S. and Chanda, A., 2017. The *Aspergillus flavus* homeobox gene, hbx1, is required for development and aflatoxin production. *Toxins* 9: e315.
- Cary, J.W., Harris-Coward, P.Y., Ehrlich, K.C., Mack, B.M., Kale, S.P., Larey, C. and Calvo, A.M., 2012. NsdC and NsdD affect *Aspergillus flavus* morphogenesis and aflatoxin production. *Eukaryotic Cell* 11: 1104-1111.
- Cary, J.W., Rajasekaran, K., Brown, R.L., Luo, M., Chen, Z.Y. and Bhatnagar, D., 2011. Developing resistance to aflatoxin in maize and cottonseed. *Toxins* 3: 678-696.
- Cary, J.W., Rajasekaran, K., Jaynes, J.M. and Cleveland, T.E., 2000. Transgenic expression of a gene encoding a synthetic antimicrobial peptide results in inhibition of fungal growth *in vitro* and *in planta*. *Plant Science* 154: 171-181.

- Cary, J.W., Rajasekaran, K., Yu, J., Brown, R.L., Bhatnagar, D. and Cleveland, T.E., 2009. Transgenic approaches for pre-harvest control of mycotoxin contamination in crop plants. *World Mycotoxin Journal* 2: 203-214.
- Cary, J.W., Uka, V., Han, Z., Buyst, D., Harris-Coward, P.Y., Ehrlich, K.C., Wei, Q., Bhatnagar, D., Dowd, P.F., Martens, S.L., Calvo, A.M., Martins, J.C., Vanhaecke, L., Coenye, T., De Saeger, S. and Di Mavungu, J.D., 2015b. An *Aspergillus flavus* secondary metabolic gene cluster containing a hybrid PKS-NRPS is necessary for synthesis of the 2-pyridones, leporins. *Fungal Genetics and Biology* 81: 88-97.
- Council for Agricultural and Science Technology (CAST), 2003. Aflatoxins and other mycotoxins: an agricultural perspective. CAST Reports, No. 80.
- Chalivendra, S.C., DeRobertis, C., Chang, P.K. and Damann, K.E., 2017. Cyclopiazonic acid is a pathogenicity factor for *Aspergillus flavus* and a promising target for screening germplasm for ear rot resistance. *Molecular Plant-Microbe Interactions* 30: 361-373.
- Chang, P.K., 2003. The *Aspergillus parasiticus* protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR. *Molecular Genetics and Genomics* 268: 711-719.
- Chang, P.K., Bennett, J.W. and Cotty, P.J., 2002. Association of aflatoxin biosynthesis and sclerotial development in *Aspergillus parasiticus*. *Mycopathologia* 153: 41-48.
- Chang, P.K., Ehrlich, K.C. and Fujii, I., 2009a. Cyclopiazonic acid biosynthesis of *Aspergillus flavus* and *Aspergillus oryzae*. *Toxins* 1: 74-99.
- Chang, P.K., Horn, B.W. and Dorner, J.W., 2009b. Clustered genes involved in cyclopiazonic acid production are next to the aflatoxin biosynthesis gene cluster in *Aspergillus flavus*. *Fungal Genetics and Biology* 46: 176-182.
- Chang, P.K., Scharfenstein, L.L., Li, P. and Ehrlich, K.C., 2013. *Aspergillus flavus* VelB acts distinctly from VeA in conidiation and may coordinate with FluG to modulate sclerotial production. *Fungal Genetics and Biology* 58-59: 71-79.
- Chang, P.K., Scharfenstein, L.L., Mack, B. and Ehrlich, K.C., 2012. Deletion of the *Aspergillus flavus* orthologue of *A. nidulans* fluG reduces conidiation and promotes production of sclerotia but does not abolish aflatoxin biosynthesis. *Applied and Environmental Microbiology* 78: 7557-7563.
- Chang, P.-K., Scharfenstein, L.L., Mack, B., Yu, J. and Ehrlich, K.C., 2014. Transcriptomic profiles of *Aspergillus flavus* CA42, a strain that produces small sclerotia, by decanal treatment and after recovery. *Fungal Genetics and Biology* 68, Suppl. C: 39-47.
- Chen, Z.Y., Brown, R.L., Damann, K.E. and Cleveland, T.E., 2002. Identification of unique or elevated levels of kernel proteins in aflatoxin-resistant maize genotypes through proteome analysis. *Phytopathology* 92: 1084-1094.
- Chen, Z.Y., Brown, R.L., Damann, K.E. and Cleveland, T.E., 2007. Identification of maize kernel endosperm proteins associated with resistance to aflatoxin contamination by *Aspergillus flavus*. *Phytopathology* 97: 1094-1103.
- Chen, Z.Y., Brown, R.L., Damann, K.E. and Cleveland, T.E., 2010. PR10 expression in maize and its effect on host resistance against *Aspergillus flavus* infection and aflatoxin production. *Molecular Plant Pathology* 11: 69-81.
- Chen, Z.Y., Brown, R.L., Lax, A.R., Guo, B.Z., Cleveland, T.E. and Russin, J.S., 1998. Resistance to *Aspergillus flavus* in corn kernels is associated with a 14-kDa protein. *Phytopathology* 88: 276-281.
- Chen, Z.Y., Brown, R.L., Rajasekaran, K., Damann, K.E. and Cleveland, T.E., 2006. Identification of a maize kernel pathogenesis-related protein and evidence for its involvement in resistance to *Aspergillus flavus* infection and aflatoxin production. *Phytopathology* 96: 87-95.
- Chen, Z.Y., Rajasekaran, K., Brown, R.L., Sayler, R.J. and Bhatnagar, D., 2015. Discovery and confirmation of genes/proteins associated with maize aflatoxin resistance. *World Mycotoxin Journal* 8: 211-224.
- Chen, Z.Y., Warburton, M.L., Hawkins, L., Wei, Q., Ruarung, Y., Brown, R.L., Zhang, L. and Bhatnagar, D., 2016. Production of the 14 kDa trypsin inhibitor protein is important for maize resistance against *Aspergillus flavus* infection/aflatoxin accumulation. *World Mycotoxin Journal* 9: 215-228.
- Chitarrini, G., Nobili, C., Pinzari, F., Antonini, A., De Rossi, P., Del Fiore, A., Procacci, S., Tolaini, V., Scala, V., Scarpari, M. and Reverberi, M., 2014. Buckwheat achenes antioxidant profile modulates *Aspergillus flavus* growth and aflatoxin production. *International Journal of Food Microbiology* 189: 1-10.
- Cleveland, T.E., Yu, J., Fedorova, N., Bhatnagar, D., Payne, G.A., Nierman, W.C. and Bennett, J.W., 2009. Potential of *Aspergillus flavus* genomics for applications in biotechnology. *Trends in Biotechnology* 27: 151-157.
- Clevenger, J., Marasigan, K., Liakos, V., Sobolev, V., Vellidis, G., Holbrook, C. and Ozias-Akins, P., 2016. RNA sequencing of contaminated seeds reveals the state of the seed permissive for pre-harvest aflatoxin contamination and points to a potential susceptibility factor. *Toxins* 8: 317.
- Cole, R.J., Jarvis, B.B. and Schweikert, M.A., 2003. Handbook of secondary fungal metabolites. Elsevier Science, New York, NY, USA.
- Cotty, P., 1989. Virulence and cultural characteristics of two *Aspergillus flavus* strains pathogenic on cotton. *Phytopathology* 79: 808-814.
- Cross, H.Z. and Zuber, M.S., 1972. Prediction of flowering dates in maize based on different methods of estimating thermal units. *Agronomy Journal* 64: 351-355.
- DaMatta, F.M., Grandis, A., Arenque, B.C. and Buckeridge, M.S., 2010. Impacts of climate changes on crop physiology and food quality. *Food Research International* 43: 1814-1823.
- Davis, N.D., Currier, C.G. and Diener, U.L., 1986. Aflatoxin contamination of corn hybrids in Alabama. *Cereal Chemistry* 60: 46-50.
- De Lucca, A.J., Jacks, T.J. and Broekaert, W.J., 1999. Fungicidal and binding properties of three plant peptides. *Mycopathologia* 144: 87-91.
- De Vries, R.P., Riley, R., Wiebenga, A., Aguilar-Osorio, G., Amillis, S., Uchima, C.A., Anderluh, G., Asadollahi, M., Askin, M., Barry, K., Battaglia, E., Bayram, O., Benocci, T., Braus-Stromeier, S.A., Caldana, C., Canovas, D., Cerqueira, G.C., Chen, F., Chen, W., Choi, C., Clum, A., Dos Santos, R.A., Damasio, A.R., Diallinas, G., Emri, T., Fekete, E., Flipphi, M., Freyberg, S., Gallo, A., Gournas, C., Habgood, R., Hainaut, M., Harispe, M.L., Henrissat, B., Hilden, K.S., Hope, R., Hossain, A., Karabika, E., Karaffa, L., Karanyi, Z., Krasevec, N., Kuo, A., Kusch, H., LaButti, K., Lagendijk, E.L., Lapidus, A., Lévassieur, A., Lindquist, E., Lipzen, A., Logrieco, A.F., MacCabe, A.,

- Makela, M.R., Malavazi, I., Melin, P., Meyer, V., Mielnichuk, N., Miskei, M., Molnar, A.P., Mule, G., Ngan, C.Y., Orejas, M., Orosz, E., Ouedraogo, J.P., Overkamp, K.M., Park, H.S., Perrone, G., Piumi, F., Punt, P.J., Ram, A.F., Ramon, A., Rauscher, S., Record, E., Riano-Pachon, D.M., Robert, V., Rohrig, J., Ruller, R., Salamov, A., Salih, N.S., Samson, R.A., Sandor, E., Sanguinetti, M., Schutze, T., Sepcic, K., Shelest, E., Sherlock, G., Sophianopoulou, V., Squina, F.M., Sun, H., Susca, A., Todd, R.B., Tsang, A., Unkles, S.E., Van de Wiele, N., Van Rossen-Uffink, D., Oliveira, J.V., Vesth, T.C., Visser, J., Yu, J.H., Zhou, M., Andersen, M.R., Archer, D.B., Baker, S.E., Benoit, I., Brakhage, A.A., Braus, G.H., Fischer, R., Frisvad, J.C., Goldman, G.H., Houbraken, J., Oakley, B., Pocsí, I., Scazzocchio, C., Seiboth, B., VanKuyk, P.A., Wortman, J., Dyer, P.S. and Grigoriev, I.V., 2017. Comparative genomics reveals high biological diversity and specific adaptations in the industrially and medically important fungal genus *Aspergillus*. *Genome Biology* 18: 28.
- DeGray, G., Rajasekaran, K., Smith, F., Sanford, J. and Daniell, H., 2001. Expression of an antimicrobial peptide via the chloroplast genome to control phytopathogenic bacteria and fungi. *Plant Physiology* 127: 852-862.
- Diener, U.L., Cole, R.J., Sanders, T.H., Payne, G.A., Lee, L.S. and Klich, M.A., 1987. Epidemiology of aflatoxin formation by *Aspergillus flavus*. *Annual Review of Phytopathology* 25: 249-270.
- Donner, M., Atehnkeng, J., Sikora, R.A., Bandyopadhyay, R. and Cotty, P.J., 2010. Molecular characterization of atoxigenic strains for biological control of aflatoxins in Nigeria. *Food Additives and Contaminants Part A* 27: 576-590.
- Dowd, P.F., 1988. Synergism of aflatoxin B<sub>1</sub> toxicity with the co-occurring fungal metabolite kojic acid to two caterpillars. *Entomologia Experimentalis et Applicata* 47: 69-71.
- Driever, S.M. and Kromdijk, J., 2013. Will C<sub>3</sub> crops enhanced with the C<sub>4</sub> CO<sub>2</sub>-concentrating mechanism live up to their full potential (yield)? *Journal of Experimental Botany* 64: 3925-3935.
- Du, W., Obrian, G.R. and Payne, G.A., 2007. Function and regulation of aflJ in the accumulation of aflatoxin early pathway intermediate in *Aspergillus flavus*. *Food Additives and Contaminants* 24: 1043-1050.
- Duran, R.M., Cary, J.W. and Calvo, A.M., 2007. Production of cyclopiazonic acid, aflatrem, and aflatoxin by *Aspergillus flavus* is regulated by veA, a gene necessary for sclerotial formation. *Applied Microbiology and Biotechnology* 73: 1158-1168.
- Egner, P.A., Yu, X., Johnson, J.K., Nathasingh, C.K., Groopman, J.D., Kensler, T.W. and Roebuck, B.D., 2003. Identification of aflatoxin M<sub>1</sub>-N<sub>7</sub>-guanine in liver and urine of tree shrews and rats following administration of aflatoxin B<sub>1</sub>. *Chemical Research Toxicology* 16: 1174-1180.
- Ehmann, D.E., Gehring, A.M. and Walsh, C.T., 1999. Lysine biosynthesis in *Saccharomyces cerevisiae*: mechanism of  $\alpha$ -amino adipate reductase (Lys2) involves posttranslational phosphopantetheinylation by Lys5. *Biochemistry* 38: 6171-6177.
- Ehrlich, K.C. and Mack, B.M., 2014. Comparison of expression of secondary metabolite biosynthesis cluster genes in *Aspergillus flavus*, *A. parasiticus*, and *A. oryzae*. *Toxins* 6: 1916-1928.
- Ehrlich, K.C., Montalbano, B.G. and Cary, J.W., 1999. Binding of the C6-zinc cluster protein, AFLR, to the promoters of aflatoxin pathway biosynthesis genes in *Aspergillus parasiticus*. *Gene* 230: 249-257.
- Fakhoury, A.M. and Woloshuk, C.P., 2001. Inhibition of growth of *Aspergillus flavus* and fungal alpha-amylases by a lectin-like protein from *Lablab purpureus*. *Molecular Plant-Microbe Interactions* 14: 955-961.
- Faustinelli, P.C., Palencia, E.R., Sobolev, V.S., Horn, B.W., Sheppard, H.T., Lamb, M.C., Wang, X.M., Scheffler, B.E., Martinez Castillo, J. and Arias, R.S., 2017. Study of the genetic diversity of the aflatoxin biosynthesis cluster in *Aspergillus* section Flavi using insertion/deletion markers in peanut seeds from Georgia, USA. *Mycologia* 109: 200-209.
- Faustinelli, P.C., Wang, X.M., Palencia, E.R. and Arias, R.S., 2016. Genome sequences of eight *Aspergillus flavus* spp. and one *A. parasiticus* sp., isolated from peanut seeds in Georgia. *Genome Announcements* 4: e00278-00216.
- Fernandes, M., Keller, N.P. and Adams, T.H., 1998. Sequence-specific binding by *Aspergillus nidulans* AflR, a C6 zinc cluster protein regulating mycotoxin biosynthesis. *Molecular Microbiology* 28: 1355-1365.
- Flaherty, J.E., Weaver, M.E., Payne, G.A. and Woloshuk, C.P., 1995. A beta-glucuronidase reporter gene construct for monitoring aflatoxin biosynthesis in *A. flavus*. *Applied and Environmental Microbiology* 61: 2482-2486.
- Forseth, R.R., Amaike, S., Schwenk, D., Affeldt, K.J., Hoffmeister, D., Schroeder, F.C. and Keller, N.P., 2013. Homologous NRPS-like gene clusters mediate redundant small-molecule biosynthesis in *Aspergillus flavus*. *Angewandte Chemie* 52: 1590-1594.
- Fountain, J.C., Bajaj, P., Nayak, S.N., Yang, L., Pandey, M.K., Kumar, V., Jayale, A.S., Chitkani, A., Lee, R.D., Kemerait, R.C., Varshney, R.K. and Guo, B., 2016. Responses of *Aspergillus flavus* to oxidative stress are related to fungal development regulator, antioxidant enzyme, and secondary metabolite biosynthetic gene expression. *Frontiers in Microbiology* 7: 2048.
- Gallagher, R.T. and Wilson, B.J., 1979. Aflatrem, the tremorgenic mycotoxin from *Aspergillus flavus*. *Mycopathologia* 66: 183-185.
- Georgianna, D.R. and Payne, G.A., 2009. Genetic regulation of aflatoxin biosynthesis: from gene to genome. *Fungal Genetics and Biology* 46: 113-125.
- Georgianna, D.R., Hawkrigde, A.M., Muddiman, D.C. and Payne, G.A., 2008. Temperature-dependent regulation of proteins in *Aspergillus flavus*: whole organism stable isotope labelling by amino acids. *Journal of Proteome Research* 7: 2973-2979.
- Georgianna, D.R., Fedorova, N.D., Burroughs, J.L., Dolezal, A.L., Bok, J.W., Horowitz-Brown, S., Woloshuk, C.P., Yu, J., Keller, N.P. and Payne, G.A., 2010. Beyond aflatoxin: four distinct expression patterns and functional roles associated with *Aspergillus flavus* secondary metabolism gene clusters. *Molecular Plant Pathology* 11: 213-226.
- Gibbons, J.G., Salichos, L., Slot, J.C., Rinker, D.C., McGary, K.L., King, J.G., Klich, M.A., Tabb, D.L., McDonald, W.H. and Rokas, A., 2012. The evolutionary imprint of domestication on genome variation and function of the filamentous fungus *Aspergillus oryzae*. *Current Biology* 22: 1403-1409.
- Gilbert, M.K., Mack, B.M., Wei, Q., Bland, J.M., Bhatnagar, D. and Cary, J.W., 2016. RNA sequencing of an nsdC mutant reveals global regulation of secondary metabolic gene clusters in *Aspergillus flavus*. *Microbiological Research* 182: 150-161.



- Giordana, B.E.G. Nones, J. and Scussel, V.M., 2012. Susceptibility of the in-shell Brazil Nut mycoflora and aflatoxin contamination to ozone gas treatment during storage. *Journal of Agricultural Sciences* 4: 8.
- Giorni, P., Battilani, P. and Magan, N., 2008a. Effect of solute and matric potential on *in vitro* growth and sporulation of strains from a new population of *Aspergillus flavus* isolated in Italy. *Fungal Ecology* 1: 102-106.
- Giorni, P., Battilani, P., Pietri, A. and Magan, N., 2008b. Effect of  $a_w$  and CO<sub>2</sub> level on *Aspergillus flavus* growth and aflatoxin production in high moisture maize post-harvest. *International Journal of Food Microbiology* 122: 109-113.
- Giorni, P., Camardo Legierri, M., Magan, N. and Battilani, P., 2012. Comparison of temperature and moisture requirements for sporulation of *Aspergillus flavus* sclerotia on natural and artificial substrates. *Fungal Biology* 116: 637-642.
- Giorni, P., Magan, N., Pietri, A., Bertuzzi, T. and Battilani, P., 2007. Studies on *Aspergillus* section Flavi isolated from maize in northern Italy. *International Journal of Food Microbiology* 113: 330-338.
- Goodwin, S., McPherson, J.D. and McCombie, W.R., 2016. Coming of age: ten years of next-generation sequencing technologies. *Nature Reviews, Genetics* 17: 333-351.
- Greaves, J.A., 1996. Improving suboptimal temperature tolerance in maize—the search for variation. *Journal of Experimental Botany* 47: 307-324.
- Gressel, J. and Polturak, G., 2017. Suppressing aflatoxin biosynthesis is not a breakthrough if not useful – a perspective. *Pest Management Science* 74: 17-21.
- Griffin, J., Smith, G.P., Smith, E. and Robinson, T., 2001. Population patterns of *Aspergillus flavus* group and *A. niger* group in field soils. *Soil Biology and Biochemistry* 33: 253-257.
- Guo, B.Z., Yu, J., Holbrook, C.C., Lee, R.D. and Lynch, R.E., 2003. Application of differential display RT-PCR and EST/Microarray technologies to the analysis of gene expression in response to drought stress and elimination of aflatoxin contamination in corn and peanut. *Journal of Toxicology* 22: 287-312.
- Guo, B.Z., Chen, Z.Y., Brown, R.L., Lax, A.R., Cleveland, T.E., Russin, J.S., Mehta, A.D., Selitrennikoff, C.P. and Widstrom, N.W., 1997. Germination induces accumulation of specific proteins and antifungal activities in corn kernels. *Phytopathology* 87: 1174-1178.
- Guo, B.Z., Krakowsky, M.D., Ni, X., Scully, B.T., Lee, R.D., Coy, A.E. and Widstrom, N.W., 2011. Registration of maize inbred line GT603. *Journal of Plant Registrations* 5: 211-214.
- Hicks, J.K., Yu, J.H., Keller, N.P. and Adams, T.H., 1997. *Aspergillus* sporulation and mycotoxin production both require inactivation of the FadA G alpha protein-dependent signalling pathway. *EMBO Journal* 16: 4916-4923.
- Holbrook, C.C., Kvien, C.K., Rucker, K.S., Wilson, D.M., Hook, J.E. and Matheron, M.E., 2000. Preharvest aflatoxin contamination in drought-tolerant and drought-intolerant peanut genotypes. *Peanut Science* 27: 45-48.
- Holmquist, G.U., Walker, H.W. and Stahr, H.M., 1983. Influence of temperature, pH, water activity and antifungal agents on growth of *Aspergillus flavus* and *A. parasiticus*. *Journal of Food Science* 48: 778-782.
- Horn, B.W., Greene, R.L., Sobolev, V.S., Dorner, J.W., Powell, J.H. and Layton, R.C., 1996. Association of morphological and mycotoxin production with vegetative compatibility groups in *Aspergillus flavus*, *A. parasiticus*, and *A. tamarii*. *Mycologia* 88: 574-587.
- Horn, B.W., Moore, G.G. and Carbone, I., 2009. Sexual reproduction in *Aspergillus flavus*. *Mycologia* 101: 423-429.
- Hruska, Z., Rajasekaran, K., Yao, H., Kincaid, R., Darlington, D., Brown, R.L., Bhatnagar, D. and Cleveland, T.E., 2014. Co-inoculation of aflatoxigenic and non-aflatoxigenic strains of *Aspergillus flavus* to study fungal invasion, colonization, and competition in maize kernels. *Frontiers in Microbiology* 5: 122.
- Intergovernmental Panel on Climate Change (IPCC), 2013. Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, UK and New York, NY, USA, 1535 pp.
- Intergovernmental Panel on Climate Change (IPCC), 2014. Summary for policymakers. In: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R. and White, L.L. (eds.) Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. Contribution of working group II to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 1-32.
- Jacks, T.J., De Lucca, A.J., Rajasekaran, K., Stromberg, K. and Van Pee, K., 2000. Antifungal and peroxidative activities of nonheme chloroperoxidase in relation to transgenic plant protection. *Journal of Agricultural and Food Chemistry* 48: 4561-4564.
- Jones, R.K., Duncan, H.E. and Hamilton, P.B., 1981. Planting date, harvest date, and irrigation effects on infection and aflatoxin production by *Aspergillus flavus* in field corn. *Phytopathology* 71: 810-816.
- Jones, R.K., Duncan, H.E., Payne, G.A. and Leonard, K.J., 1980. Factors influencing infection by *Aspergillus flavus* in silk-inoculated corn. *Plant Disease* 64: 859-863.
- Kale, S.P., Milde, L., Trapp, M.K., Frisvad, J.C., Keller, N.P. and Bok, J.W., 2008. Requirement of LaeA for secondary metabolism and sclerotial production in *Aspergillus flavus*. *Fungal Genetics and Biology* 45: 1422-1429.
- Kato, N., Tokuoaka, M., Shinohara, Y., Kawatani, M., Uramoto, M., Seshime, Y., Fujii, I., Kitamoto, K., Takahashi, T., Takahashi, S., Koyama, Y. and Osada, H., 2011. Genetic safeguard against mycotoxin cyclopiazonic acid production in *Aspergillus oryzae*. *ChemBioChem* 12: 1376-1382.
- Khatodia, S., Bhatotia, K., Passricha, N., Khurana, S.M.P. and Tuteja, N., 2016. The CRISPR/Cas genome-editing tool: application in improvement of crops. *Frontiers in Plant Science* 7: 506.
- Kim, H., Han, K., Kim, K., Han, D., Jahng, K. and Chae, K., 2002. The veA gene activates sexual development in *Aspergillus nidulans*. *Fungal Genetics and Biology* 37: 72-80.
- Kim, J.H., Yu, J., Mahoney, N., Chan, K.L., Molyneux, R.J., Varga, J., Bhatnagar, D., Cleveland, T.E., Nierman, W.C. and Campbell, B.C., 2008. Elucidation of the functional genomics of antioxidant-based inhibition of aflatoxin biosynthesis. *International Journal of Food Microbiology* 122: 49-60.

- Koiso, Y., Li, Y., Iwasaki, S., Hanaoka, K., Kobayashi, T., Sonoda, R., Fujita, Y., Yaegashi, H. and Sato, Z., 1994. Ustiloxins, antimitotic cyclic peptides from false smut balls on rice panicles caused by *Ustilagoidea virens*. *Journal of Antibiotics* 47: 765-773.
- Kos, J., Mastilović, J., Hajnal, E.J. and Šarić, B., 2013. Natural occurrence of aflatoxins in maize harvested in Serbia during 2009-2012. *Food Control* 34: 31-34.
- Kumar, P., Mahato, D.K., Kamle, M., Mohanta, T.K. and Kang, S.G., 2016. Aflatoxins: a global concern for food safety, human health and their management. *Frontiers in Microbiology* 7: 2170.
- Langfelder, P. and Horvath, S., 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* 9: 559.
- Lanubile, A., Pasini, L. and Marocco, A., 2010. Differential gene expression in kernels and silks of maize lines with contrasting levels of ear rot resistance after *Fusarium verticillioides* infection. *Journal of Plant Physiology* 167: 1398-1406.
- Lee, S., Rajasekaran, K., Ramanarao, M.V., Bedre, R., Bhatnagar, D. and Baisakh, N., 2012. Identifying cotton (*Gossypium hirsutum* L.) genes induced in response to *Aspergillus flavus* infection. *Physiological and Molecular Plant Pathology* 80: 35-40.
- Lin, J.Q., Zhao, X.X., Zhi, Q.Q., Zhao, M. and He, Z.M., 2013. Transcriptomic profiling of *Aspergillus flavus* in response to 5-azacytidine. *Fungal Genetics and Biology* 56: 78-86.
- Liu, S.Y., Lin, J.Q., Wu, H.L., Wang, C.C., Huang, S.J., Luo, Y.F., Sun, J.H., Zhou, J.X., Yan, S.J., He, J.G., Wang, J. and He, Z.M., 2012. Bisulfite sequencing reveals that *Aspergillus flavus* holds a hollow in DNA methylation. *PLoS ONE* 7: e30349.
- Lohmar, J.M., Harris-Coward, P.Y., Cary, J.W., Dhingra, S. and Calvo, A.M., 2016. *rtfA*, a putative RNA-Pol II transcription elongation factor gene, is necessary for normal morphological and chemical development in *Aspergillus flavus*. *Applied Microbiology and Biotechnology* 100: 5029-5041.
- Luo, M., Liu, J., Lee, R.D., Scully, B.T. and Guo, B., 2010. Monitoring the expression of maize genes in developing kernels under drought stress using oligo-microarray. *Journal of Integrative Plant Biology* 52: 1059-1074.
- Luo, X., Affeldt, K.J. and Keller, N.P., 2016. Characterization of the far transcription factor family in *Aspergillus flavus*. *G3: Genes Genomes Genetics* 6: 3269-3281.
- Macheleidt, J., Mattern, D.J., Fischer, J., Netzke, T., Weber, J., Schroeckh, V., Valiante, V. and Brakhage, A.A., 2016. Regulation and role of fungal secondary metabolites. *Annual Review of Genetics* 50: 371-392.
- Magan, N., 1993. Use of sulphur dioxide to control fungi in stored grain. In: Navarro, S. and Donahaye, E. (eds.) *Proceedings of the International Conference on controlled atmosphere and fumigation in grain storages*. Capital Press, Jerusalem, Israel, pp. 163-171.
- Majumdar, R., Rajasekaran, K. and Cary, J.W., 2017a. RNA interference (RNAi) as a potential tool for control of mycotoxin contamination in crop plants: concepts and considerations. *Frontiers in Plant Science* 8: 200.
- Majumdar, R., Rajasekaran, K., Sickler, C., Lebar, M., Musungu, B.M., Fakhoury, A.M., Payne, G.A., Geisler, M., Carter-Wientjes, C., Wei, Q., Bhatnagar, D. and Cary, J.W., 2017b. The pathogenesis-related maize seed (*PRms*) gene plays a role in resistance to *Aspergillus flavus* infection and aflatoxin contamination. *Frontiers in Plant Science* 8: 1758.
- Marín, S., Sanchis, V., Ramos, A.J. and Magan, N., 1998. Ecological determinants for germination and growth of some *Aspergillus* and *Penicillium* spp. from maize grain. *Journal of Applied Microbiology* 84: 25-36.
- Martin Castaño, S., Medina, A. and Magan, N., 2017. Comparison of dry matter losses and aflatoxin B<sub>1</sub> contamination of paddy and brown rice stored naturally or after inoculation with *Aspergillus flavus* at different environmental conditions, *Journal of Stored Products Research* 73: 47-53.
- Marui, J., Yamane, N., Ohashi-Kunihiro, S., Ando, T., Terabayashi, Y., Sano, M., Ohashi, S., Ohshima, E., Tachibana, K., Higa, Y., Nishimura, M., Koike, H. and Machida, M., 2011. Kojic acid biosynthesis in *Aspergillus oryzae* is regulated by a Zn(II)(2) Cys(6) transcriptional activator and induced by kojic acid at the transcriptional level. *Journal of Bioscience and Bioengineering* 112: 40-43.
- Masanga, J.O., Matheka, J.M., Omer, R.A., Ommeh, S.C., Monda, E.O. and Alakonya, A.E., 2015. Downregulation of transcription factor aflR in *Aspergillus flavus* confers reduction to aflatoxin accumulation in transgenic maize with alteration of host plant architecture. *Plant Cell Reports* 34: 1379-1387.
- Medema, M.H., Blin, K., Cimermancic, P., De Jager, V., Zakrzewski, P., Fischbach, M.A., Weber, T., Takano, E. and Breitling, R., 2011. antiSMASH: rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. *Nucleic Acids Research* 39: W339-346.
- Medina, A., Akbar, A., Baazeem, A., Rodriguez, A. and Magan, N., 2017a. Climate change, food security and mycotoxins: do we know enough? *Fungal Biology Reviews* 31: 143-154.
- Medina, A., Gilbert, M.K., Mack, B.M., O'Brian, G.R., Rodriguez, A., Bhatnagar, D., Payne, G. and Magan, N., 2017b. Interactions between water activity and temperature on the *Aspergillus flavus* transcriptome and aflatoxin B<sub>1</sub> production. *International Journal of Food Microbiology* 256: 36-44.
- Medina, A., Rodríguez, A., Sultan, Y. and Magan, N., 2015. Climate change factors and *Aspergillus flavus*: effects on gene expression, growth and aflatoxin production. *World Mycotoxin Journal* 8: 171-179.
- Menkir, A., Brown, R.L., Bandyopadhyay, R. and Cleveland, T.E., 2008. Registration of six tropical maize germplasm lines with resistance to aflatoxin contamination. *Journal of Plant Registrations* 2: 246-250.
- Meyers, D.M., O'Brian, G., Du, W.L., Bhatnagar, D. and Payne, G.A., 1998. Characterization of *afII*, a gene required for conversion of pathway intermediates to aflatoxin. *Applied and Environmental Microbiology* 64: 3713-3717.
- Miglani, G.S., 2017. Genome editing in crop improvement: present scenario and future prospects. *Journal of Crop Improvement* 31: 453-559.

- Moore, G.G., Elliott, J.L., Singh, R., Dorner, J.W., Stone, E.A., Chulze, S.N., Barros, G.G., Naik, M.K., Wright, G.C., Hell, K. and Carbone, I., 2013. Sexuality generates diversity in the aflatoxin gene cluster: evidence on a global scale. *PLoS Pathogens* 9: e1003574.
- Moore, G.G., Mack, B.M. and Beltz, S.B., 2015. Genomic sequence of the aflatoxigenic filamentous fungus *Aspergillus nomius*. *BMC Genomics* 16: 551.
- Moore, G.G., Mack, B.M., Beltz, S.B. and Gilbert, M.K., 2016. Draft genome sequence of an aflatoxigenic *Aspergillus* species, *A. bombycis*. *Genome Biology and Evolution* 8: 3297-3300.
- Moore, J., Rajasekaran, K., Cary, J. and Chlan, C., 2017. Identification of resistance to *Aspergillus flavus* infection in cotton germplasm. *Journal of Crop Improvement* 31: 727-741.
- Moore, K.G., Price, M.S., Boston, R.S., Weissinger, A.K. and Payne, G.A., 2004. A chitinase from Tex6 maize kernels inhibits growth of *Aspergillus flavus*. *Phytopathology* 94: 82-87.
- Musungu, B., Bhatnagar, D., Brown, R.L., Fakhoury, A.M. and Geisler, M., 2015. A predicted protein interactome identifies conserved global networks and disease resistance subnetworks in maize. *Frontiers in Genetics* 6: 201.
- Musungu, B.M., Bhatnagar, D., Brown, R.L., Payne, G.A., O'Brian, G., Fakhoury, A.M. and Geisler, M., 2016. A network approach of gene co-expression in the *Zea mays/Aspergillus flavus* pathosystem to map host/pathogen interaction pathways. *Frontiers in Genetics* 7: 206.
- Mylona, K., Kogkaki, E., Sulyok, M. and Magan, N., 2014. Efficacy of gaseous ozone treatment on spore germination, growth and fumonisin production by *Fusarium verticillioides* *in vitro* and *in situ* in maize. *Journal of Stored Product Research* 59: 178-184.
- Mylroie, J.E., Ozkan, S., Shivaji, R., Windham, G.L., Alpe, M.N. and Williams, W.P., 2016. Identification and quantification of a toxigenic and non-toxigenic *Aspergillus flavus* strain in contaminated maize using quantitative real-time PCR. *Toxins* 8: e15.
- Nagano, N., Umemura, M., Izumikawa, M., Kawano, J., Ishii, T., Kikuchi, M., Tomii, K., Kumagai, T., Yoshimi, A., Machida, M., Abe, K., Shin-ya, K. and Asai, K., 2016. Class of cyclic ribosomal peptide synthetic genes in filamentous fungi. *Fungal Genetics and Biology* 86: 58-70.
- Nesci, A., Etcheverry, M. and Magan, N., 2004. Osmotic and matric potential effects on growth, sugar alcohol, and sugar accumulation in *Aspergillus* section *Flavi* strains from Argentina. *Journal of Applied Microbiology* 96: 965-972.
- Nicholson, M.J., Koulman, A., Monahan, B.J., Pritchard, B.L., Payne, G.A. and Scott, B., 2009. Identification of two aflatoxin biosynthesis gene loci in *Aspergillus flavus* and metabolic engineering of *Penicillium paxilli* to elucidate their function. *Applied and Environmental Microbiology* 75: 7469-7481.
- Nielsen, K., Payne, G.A. and Boston, R.S., 2001. Maize ribosome-inactivating protein inhibits normal development of *Aspergillus nidulans* and *Aspergillus flavus*. *Molecular Plant-Microbe Interactions* 14: 164-172.
- Nierman, W.C., Yu, J., Fedorova-Abrams, N.D., Losada, L., Cleveland, T.E., Bhatnagar, D., Bennett, J.W., Dean, R. and Payne, G.A., 2015. Genome sequence of *Aspergillus flavus* NRRL 3357, a strain that causes aflatoxin contamination of food and feed. *Genome Announcements* 3: 2.
- Northolt, M.D., Vanegmond, H.P. and Paulsch, W.E., 1977. Differences between *Aspergillus flavus* strains in growth and aflatoxin B<sub>1</sub> production in relation to water activity and temperature. *Journal of Food Protection* 40: 778-781.
- O'Brian, G.R., Fakhoury, A.M. and Payne, G.A., 2003. Identification of genes differentially expressed during aflatoxin biosynthesis in *Aspergillus flavus* and *Aspergillus parasiticus*. *Fungal Genetics and Biology* 39: 118-127.
- O'Brian, G.R., Georgianna, D.R., Wilkinson, J.R., Yu, J., Abbas, H.K., Bhatnagar, D., Cleveland, T.E., Nierman, W. and Payne, G.A., 2007. The effect of elevated temperature on gene transcription and aflatoxin biosynthesis. *Mycologia* 99: 232-239.
- Parisi, C., Tillie, P. and Rodriguez-Cerezo, E., 2016. The global pipeline of GM crops out to 2020. *Nature Biotechnology* 34: 31-36.
- Passone, M.A., Rosso, L.C., Ciancio, A. and Etcheverry, M., 2010. Detection and quantification of *Aspergillus* section *Flavi* spp. in stored peanuts by real-time PCR of *nor-1* gene, and effects of storage conditions on aflatoxin production. *International Journal of Food Microbiology* 138: 276-281.
- Paterson, R.R.M. and Lima, N., 2011. Further mycotoxin effects from climate change. *Food Research International* 44: 2555-2566.
- Payne, G.A., Nierman, W.C., Wortman, J.R., Pritchard, B.L., Brown, D., Dean, R.A., Bhatnagar, D., Cleveland, T.E., Machida, M. and Yu, J., 2006. Whole genome comparison of *Aspergillus flavus* and *A. oryzae*. *Medical Mycology* 44: S9-S11.
- Payne, G.A., Thompson, D.L., Lillehoj, E.B., Zuber, M.S. and Adkins, C.R., 1988. Effect of temperature on the preharvest infection of maize kernels by *Aspergillus flavus*. *Phytopathology* 78: 1376-1380.
- Pechanova, O., Pechan, T., Rodriguez, J.M., Williams, W.P. and Brown, A.E., 2013. A two-dimensional proteome map of the aflatoxigenic fungus *Aspergillus flavus*. *Proteomics* 13: 1513-1518.
- Piva, G., Battilani, P. and Pietri, A., 2006. Emerging issues in southern Europe: aflatoxins in Italy. In: Barug, D., Bhatnagar, D., Van Egmond, H.P., Van der Kamp, J.W., Van Ossenbruggen, W.A. and Visconti, A. (eds.) *The mycotoxin fact book, food and feed topics*. Wageningen Academic Publishers, Wageningen, the Netherlands, pp. 139-153.
- Porter, J.R. and Semenov, M.A., 2005. Crop responses to climatic variation. *Philosophical Transactions of the Royal Society B: Biological Sciences* 360: 2021-2035.
- Power, I.L., Dang, P.M., Sobolev, V.S., Orner, V.A., Powell, J.L., Lamb, M.C. and Arias, R.S., 2017. Characterization of small RNA populations in non-transgenic and aflatoxin-reducing-transformed peanut. *Plant Science* 257: 106-125.
- Price, M.S., Yu, J., Nierman, W.C., Kim, H.S., Pritchard, B., Jacobus, C.A., Bhatnagar, D., Cleveland, T.E. and Payne, G.A., 2006. The aflatoxin pathway regulator AflR induces gene transcription inside and outside of the aflatoxin biosynthetic cluster. *FEMS Microbiology Letters* 255: 275-279.
- Rajasekaran, K., Bhatnagar, D., Brown, R., Chen, Z.Y., Cary, J.W. and Cleveland, T.E., 2005a. Enhancing food safety: prevention of preharvest aflatoxin contamination. In: Ramasamy, C., Ramanathan, S. and Dhakshinamoorthy, M. (eds.) *Perspectives of agricultural research and development*. Tamil Nadu Agricultural University, Coimbatore, India, pp. 434-467.



- Rajasekaran, K., Cary, J. and Cleveland, T.E., 2006. Prevention of preharvest aflatoxin contamination through genetic engineering of crops. *Mycotoxin Research* 22: 118-124.
- Rajasekaran, K., Cary, J.W., Chen, Z.-Y., Brown, R.L. and Cleveland, T.E., 2008. Antifungal traits of a 14 kDa maize kernel trypsin inhibitor protein in transgenic cotton. *Journal of Crop Improvement* 22: 1-16.
- Rajasekaran, K., Cary, J.W., Chlan, C.A., Jaynes, J.M. and Bhatnagar, D., 2012. Strategies for controlling plant diseases and mycotoxin contamination using antimicrobial synthetic peptides. In: Rajasekaran, K., Cary, J.W., Jaynes, J.M. and Montesinos, E. (eds.) *Small wonders: peptides for disease control*. ACS symposium series. Vol. 1095. American Chemical Society, Washington, DC, USA, pp. 295-315.
- Rajasekaran, K., Cary, J.W., Jacks, T.J., Stromberg, K. and Cleveland, T.E., 2000. Inhibition of fungal growth *in planta* and *in vitro* by transgenic tobacco expressing a bacterial nonheme chloroperoxidase gene. *Plant Cell Reports* 19: 333-338.
- Rajasekaran, K., Cary, J.W., Jaynes, J.M. and Cleveland, T.E., 2005b. Disease resistance conferred by the expression of a gene encoding a synthetic peptide in transgenic cotton (*Gossypium hirsutum* L.) plants. *Plant Biotechnology Journal* 3: 545-554.
- Rajasekaran, K., De Lucca, A.J. and Cary, J.W., 2009a. Aflatoxin control through transgenic approaches. *Toxin Reviews* 28: 89-101.
- Rajasekaran, K., Jaynes, J.M. and Cary, J.W., 2009b. Transgenic expression of lytic peptides in food and feed crops to control phytopathogens and preharvest mycotoxin contamination. In: Appell, M., Kendra, D.F. and Trucksess, M.W. (eds.) *Mycotoxin prevention and control in agriculture*. ACS symposium series. Vol. 1031. American Chemical Society, Washington, DC, USA, pp. 119-142.
- Rajasekaran, K., Sayler, R.J., Sickler, C.M., Majumdar, R., Jaynes, J.M. and Cary, J.W., 2018. Control of *Aspergillus flavus* growth and aflatoxin production in transgenic maize kernels expressing a tachyplestin-derived synthetic peptide, AGM182. *Plant Science*: in press.
- Rank, C., Klejnstrup, M.L., Petersen, L.M., Kildgaard, S., Frisvad, J.C., Held Gotfredsen, C. and Ostenfeld Larsen, T., 2012. Comparative chemistry of *Aspergillus oryzae* (RIB40) and *A. flavus* (NRRL 3357). *Metabolites* 2: 39-56.
- Reverberi, M., Punelli, M., Scala, V., Scarpari, M., Uva, P., Mentzen, W.I., Dolezal, A.L., Woloshuk, C., Pinzari, F., Fabbri, A.A., Fanelli, C. and Payne, G.A., 2013. Genotypic and phenotypic versatility of *Aspergillus flavus* during maize exploitation. *PLoS ONE* 8: e68735.
- Reyes-Dominguez, Y., Bok, J.W., Berger, H., Shwab, E.K., Basheer, A., Gallmetzer, A., Scazzocchio, C., Keller, N. and Strauss, J., 2010. Heterochromatic marks are associated with the repression of secondary metabolism clusters in *Aspergillus nidulans*. *Molecular Microbiology* 76: 1376-1386.
- Riley, R.T., Goeger, D.E., Yoo, H. and Showker, J.L., 1992. Comparison of three tetramic acids and their ability to alter membrane function in cultured skeletal muscle cells and sarcoplasmic reticulum vesicles. *Toxicology and Applied Pharmacology* 114: 261-267.
- Rokas, A., Payne, G., Fedorova, N.D., Baker, S.E., Machida, M., Yu, J., Georgianna, D.R., Dean, R.A., Bhatnagar, D., Cleveland, T.E., Wortman, J.R., Maiti, R., Joardar, V., Amedeo, P., Denning, D.W. and Nierman, W.C., 2007. What can comparative genomics tell us about species concepts in the genus *Aspergillus*? *Studies in Mycology* 59: 11-17.
- Roze, L.V., Hong, S.Y. and Linz, J.E., 2013. Aflatoxin biosynthesis: current frontiers. *Annual Review of Food Science and Technology* 4: 293-311.
- Ruhlman, T.A., Rajasekaran, K. and Cary, J.W., 2014. Expression of chloroperoxidase from *Pseudomonas pyrocinia* in tobacco plastids for fungal resistance. *Plant Science* 228: 98-106.
- Sabburg, R., Obanor, F., Aitken, E. and Chakraborty, S., 2015. Changing fitness of a necrotrophic plant pathogen under increasing temperature. *Global Change Biology* 21: 3126-3137.
- Saikia, S., Nicholson, M.J., Young, C., Parker, E.J. and Scott, B., 2008. The genetic basis for indole-diterpene chemical diversity in filamentous fungi. *Mycological Research* 112: 184-199.
- Saikia, S., Parker, E.J., Koulman, A. and Scott, B., 2006. Four gene products are required for the fungal synthesis of the indole-diterpene, paspaline. *FEBS Letters* 580: 1625-1630.
- Sanchis, V. and Magan, N., 2004. Environmental profiles for growth and mycotoxin production. In: Magan, N. and Olsen, M. (eds.) *Mycotoxins in food: detection and control*. Woodhead Publishing Ltd., Cambridge, UK, pp. 174-189.
- Saruwatari, T., Yagishita, F., Mino, T., Noguchi, H., Hotta, K. and Watanabe, K., 2014. Cytochrome P450 as dimerization catalyst in diketopiperazine alkaloid biosynthesis. *ChemBioChem* 15: 656-659.
- Satterlee, T., Cary, J.W. and Calvo, A.M., 2016. RmtA, a putative arginine methyltransferase, regulates secondary metabolism and development in *Aspergillus flavus*. *PLoS ONE* 11: e0155575.
- Schmidt-Heydt, M., Abdel-Hadi, A., Magan, N. and Geisen, R., 2009. Complex regulation of the aflatoxin biosynthesis gene cluster of *Aspergillus flavus* in relation to various combinations of water activity and temperature. *International Journal of Food Microbiology* 135: 231-237.
- Schmidt-Heydt, M., Rufer, C.E., Abdel-Hadi, A., Magan, N. and Geisen, R., 2010. The production of aflatoxin B<sub>1</sub> or G<sub>1</sub> by *Aspergillus parasiticus* at various combinations of temperature and water activity is related to the ratio of aflS to aflR expression. *Mycotoxin Research* 26: 241-246.
- Schubert, M., Houdelet, M., Kogel, K.H., Fischer, R., Schillberg, S. and Nolke, G., 2015. Thanatin confers partial resistance against aflatoxigenic fungi in maize (*Zea mays*). *Transgenic Research* 24: 885-895.
- Sharma, K.K., Pothana, A., Prasad, K., Shah, D., Kaur, J., Bhatnagar, D., Chen, Z.Y., Raruang, Y., Cary, J.W., Rajasekaran, K., Sudini, H.K. and Bhatnagar-Mathur, P., in press. Peanuts that keep aflatoxin at bay: a threshold that matters. *Plant Biotechnology Journal*. <https://doi.org/10.1111/pbi.12846>.
- Smith, C.A., Robertson, D., Yates, B., Nielsen, D.M., Brown, D., Dean, R.A. and Payne, G.A., 2008. The effect of temperature on natural antisense transcript (NAT) expression in *Aspergillus flavus*. *Current Genetics* 54: 241-269.

- Sobolev, V.S., Cole, R.J., Dorner, J.W., Horn, B.W., Harrigan, G.G. and Gloer, J.B., 1997. Isolation and structure elucidation of a new metabolite produced by *Aspergillus parasiticus*. *Journal of Natural Products* 60: 847-850.
- Strauss, J. and Reyes-Dominguez, Y., 2011. Regulation of secondary metabolism by chromatin structure and epigenetic codes. *Fungal Genetics and Biology* 48: 62-69.
- Sultan, Y., 2010. Biodiversity of mycotoxigenic *Aspergillus* species in Egyptian peanuts and strategies for minimizing aflatoxin contamination. PhD-thesis, Cranfield University, Cranfield, UK.
- Sundaresha, S., Manoj Kumar, A., Rohini, S., Math, S.A., Keshamma, E., Chandrasekhar, S.C. and Udayakumar, M., 2010. Enhanced protection against two major fungal pathogens of groundnut, *Cecospora arachidicola* and *Aspergillus flavus* in transgenic groundnut over-expressing a tobacco b-1-3 glucanase. *European Journal of Plant Pathology* 126: 497-508.
- †t Hoen, P.A.C., Ariyurek, Y., Thygesen, H.H., Vreugdenhil, E., Vossen, R.H., De Menezes, R.X., Boer, J.M., Van Ommen, G.J. and Den Dunnen, J.T., 2008. Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms. *Nucleic Acids Research* 36: e141.
- Tang, M.C., Lin, H.C., Li, D., Zou, Y., Li, J., Xu, W., Cacho, R.A., Hillenmeyer, M.E., Garg, N.K. and Tang, Y., 2015. Discovery of unclustered fungal indole diterpene biosynthetic pathways through combinatorial pathway reassembly in engineered yeast. *Journal of the American Chemical Society* 137: 13724-13727.
- TePaske, M.R., Gloer, J.B., Wicklow, D.T. and Dowd, P.F., 1991. Leporin A: an antiinsectan N-alkoxy-pyridone from the sclerotia of *Aspergillus leporis*. *Tetrahedron Letters* 32: 5687-5690.
- TePaske, M.R., Gloer, J.B., Wicklow, D.T. and Dowd, P.F., 1992. Aflavarin and beta-aflatrein: new anti-insectan metabolites from the sclerotia of *Aspergillus flavus*. *Journal of Natural Products* 55: 1080-1086.
- Terabayashi, Y., Sano, M., Yamane, N., Marui, J., Tamano, K., Sagara, J., Dohmoto, M., Oda, K., Ohshima, E., Tachibana, K., Higa, Y., Ohashi, S., Koike, H. and Machida, M., 2010. Identification and characterization of genes responsible for biosynthesis of kojic acid, an industrially important compound from *Aspergillus oryzae*. *Fungal Genetics and Biology* 47: 953-961.
- Thakare, D., Zhang, J., Wing, R.A., Cotty, P.J. and Schmidt, M.A., 2017. Aflatoxin-free transgenic maize using host-induced gene silencing. *Science Advances* 3: 1-8.
- Turner, W.B. and Aldridge, D.C., 1983. *Fungal metabolites*. Vol. II. Academic Press, New York, NY, USA, 631 pp.
- Umemura, M., Nagano, N., Koike, H., Kawano, J., Ishii, T., Miyamura, Y., Kikuchi, M., Tamano, K., Yu, J., Shin-ya, K. and Machida, M., 2014. Characterization of the biosynthetic gene cluster for the ribosomally synthesized cyclic peptide ustiloxin B in *Aspergillus flavus*. *Fungal Genetics and Biology* 68: 23-30.
- Vary, Z., Mullins, E., McElwain, J.C. and Doohan, F.M., 2015. The severity of wheat diseases increases when plants and pathogens are acclimatized to elevated carbon dioxide. *Global Change Biology* 21: 2661-2669.
- Vesth, T.C., Brandl, J. and Andersen, M.R., 2016. FunGeneClusterS: predicting fungal gene clusters from genome and transcriptome data. *Synthetic and Systems Biotechnology* 1: 122-129.
- Waliyar, E., Osiru, M., Ntare, B.R., Vijay Krishna Kumar, K., Sudini, H., Traore, A. and Diarra, B., 2015. Post-harvest management of aflatoxin contamination in groundnuts. *World Mycotoxin Journal* 8: 245-252.
- Wang, H., Lei, Y., Yan, L., Cheng, K., Dai, X., Wan, L., Guo, W., Cheng, L. and Liao, B., 2015. Deep sequencing analysis of transcriptomes in *Aspergillus flavus* in response to resveratrol. *BMC Microbiology* 15: 182.
- Weaver, M.A., Scheffler, B.E., Duke, M., Ballard, L., Abbas, H.K. and Grodowitz, M.J., 2017. Genome sequences of three strains of *Aspergillus flavus* for the biological control of aflatoxin. *Genome Announcement* 5: 44.
- Weber, T. and Kim, H.U., 2016. The secondary metabolite bioinformatics portal: computational tools to facilitate synthetic biology of secondary metabolite production. *Synthetic and Systems Biotechnology* 1: 69-79.
- Weissinger, A., Sampson, K., Urban, L., Ingram, K., Payne, G., Scanlon, S., Liu, Y.-S. and Cleveland, T.E., 2000. Transformation with genes encoding peptidyl MIM<sup>®</sup>, as a means of reducing aflatoxin contamination in peanut. In: USDA-ARS (eds.) Proceedings of the 2000 USDA-ARS aflatoxin elimination workshop. Yosemite, CA, USA, pp. 112-114.
- Weissinger, A., Wu, M., Wang, X., Isleib, T., Stalker, T., Shew, B., Rajasekaran, K., Cary, J. and Cleveland, T.E., 2007. Advancement and testing of transgenic peanuts with enhanced resistance to *A. flavus*. In: USDA-ARS (eds.) Proceedings of the 2007 annual aflatoxin/fumonisin workshop. Atlanta, GA, USA, pp. 106-107.
- Wilkinson, J.R., Kale, S.P., Bhatnagar, D., Yu, J. and Ehrlich, K.C., 2011. Expression profiling of non-aflatoxigenic *Aspergillus parasiticus* mutants obtained by 5-azacytosine treatment or serial mycelial transfer. *Toxins* 3: 932.
- Wilkinson, J.R., Yu, J., Bland, J.M., Nierman, W.C., Bhatnagar, D. and Cleveland, T.E., 2007. Amino acid supplementation reveals differential regulation of aflatoxin biosynthesis in *Aspergillus flavus* NRRL 3357 and *Aspergillus parasiticus* SRRC 143. *Applied Microbiology and Biotechnology* 74: 1308-1319.
- Williams, W.P. and Windham, G.L., 2012. Registration of Mp718 and Mp719 germplasm lines of maize. *Journal of Plant Registrations* 6: 200-202.
- Williams, W.P., Krakowsky, M.D., Scully, B.T., Brown, R.L., Menkir, A., Warburton, M.L. and Windham, G.L., 2015. Identifying and developing maize germplasm with resistance to accumulation of aflatoxins. *World Mycotoxin Journal* 8: 193-209.
- Woloshuk, C.P., Foutz, K.R., Brewer, J.F., Bhatnagar, D., Cleveland, T.E. and Payne, G.A., 1994. Molecular characterization of aflR, a regulatory locus for aflatoxin biosynthesis. *Applied and Environmental Microbiology* 60: 2408-2414.
- Wolt, J.D., Wang, K. and Yang, B., 2016. The Regulatory status of genome-edited crops. *Plant Biotechnology Journal* 14: 510-518.
- World Health Organization (WHO), 2017. Evaluation of certain contaminants in food: eighty-third report of the Joint FAO/WHO Expert Committee on Food Additives. WHO technical report series, No. 1002. WHO Press, Geneva, Switzerland, 167 pp.

- Wu, F., Bhatnagar, D., Bui-Klimke, T., Carbone, I., Hellmich, R., Munkvold, G., Paul, P., Payne, G. and Takle, E., 2011. Climate change impacts on mycotoxin risks in US maize. *World Mycotoxin Journal* 4: 79-93.
- Young, M.D., Wakefield, M.J., Smyth, G.K. and Oshlack, A., 2010. Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biology* 11: R14.
- Yu, J., 2012. Current understanding on aflatoxin biosynthesis and future perspective in reducing aflatoxin contamination. *Toxins* 4: 1024-1057.
- Yu, J., Chang, P.K., Cary, J.W., Wright, M., Bhatnagar, D., Cleveland, T.E., Payne, G.A. and Linz, J.E., 1995. Comparative mapping of aflatoxin pathway gene clusters in *Aspergillus parasiticus* and *Aspergillus flavus*. *Applied and Environmental Microbiology* 61: 2365-2371.
- Yu, J., Chang, P.-K., Ehrlich, K.C., Cary, J.W., Bhatnagar, D., Cleveland, T.E., Payne, G.A., Linz, J.E., Woloshuk, C.P. and Bennett, J.W., 2004. Clustered pathway genes in aflatoxin biosynthesis. *Applied and Environmental Microbiology* 70: 1253-1262.
- Yu, J., Cleveland, T.E., Wilkinson, J.R., Campbell, B.C., Kim, J.H., Kim, H.S., Bhatnagar, D., Payne, G.A. and Nierman, W.C., 2006. *Aspergillus flavus* expressed sequence tags and microarray as tools in understanding aflatoxin biosynthesis. *Mycotoxin Research* 22: 16-21.
- Yu, J., Fedorova, N.D., Montalbano, B.G., Bhatnagar, D., Cleveland, T.E., Bennett, J.W. and Nierman, W.C., 2011. Tight control of mycotoxin biosynthesis gene expression in *Aspergillus flavus* by temperature as revealed by RNA-Seq. *FEMS Microbiology Letters* 322: 145-149.
- Yu, J., Ronning, C.M., Wilkinson, J.R., Campbell, B.C., Payne, G.A., Bhatnagar, D., Cleveland, T.E. and Nierman, W.C., 2007. Gene profiling for studying the mechanism of aflatoxin biosynthesis in *Aspergillus flavus* and *A. parasiticus*. *Food Additives and Contaminants* 24: 1035-1042.
- Yu, J.H., Butchko, R.A., Fernandes, M., Keller, N.P., Leonard, T.J. and Adams, T.H., 1996. Conservation of structure and function of the aflatoxin regulatory gene aflR from *Aspergillus nidulans* and *A. flavus*. *Current Genetics* 29: 549-555.
- Yu, L.X. and Setter, T.L., 2003. Comparative transcriptional profiling of placenta and endosperm in developing maize kernels in response to water deficit. *Plant Physiology* 131: 568-582.
- Zablotowicz, R.M., Abbas, H.K. and Locke, M.A., 2007. Population ecology of *Aspergillus flavus* associated with Mississippi Delta soils. *Food Additives and Contaminants* 24: 1102-1108.
- Zhang, F., Guo, Z., Zhong, H., Wang, S., Yang, W., Liu, Y. and Wang, S., 2014. RNA-Seq-based transcriptome analysis of aflatoxigenic *Aspergillus flavus* in response to water activity. *Toxins* 6: 3187-3207.
- Zhang, K., Raboanatahiry, N., Zhu, B. and Li, M., 2017. Progress in genome editing technology and its application in plants. *Frontiers in Plant Science* 8: 177.
- Zhang, S., Monahan, B.J., Tkacz, J.S. and Scott, B., 2004. Indole-diterpene gene cluster from *Aspergillus flavus*. *Applied and Environmental Microbiology* 70: 6875-6883.
- Zhao, S., Fung-Leung, W.P., Bittner, A., Ngo, K. and Liu, X., 2014. Comparison of RNA-Seq and microarray in transcriptome profiling of activated T cells. *PLoS ONE* 9: e78644.
- Zhuang, Z., Lohmar, J.M., Satterlee, T., Cary, J.W. and Calvo, A.M., 2016. The master transcription factor mtfa governs aflatoxin production, morphological development and pathogenicity in the fungus *Aspergillus flavus*. *Toxins* 8: 1



# Advances in molecular and genomic research to safeguard food and feed supply from aflatoxin contamination

Bhatnagar, D

2018-02-23

Attribution-NonCommercial-ShareAlike 4.0 International

---

Bhatnagar D, Rajasekaran K, Gilbert M, Cary JW, Magan N, Advances in molecular and genomic research to safeguard food and feed supply from aflatoxin contamination, Vol. 11, Issue 1, 2018, pp. 47-72

<http://dx.doi.org/10.3920/WMJ2017.2283>

*Downloaded from CERES Research Repository, Cranfield University*