

## **A loss-of-function allele of a *TAC1*-like gene (*SITAC1*) located on tomato chromosome 10 is a candidate for the *Erectoid leaf* (*Erl*) mutation**

Matías González-Arcos<sup>1\*</sup>; Maria Esther de Noronha Fonseca<sup>2</sup> Daniel Basílio Zandonadi<sup>3</sup>; Lázaro E. P. Peres<sup>4</sup>; Ana Arruabarrena<sup>1</sup>; Demetrys S. Ferreira<sup>5</sup>; Zoltan Kevei<sup>5</sup>; Fady Mohareb<sup>5</sup>; Andrew J. Thompson<sup>5</sup> & Leonardo S. Boiteux<sup>2</sup>

<sup>1</sup>National Research Program on Horticultural Production, National Institute of Agricultural Research (INIA), Estación Experimental INIA Salto Grande, CP 50000, Salto, Uruguay.

<sup>2</sup>Nacional Center for Vegetable Crops Research (CNPH) – Embrapa Vegetable Crops (Hortaliças), Brasília–DF, Brazil.

<sup>3</sup>Universidade Federal do Rio de Janeiro (UFRJ), Nupem, 27965045, Macaé–RJ, Brazil.

<sup>4</sup>Laboratory of Hormonal Control of Plant Development, Departamento de Ciências Biológicas, Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo (ESALQ/USP), Piracicaba–SP, Brazil.

<sup>5</sup>Cranfield Soil and Agrifood Institute, Cranfield University, Cranfield, MK43 0AL, UK.

\*Corresponding author (matgon@inia.org.uy)

**Acknowledgements** This work was done in the context of MG-A doctoral studies program at the Facultad de Agronomía, Universidad de la República Oriental del Uruguay. We thank A. Manzoni, I. Laxague and N. Zunini of INIA Salto Grande, and W. P. Dutra and A. F. Costa of Embrapa Vegetable Crops, for their assistance in conducting some of the experiments. LSB and MENF were supported by CNPq and CAPES grants. AJT and FM were supported by BBSRC Research Grant BB/L011611/1.

## SUMMARY

The genetic basis of an erectoid leaf phenotype was investigated in distinct tomato breeding populations, including one derived from *Solanum lycopersicum* ‘LT05’ (with the erectoid leaf phenotype and uniform ripening, genotype *uu*) × *S. pimpinelifolium* ‘TO-937’ (with the wild-type leaf phenotype and green fruit shoulder, genotype *UU*). The erectoid leaf phenotype was inherited as a semi-dominant trait and it co-segregated with the *u* allele of gene *SLGLK2* (*Solyc10g008160*). This genomic location coincides with a previously described semi-dominant mutation named as *Erectoid leaf* (*Erl*). The genomes of ‘LT05’, ‘TO-937’, and three other unrelated accessions (with the wild-type *Erl*<sup>+</sup> allele) were resequenced with the aim of identifying candidate genes. Comparative genomic analyses, including the reference genome ‘Heinz 1706’ (*Erl*<sup>+</sup> allele), identified an *Erectoid leaf*-specific single nucleotide polymorphism (SNP) in the gene *Solyc10g009320*. This SNP caused a change of a glutamine (CAA) codon (present in all the wild-type genomes) to a TAA (= ochre stop-codon) in the *Erl* allele, resulting in a smaller version of the predicted mutant protein (221 versus 279 amino acids). *Solyc10g009320*, previously annotated as an ‘unknown protein’, was identified as a *TILLER ANGLE CONTROL1* (*TAC1*)-like gene. Linkage between the *Erl* and *Solyc10g009320* was confirmed via Sanger sequencing of the PCR amplicons of the two variant alleles. No recombinants were detected in 265 F<sub>2</sub> individuals. Contrasting S<sub>7</sub> near-isogenic lines were also homozygous for each of the alternate alleles, reinforcing *Solyc10g009320* as a strong *Erl* candidate gene and opening the possibility for fine-tuning manipulation of tomato architecture in breeding programs.

**Key words:** *Solanum lycopersicum*; resequencing; comparative genomic analysis; plant architecture; breeding.

## INTRODUCTION

Genetic factors are the major determinants of plant architecture, even though the spatial structure of a plant might also be influenced by various environmental stimuli such as light, temperature, humidity, mineral and organic nutrition (Wang et al. 2018). For this reason, breeding programs have placed great emphasis in exploiting the genetic diversity associated with plant architecture as a strategy to develop high-yielding cultivars with greater adaptation to wide array of environmental conditions and cropping systems (Coyne 1980; Huyghe 1998; Jiao et al. 2010).

In tomato (*Solanum lycopersicum* L.), plant growth habit (i.e. determinate, semi-determinate, and indeterminate), foliar insertion angle, leaf size and internode length are important plant architectural traits for breeding, especially because of their impact on the vertical distribution of light through the crop canopy and the consequent effects on the efficiency of light interception (Sarlikioti et al. 2011a, 2011b; Silva et al. 2018). Currently, tomato breeding for indeterminate growth habit is focused on reducing the length of internodes to increase the number of trusses per stem length (Zsögön et al. 2017). However, this can increase self-shading and reduce the efficiency of light absorption (Sarlikioti et al. 2011b). At a given light intensity, the leaf insertion angle has direct implications on the amount of light received per leaf surface unit (Ehleringer & Werk 1986; Ezcurra et al. 1991) whereby a smaller leaf insertion angle (i.e. more erect leaves) can lead to more uniform light intensities vertically through the canopy, reducing light stress at the upper levels and increasing photosynthesis in the lower levels. More erect leaves may also improve the efficacy of contact pesticide applications by improving penetration through the crop canopy and reaching more efficiently abaxial leaf surfaces.

Although the control of lateral growth has been extensively studied at the molecular level in some species, the genetic and physiological mechanisms that define the insertion angle of distinct organs (including leaves) have not been properly characterized in many taxa, including Solanaceae (for general review see Wang and Li 2008; Teichmann and Muhr 2015; Roychoudhry and Kepinski 2015). So far, all physiological models converge to the central role of auxin content and distribution in the response to the continuous growth of shoots and roots (Friml et al. 2003; Roychoudhry and Kepinski 2015). Polar transport mediated by PIN and AUX/LAX proteins is a major mechanism that regulates auxin distribution in plants. These gene products control cellular auxin efflux and influx, respectively, through their subcellular localization at the plasma membrane (Wiśniewska et al. 2006; Vanneste and Friml 2009).

In tomato, characterization of genes that control auxin fluxes and directional growth of organs is still quite limited. Pattison and Catalá (2012) studied the function of the genes in tomato homologous to *PIN* and *AUX/LAX* and they verified that *SIPIN4* and *SIPIN3* have specific roles in the regulation of vegetative shoot architecture. Recently, Shi et al. (2017) proposed a model to explain how the auxin polar transport mediated by *PIN1* is critical in tomato leaf polarity formation. In other plant species, a distinct set of genes associated with plant architecture has been identified. In monocotyledons, *TILLER ANGLE CONTROL1 (TAC1)* (Yu et al. 2007) and *LAZY1*, a gravitropism-related gene (Li et al. 2007), are the main genetic factors identified as being involved in the regulation of shoot angle in rice (*Oryza sativa* L.). In the case of *TAC1*, a mutation that reduces gene expression is responsible for a lower insertion angle of the lateral shoots (i.e. more erect). The *LAZY1* gene has sequence motifs similar to *TAC1* with the addition of an Ethylene-responsive element binding factor-associated Amphiphilic Repression (EAR) domain. *LAZY1* loss-of-function mutants are associated with larger insertion angle of lateral shoots. It has been demonstrated that the recessive *lazy1* mutant increases the polar (apical–basal) auxin transport and decreases lateral transport, generating an abnormal auxin flow/distribution. The associated phenotype is due to the loss of gravitropism, leading to larger leaf insertion angles (Roychoudhry and Kepinski, 2015). In peach [*Prunus persicae* (L.) Batsch], the semi-dominant allele “*broomy*” (*br* – which was later designated as “*pillar*”) is responsible for a more vertical growth of the branches (Scorza et al. 1989, 2002). Subsequent genetic/genomic characterization determined that the recessive *br* allele is a non-functional mutation of a homologue of the monocotyledonous gene *TAC1*, which they denominated *PpeTAC1* (Dardick et al. 2013). Plants with the homozygous *br* allele showed higher auxin concentration in shoots when compared with wild-type plants with a horizontal branch pattern (Tworkoski et al. 2006). The auxin content in peach was found to be inversely proportional to the expression of the *TAC1* gene (Tworkoski et al. 2015). The silencing of *PpeTAC1* orthologue gene in *Arabidopsis thaliana* generated a “more erect plant”, suggesting that this class of genes works universally in promoting vertical growth. *TAC1* and *LAZY1* belong to a superfamily of genes defined by an IGT (GφL(A/T)GT) domain, which is present in a wide array of plant genomes and is related to the vertical growth of the shoots through the regulation of auxin polar transport as demonstrated in rice, maize, *A. thaliana*, and peach (Dardick et al 2013, Roychoudhry and Kepinski 2015). The understanding of the genetic control of shoot architecture could provide breeding tools for selection of cultivars with improved

utilization of light, more adapted to high planting densities (Testa et al., 2016), and that also increases the efficacy of pesticide applications.

We have observed in some segregating tomato breeding lines a peculiar erect leaf phenotype, which apparently affects all aerial organs, especially young shoots. A highly endogamic breeding line (named ‘LT05’) was recovered and it showed a genetically stable erect leaf phenotype. Two apparently similar phenotypes have already been described in tomato: a radiation-induced recessive mutant *erecta* (*er*) (Tomato Genetics Resource Center: <http://tgrc.ucdavis.edu/>, last accessed 12 March, 2018), present in the accession *S. lycopersicum* LA600 (derived from the ‘Codine Red’ cultivar) and a spontaneous semi-dominant mutation named *Erectoid leaf* (*Erl*) (Georgiev and Kraptchev 1992; Tomato Genetics Resource Center: <http://tgrc.ucdavis.edu/>, last accessed 12 March, 2018). Additional observations indicated that the *Erl* mutation co-segregated with the *uniform ripening* (*u*) mutation (Georgiev and Kraptchev 1992). The *u* mutation was identified as a loss-of-function allele of the *GOLDEN2-LIKE* (*GLK2*) gene located on tomato chromosome 10 (Kinzer et al. 1990; Powell et al. 2012). Here, we investigate the genetic basis and chromosomal location of the erect leaf phenotype observed in the inbred line ‘LT05’. Based on a further genomic analysis, we report a strong candidate gene at the *Erl* locus containing a loss-of-function mutation. The identification of genetic variability associated with leaf angle in tomato opens the possibility for fine-tuning manipulation of plant architecture in breeding programs.

## MATERIALS AND METHODS

### Accessions employed as parental lines and development of segregating populations

The genetic basis of the erectoid leaf mutation was investigated using three distinct segregating F<sub>2</sub> populations. The first segregating F<sub>2</sub> population was generated from a cross between the *S. lycopersicum* inbred line ‘LT05’ (with the erectoid leaf phenotype and uniform ripening, genotype *uu*) and *S. lycopersicum* ‘LT17’ (an inbred line with the wild-type horizontal leaf phenotype). Analyses were conducted with the contrasting parental lines (13 plants each) and five crossing generations: F<sub>1</sub> (n=13), reciprocal F<sub>1</sub>’ (n=13); backcross (BC) to ‘LT05’ (n=32), BC to ‘LT17’ (n=37), and F<sub>2</sub> (n=138). The second segregating F<sub>2</sub> population (n=274) was obtained from the cross ‘LT05’ × *S. pimpinelifolium* ‘TO-937’ (with green fruit shoulder, genotype *UU*) (Powell et al. 2012). A third F<sub>2</sub> population was produced by first generating a pair of near-isogenic lines (NILs) and then crossing them. These NILs were created as follows: from the *S. lycopersicum*

‘LT05’ × *S. lycopersicum* ‘LT17’ cross, three putative heterozygous F<sub>2</sub> plants (with intermediate leaf phenotype) were visually selected and selfed to generate three segregating F<sub>3</sub> families of 20 plants each. Individual F<sub>3</sub> plants with intermediate leaf phenotype (i.e. putative heterozygous) were then chosen to continue a consecutive progeny testing-based process of selection and subsequent selfing. This process was repeated until obtaining segregating F<sub>5</sub> families. In this step, three individual F<sub>5</sub>:F<sub>6</sub> plants with erect leaf phenotype and three with normal leaf phenotype were visually selected within the same segregating progeny and then selfed. Single F<sub>6</sub> plants able to generate progenies with stability for each of the opposing traits (i.e. the erect leaf versus normal leaf) were chosen as the contrasting near-isogenic lines and named as ‘IsoL-EL’ (with stable erectoid leaf phenotype) and ‘IsoL-WTL’ (with stable wild-type leaf phenotype). The NILs were then crossed (‘IsoL-EL’ × ‘IsoL-WTL’) and the F<sub>1</sub> plants were selfed to generate an F<sub>2</sub> population (IsoL-F<sub>2</sub>) composed by 127 plants (see the representation of the process in **Online resource 1**). The IsoL-F<sub>2</sub> population was, therefore, segregating mainly for the erect/normal leaf phenotype, whereas outside this locus the genetic background was predominantly a homozygous non-segregating mosaic of the genomes of the two parental lines ‘LT05’ and ‘LT17’. The IsoL-F<sub>2</sub> population was also used in candidate gene validation analyses (see section below).

### **Evaluation of the leaf growth pattern (erectoid versus wild-type)**

All evaluated plants were cultivated under greenhouse conditions in 5L pots filled with a mixture of soil and commercial peat. Individual plants were pruned to a single main stem. For inheritance studies, the leaf insertion angle ( $\alpha$ ) between the leaf petiole and the main stem (**Online Resource 2**) was measured in fully developed leaves and employed as a phenotypic indicator of each individual plant. In the case of the F<sub>2</sub> population derived from the cross *S. lycopersicum* ‘LT05’ × *S. lycopersicum* ‘LT17’, two measurements were made at 80 days after sowing. One measurement was done in the lower leaf (immediately below the first floral truss) and other in the first fully developed leaf (counting from the apex). For analyses, both measures were averaged to generate a mean  $\alpha$  angle value that was converted to an ordinal scale according to the following criteria: mean  $\alpha$  angle < 100° = erectoid leaf, mean  $\alpha$  angle between 100-125° = intermediate, mean  $\alpha$  angle > 125° = wild-type with standard leaf phenotype. For all segregating F<sub>2</sub> populations used in trait chromosomal location and candidate gene validation studies, the classification of the leaf growth trait was done by directly assessing the general aspect of the plants under

greenhouse conditions. The plants were classified as either erect leaf or non-erect leaf. This last category involved intermediate as well as wild-type leaf phenotypes.

### **Chromosome mapping of the erectoid leaf trait**

The F<sub>2</sub> population from the cross *S. lycopersicum* ‘LT05’ × *S. pimpinelifolium* ‘TO-937’ was employed to verify the linkage of the erectoid leaf phenotype with the uniform ripening *SlGLK2* gene (*u*, *Solyc10g008160*) (Powell et al. 2012). The green fruit shoulder phenotype (presence of the *U* allele) or its absence (due to the homozygous presence of the *u* allele) was used as a phenotypic marker to evaluate co-segregation with the erectoid leaf trait observed in our populations. Evaluation was carried out visually employing a simple scale of presence/absence for both traits.

### **Genetic and statistical analyses**

For leaf insertion angle ( $\alpha$ ), the standard error and ANOVA were calculated using the software InfoStat version 2014 (Di Rienzo et al. 2013). For ANOVA, significant differences were claimed for  $P < 0.01$  in a Tukey and Dunn’s post-hoc test. A chi-squared test was applied to: (a) verify across the F<sub>2</sub> populations the goodness-of-fit of the erectoid vs. wild-type segregation ratios to Mendelian segregation models, and (b) to confirm the linkage between the erectoid leaf trait and the phenotypic marker green fruit shoulder, searching for a statistical difference from a 3:1 segregation ratio (i.e. independence) in a sub-group of erectoid and wild-type F<sub>2</sub> genotypes. In all cases a probability level ( $P$ -value) is given as the value for the null hypothesis.

### **Comparative genomic analyses of the chromosome 10 and variant screening**

Resequencing information was obtained using genomic DNA from the erectoid leaf line ‘LT05’ and from four genetically diverse wild-type leaf accessions: ‘TO-937’ (Powell et al. 2012), ‘CNPH498’ (data not shown), ‘Santa Clara’ (Carmo et al., 2017), and ‘Viradoro’ (Giordano et al. 2000). Genomic DNA was extracted from leaf tissue of these accessions using DNeasy PowerPlant Pro Kit (QIAGEN Hilden, Germany). Whole genome sequencing of individual samples was sequenced on one lane of a HiSeq 2500 (Illumina Inc., San Diego, CA), at the Centro de Biotecnologia Animal (ESALQ/USP). Sequencing was performed with 100 bp paired-end reads. Quality control was done using

FastQC (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Sequencing data from each of the five accessions was assembled separately against the tomato reference genome (version SL2.50) using SeqMan NGen version 14 software with default parameters (Lasergene, DNASTAR, Madison, WI, USA). An in-line Bayesian modeled variant detector based on the MAQ caller (Li, 2008) was used to tabulate SNPs and small indels relative to the reference genome in each accession. Variant calls from the five assemblies were then combined for further analysis in ArrayStar 14 (Lasergene Suite 14). Based on the chromosome mapping studies of the erect leaf trait, we carried out a search for gene variants only within chromosome 10 using variant calls from the five accessions. Since we also found the reference genome, ‘Heinz 1706’ (Tomato Genome Consortium, 2012), which have a standard, wild-type (non-erect) leaf pattern, we focused on identifying non-synonymous variants that were exclusively found in the erect leaf line ‘LT05’. To do so, the following series of filtering criteria was used in ArrayStar: (1) non-synonymous variants occurring in ‘LT05’ but none of four non-erect leaf accessions (and by definition, not in the reference sequence); (2) variant positions with a minimum depth of coverage of 10; (3) SNP% of 100 and (4) Qcall  $\geq 7$ . Summary variant information was exported for individual accessions and then imported into Microsoft Excel (Office 2016; Microsoft Corp., Redmond, WA) for further filtering and analyses. These final variants were manually verified by inspection of the corresponding sequence assemblies to eliminate artifacts arising from assembly differences. Finally, the best candidate genes were chosen considering: the physical/genetic position respect to *SlGLK2*, the protein annotation (ITAG 2.40; <http://solgenomics.net/>, last accessed 18 April, 2018), the predicted effects of amino acid substitutions on protein function using the PROVEAN tool (Choi et al. 2012) (<http://provean.jcvi.org/index.php>, last accessed 18 April, 2018) and the expression pattern according to TomExpress RNAseq database (Zouine et al. 2017) (<http://gbf.toulouse.inra.fr/tomexpress/>, last accessed 12 March, 2018).

### **Multiple alignments of the predicted protein sequences from *TAC1*-like genes of tomato and other plant species**

An erectoid leaf locus-specific non-synonymous, single nucleotide polymorphism (SNP) was found in the putative tomato *TAC1*-like gene located on the chromosome 10 (XP\_004248091 = *Solyc10g009320*). For this reason, multiple alignments of the predicted TAC1-like amino acid sequences from the wild-type and erectoid leaf tomato lines as well as from different plant species were carried out. TAC1-like amino acid



sequences of peach (XP\_020413395), soybean (KRH06413), *A. thaliana* (OAP08981.1), maize (NP\_001170644), and rice (BAF25656.2) were retrieved from GenBank database and aligned employing the Muscle algorithm with the default parameters on Geneious software v7.1.5.

### **Validation of *Solyc10g009320* as a candidate gene of the erectoid leaf phenotype**

Individuals of two F<sub>2</sub> populations were used: the 127 individuals of the IsolF<sub>2</sub> and 71 erect and 67 non-erect individuals of the ('LT05' × 'TO-937') F<sub>2</sub>, comprising a total of 265 F<sub>2</sub> individual plants. Genomic DNA was extracted from leaf samples collected at the apex of each individual plants according to a modified methodology employing 2X CTAB and organic solvents (Boiteux et al. 1999). From the sequence variants of *Solyc10g009320*, a primer pair TAC-Tom (F/R) flanking the identified SNP site was designed using PrimerSelect (Lasergene, DNASTAR, Madison, WI, USA). PCR with this primer pair amplified a 400 bp DNA fragment encompassing the SNP site detected within allelic variants of *Solyc10g009320*. Primer sequences for TAC-Tom (F/R) were: 5'-GAG-TTC-AGT-AAG-TGG-TCA-AGA-3' / 5'-AAA-GAA-AGG-ATC-ACT-CTA-GCA-GA-3'. PCR reagents were adjusted to a final volume of 12.5 µL using 5.95 µL Milli-Q water, 1.25 µL of 10X Buffer *Taq* polymerase (100 mM Tris- HCl, pH 8.3 and 500 mM KCl), 0.6 µL of MgCl<sub>2</sub> (50 mM); 0.5 µL of dNTPs (2.5 mM); 1 µL of each primer, 0.1 U/µl *Taq* DNA polymerase (Invitrogen, Gaithersburg, MD, USA), and 2 µL of DNA template (30 ng µL<sup>-1</sup>). The amplification conditions were an initial stage of denaturation at 94 °C for 2 minutes, followed by 30 cycles of: denaturation at 94 °C for 30 seconds, annealing at 60 °C for 1 minute and extension at 72 °C for 1.5 minutes; ending with an extension step at 68 °C for 10 minutes. After purification of PCR products using Wizard kit (Promega, Madison, WI, USA), the amplicons were subjected to Sanger sequencing using an ABI Prism 3130 sequencer of the Genomic Analysis Laboratory (Embrapa Vegetable Crops, Brasília–DF, Brazil) employing the ABI Prism BigDye version 3.1 Kit (Applied Biosystems Division, Foster City, CA, USA) and the primer pair TAC-Tom (F/R). Sequence quality analysis, the removal of low quality fragments, and the identification of consensus sequences were performed using the SeqMan program (Lasergene, Madison, WI, USA). Alignment of multiple protein sequences was performed using the Clustal W method of the MegAlign software package (Lasergene, Madison, WI, USA).

## RESULTS

### Phenotypic characterization and genetic basis of the erectoid leaf growth trait

The most remarkable effect of erectoid leaf phenotype present in the inbred line ‘LT05’ (our original source of this mutation) consisted of a significantly more vertical growth of leaves, leaflets, and lateral shoots when compared to the control wild-type phenotype ‘LT17’ (**Fig. 1**). This trait was also easily identifiable by visual analysis in all segregating F<sub>2</sub> populations. In fact, this mutation expressed its phenotypic effects on all aerial structures (i.e. leaflets, leaves, shoots, and trusses) with more striking manifestation in young shoots. However, due to a former description in the literature limited to the leaf phenotype (Georgiev and Kraptchev 1992; Tomato Genetics Resource Center: <http://tgrc.ucdavis.edu/>), we decided to keep the nomenclature of this mutation as “*erectoid leaf*”. The analysis of the phenotypic results of the cross ‘LT05’ × ‘LT17’ population showed a clear contrast between the parental lines ( $P < 0.0001$ ). The mean insertion angle  $\alpha$  was  $77^{\circ} \pm 2^{\circ}$  for ‘LT05’ (n=13) and  $142^{\circ} \pm 1^{\circ}$  for ‘LT17’ (n=13). The reciprocal hybrids (obtained using each parent as either male or female) displayed similar results. The reciprocal hybrids displayed an intermediate leaf insertion angle between both contrasting lines ( $p < 0.0001$ ). The average insertion angle  $\alpha$  was  $120^{\circ} \pm 1^{\circ}$  and  $119^{\circ} \pm 1^{\circ}$  for F<sub>1</sub> (n=13) and F<sub>1</sub>’ (n=13), respectively (**Fig. 2 a**). These results suggested the absence of maternal inheritance effects. For the segregating families the distribution of their individuals in three different categories of leaf insertion angle (erectoid, intermediate, and wild-type) was analyzed (**Fig. 2 b**). The backcross family ‘LT05’ × F<sub>1</sub> segregated very close to a 1:1 (intermediate : erectoid) ratio. The same was observed with the backcross family ‘LT17’ × F<sub>1</sub> with a 1:1 (intermediate : wild-type) ratio. The F<sub>2</sub> family segregated very closely to the expected 1:2:1 (erectoid : intermediate : wild-type) ratio. Finally, the F<sub>3</sub> families derived from the self-pollinating of F<sub>2</sub> plants with wild-type leaf insertion (**Fig. 2 c right**) displayed 100% individuals with wild-type leaf insertion, while in F<sub>3</sub> families derived from F<sub>2</sub> individuals with erectoid leaf insertion (**Fig. 2 c left**) had 100% of individuals with erectoid leaf insertion. These segregation patterns indicate a strong fit for a single gene model with semi-dominant inheritance of the erectoid leaf phenotype.

### Chromosome location of the gene controlling the erectoid leaf phenotype

Each plant of the F<sub>2</sub> family derived from the cross between ‘LT05’ and ‘TO-937’ was simultaneously evaluated for the erectoid leaf and the uniform fruit ripening phenotype.

A clear-cut 1:3 segregation (monogenic) was observed for both erectoid leaf growth (75:199) and uniform fruit ripening (68:206) (**Table 1**). Within the subgroup of “erectoid” leaf growth (n=75) and the subgroup of “intermediate or wild-type” leaf growth (n=199), it was possible to observe a strong deviation of the expected ratio 3:1 for green fruit shoulder (dominant) and uniform fruit ripening (recessive), indicating that the two loci were not segregating independently, thus confirming their co-location on the tomato chromosome 10. A frequency of 12% of recombinants (33/274) indicated a relatively close genetic distance between these two loci.

### **Comparative analyses of chromosome 10 in the genomes of contrasting (erectoid vs. wild-type) lines**

To identify potential polymorphisms associated with the erectoid leaf trait, a strategy based upon the comparative genome analyses of the original erectoid leaf source (= *S. lycopersicum* ‘LT05’) with the genomes of genetically unrelated accessions displaying standard (wild-type) leaf angle phenotypes (viz. ‘Viradoro’, ‘TO-937’, ‘Santa Clara’, ‘CNPH498’ and ‘Heinz 1706’) was employed. A total of 27,259 variants were found only on chromosome 10. After filtering for variants exclusively present in ‘LT05’, a total of 1,702 variants were obtained (**Fig. 3 a**). When filtering those variants for non-synonymous variants only five SNPs remained (**Fig. 3 b, Table 2**). One of these variants (corresponding to the alkaloid biosynthesis gene *Solyc10g086620*) was discarded because it was located far from the *SlGLK2* gene (*Solyc10g008160* located at position 2,293,088 in SL2.50), which was not in agreement with our data of 12% recombination (**Table 1**). The PROVEAN tool predicted that from the four remaining gene variants only one (*Solyc10g009320*) contained an amino acid change capable of inducing deleterious effect: an SNP that creates a new stop codon (**Fig. 3 c and d**). This *Erectoid leaf*-specific single nucleotide polymorphism (SNP) was found in the position 3,394,715 of chromosome 10, where a DNA substitution of a cytosine (C) for a thymine (T) (C>T) was observed. This SNP resulted in a change in the CAA codon (=coding for a glutamine) of the wild-type genomes to a TAA (= ochre stop-codon) in the genome of the line ‘LT05’ (with erectoid leaf trait). *Solyc10g009320* is about 1.1 Mbp from *SlGLK2*, in the euchromatin, consistent with the observed recombination frequency of 12%. The TomExpress RNASeq database reported a similar expression pattern in seeds, meristems, stem, leaves, and fruits for *Solyc10g009320*, but with higher expression in flowers and lower in roots (**Online**

**Resource 3).** Thus, *Solyc10g009320* became our single candidate gene for controlling the erectoid leaf growth phenotype.

### **Sequence analysis of *Solyc10g009320* and impact on the predicted proteins**

The gene *Solyc10g009320* was found to encode a predicted protein of 279 amino acids. When translated, the *Erl*-specific SNP caused a change of a glutamine (CAA) codon (present in all the wild-type genomes) to a TAA (= ochre stop-codon) at amino acid position 222 in exon 3 (Q222\*) in the ‘LT05’ genome (with erectoid leaf trait), resulting in a smaller version of the predicted mutant protein (221 amino acids) (**Fig. 3 c and d, Online Resource 4**). BLAST analysis (tBLASTx tool) of the predicted protein encoded by *Solyc10g009320* gene indicated 36% amino acid identity with the gene *PpeTAC1*, belonging to the gene family IGT, described to have a major role in determining plant architecture in peach.

### **Multiple alignments of the predicted protein sequences from *TAC1*-like genes of tomato and other plant species**

Multiple alignments of *TAC1*-like proteins from tomato and from a diverse group of plant species revealed the presence of the conserved domains described by Dardick et al. (2013), and specifically the presence of an IGT (GφL(A/T)GT) domain (**Fig. 4**), which is present in a wide array of plant genomes and is related to the vertical growth of the shoots (Dardick et al. 2013, Roychoudhry and Kepinski 2015). Due to the similarities between *TAC1* and *Solyc10g009320* we named this tomato ortholog as *S. lycopersicum TAC1* (*SITAC1*) gene.

### **Sanger sequencing validation of the *SITAC1*-derived marker**

PCR products obtained with the TAC-Tom (F/R) primers (designed to flank the identified SNP causing an early stop codon on *Solyc10g009320*) were Sanger sequenced in individual samples of two F<sub>2</sub> populations and the two contrasting near-isogenic lines. A total of 265 F<sub>2</sub> individuals were sampled from two F<sub>2</sub> populations. Sequence analyses of these individuals indicated the constant presence of the homozygous C>T substitution in all 108 F<sub>2</sub> plants with the erectoid leaf growth phenotype, in the parents, ‘LT05’ and in the near isogenic line IsoL-EL. On the other hand, the C>T substitution was found to be either absent or in heterozygous condition in all 157 F<sub>2</sub> with either intermediate or wild-

type growth phenotype, in the parental lines ‘TO-937’ and IsoL-WTL (**Online Resource 5**).

## DISCUSSION

### **The erectoid leaf phenotype is controlled by a semi-dominant locus on chromosome 10**

Our results indicated that the erectoid leaf phenotype observed in the line ‘LT05’ is controlled by a single semi-dominant gene/locus at the top of chromosome 10. This information was experimentally confirmed by the phenotypic analyses across distinct segregating populations and by linkage analysis with the uniform fruit ripening-coding *SlGLK2* gene, which is located on chromosome 10 (Kinzer et al. 1990; Powell et al. 2012). Two mutations with similar phenotypes were previously described in tomatoes: the *Erectoid leaf* (*Erl*) (Georgiev and Kraptchev 1992) and *erecta* (*er*) (LA0600). Given the similar characteristics in terms of location and pattern of inheritance, we assume that the gene that determines the erectoid leaf phenotype in the line ‘LT05’ is either the same gene or an allelic variant of the wild-type (*Erl*<sup>+</sup>) gene as previously described by Georgiev and Kraptchev (1992) (Tomato Genetics Resource Center: <http://tgrc.ucdavis.edu/>, last accessed 12 March, 2018). Our results indicated that the *Erl*<sup>+</sup> allele is associated with the wild-type phenotype, whereas the *Erl* allele is associated with the erectoid leaf trait. Heterozygous (*Erl*<sup>+</sup>/*Erl*) plants displayed an intermediate leaf growth phenotype. Genes with similar phenotypic expression have been reported in other dicot and monocot plant species. In rice, the recessive loss-of-function allele of the *OzTAC1* gene (Yu et al. 2007) is one of the determinants of erect tiller growth and it has been used, along with other allelic variants, in several modern cultivars in order to generate more efficient crops (Dong et al., 2016). In peach, the “*broomy*” (*br*) allele of *PpeTAC1* defines the plant architecture by modifying the angle of insertion of the lateral branches and also displays a semi-dominant inheritance (Scorza et al. 1989, 2002). In this case, the possibility of manipulating the degree of branch inclination with distinct doses of the allele *br* (homozygous or heterozygous form) was suggested (Tworkoski and Scorza 2001).

### **A *TAC1*-like gene is the best candidate related to the erectoid leaf phenotype**

Our genomic and genetic analyses allowed us to indicate *Solyc10g009320* (previously annotated as an ‘unknown protein’) as being the best candidate gene related to erectoid leaf growth observed in the ‘LT05’ line (zero recombinants in 265 F<sub>2</sub> plants analyzed,

**Online Resource 5).** The methodology that allowed to identify *Solyc10g009320* as the more likely candidate gene was based on the genomic comparison of a line with phenotypically stable erectoid leaf phenotype ('LT05') with the genomic information obtained from the reference genome 'Heinz 1706' (Tomato Genome Consortium, 2012) and from four accessions with wild-type leaf growth (viz. 'Viradoro', 'TO-937', 'Santa Clara', and 'CNPH498'). The previous confirmation (by mapping) of the chromosomal location of the erectoid leaf phenotype was a key information that allowed us to apply genomic filters directly to chromosome 10, starting with a total number of 27,259 variants and reaching only five candidate genes according to additional genomic and putative gene function analyses (**Table 2**). This filtering allowed a more precise landing on potential candidate genes/loci. The estimated location on the chromosome 10 (close to the *SIGLK2* gene), and the identification of loss-of-function mutation altogether allowed selection of the *Solyc10g009320* as the most likely candidate gene associated with the erectoid leaf growth phenotype. The validation using Sanger sequencing of the F<sub>2</sub> segregating population showed a 100% association of the erectoid leaf phenotype and *Solyc10g009320* mutation.

Due to the structural similarities of *Solyc10g009320* with an array of *TAC1*-like genes (see **Fig. 4**), we tentatively named this gene as *S. lycopersicum TAC1* (*SITAC1*). Our annotation of *Solyc10g009320* as an ortholog of *TAC1* genes is consistent with a recent phylogenetic analysis (Guseman et al. 2017) where this tomato gene was placed into a small cluster along with other *TAC1* genes (e.g. *PpTAC1*, *OsTAC1*, *ZmTAC1*, and *AtTAC1*) within the IGT gene family. This work also identified a second tomato gene (*Solyc01g096260*) within the cluster of *TAC1*-like genes (Guseman et al. 2017). *Solyc01g096260* is located on tomato chromosome 1 and displayed  $\approx 40\%$  amino acid identity with *SITAC1* (**Online Resource 6**). For this reason, we also examined polymorphisms within this gene across our resequenced accessions with contrasting leaf architecture. However, no variants were identified in comparative analyses of the proteins encoded by *Solyc01g096260* in the 'LT05' genome (with erectoid leaf growth phenotype) and the genomes of the four accessions with wild-type leaf phenotype (**Online Resource 7**). These analyses indicated that this evolutionary and functionally-related gene on chromosome 1 has no allelic variation in the coding sequence that could explain the phenotypic impact on the erectoid leaf growth in the germplasm employed in the present study, and indeed the genetic analysis showed that segregation of the locus on chromosome 10 was able to fully explain the occurrence of the erectoid leaf phenotype

without the need to propose the involvement of a second locus. Moreover, the expression patterns reported in the TomExpress RNA Seq database (**Online Resource 8**) show differences to *Solyc01g096260* when compared with *Solyc10g009320*, with a notably greater increase of expression in flowers and fruits and a low or no expression in vegetative parts. Although these data should be confirmed with expression and functional tests, it is likely that these genes, even though belonging to the same family, could be under control of distinct expression mechanisms across distinct plant organs.

It is not yet known how the *TAC1* genes are involved in determining the direction of lateral growth, although some evidence suggests that it would be directly or indirectly implicated in a negative regulation of *LAZY1* (Dardick et al. 2013) and this interaction, which varies depending on the position and aerial organ of the plant, would arise from a gravitropic response. In all cases described in the literature, the loss-of-function of *TAC1* genes (in both monocots and dicots plants) is associated with narrower insertion angles in tillers, leaves, and flowers (i.e. more erect posture) in comparison to the wild-type controls (Yu et al. 2007, Ku et al. 2011, Dardick et al. 2013). This fact is associated with higher auxin content in the affected organs, apparently resulting from a modification in the polar transport of this hormone (Li et al. 2007, Yoshihara and Iino 2007, Yoshihara et al. 2013). Since auxin transport/content is regulating a multiplicity of developmental processes (Reinhardt et al. 2003) mutants involved in this process might have pleiotropic effects in several aspects of agronomic interest.

Auxin also controls many aspects of fruit development, including the sequential stages of fruit formation, expansion, ripening, and abscission (Gillaspy et al., 1993; Srivastava and Handa, 2005). In tomato, Pattison and Catalá (2012) showed a coordinated action of PIN and AUX/LAX proteins in the establishment of auxin gradients during fruit development. Artificially increasing the auxin levels in the ovary can bypass fertilization and lead to the development of parthenocarpic fruits (Lipari & Paratore 1988; Ficcadenti et al. 1999). This would have an interesting effect in extreme/hostile environments where fertilization is compromised by pollen viability problems. Additionally, regulation of auxin efflux by *SIPINI* prevents flower abscission by maintaining a high auxin transport activity in the abscission zone (Shi et al. 2017).

On the other hand, little is known about possible effects of the *TAC1*-like genes at the root system. The pioneering study of Tworkoski and Scorza (2001) in peach provided information that mutations in aerial architecture (including *tac1*) is also associated with changes in root growth pattern. In rice, it was also reported that *DEEPER*

*ROOTING1 (DRO1)* controls the depth of the root system via the regulation of the insertion angle of lateral roots (Uga et al. 2013). An increase of the expression of this factor increases the depth of the root system. The deeper roots determined by *DRO1* allow a better performance and higher yield of the rice in water deficit conditions (Uga et al. 2013). Later, Guesan et al. (2017) generated evidence and hypothesize that, since the functions of *LAZY1* and *DRO1* are homologous, in shoots and roots respectively, *TAC1* could be a negative controller of both genes, and this may explain why a reduced function of *TAC1* could influence the vertical growth both of the canopy and of the roots. In this context, it would be also of interest to investigate the potential effects of the *Erl* locus on the root system in tomatoes.

### **Phenotypic and agronomic effects of the *Erectoid leaf (Erl)* mutation**

So far, no strategies have been proposed to genetically manipulate the inclination angles of lateral shoots in order to obtain tomato ideotypes with adaptation to a wide range of environmental conditions (Zsögön et al. 2017). The importance of leaf insertion angle in relation to efficiency and distribution of light absorption (and hence the photosynthetic crop capacity) has been established in theoretical models using tree/forest species (Pearcy and Yang 1998, Sinoquet et al. 2005). Most studies are in agreement that larger leaf insertion angles (i.e. more horizontal leaves) intercept a considerable greater amount of light when the sun is placed at high angle in relation to the horizon (e.g. midday, summer season, low latitude areas) while smaller leaf insertion angles (i.e. erectoid leaf) intercept a greater portion of light when the sun is at low angles in respect to the horizon (e.g. during early mornings and late afternoons, winter season, and high latitude areas). Falster and Westoby (2003) reinforced these observations by performing studies with several plant species under high radiation conditions, proposing that the cost of increased light interception of horizontal leaves can involve higher leaf temperatures, higher risk of sunscalds and photo-inhibition. Thus, under high radiation conditions, a smaller leaf insertion angle (i.e. more erect leaves) would allow greater protection against the damages caused by the excess of radiation, including the reduction of the heat levels at the leaf surface, increasing the efficiency of water use, minimizing leaf burn damage (Werner et al. 2001), decreasing photo-inhibition (Ryel et al. 1993, Valladares and Pugnaire 1999, Werner et al. 2001) and improving the water use in relation to the daily carbon gain (Cowan 1982). Our data also indicate that is possible to manage the dosage of the *Erl* allele in tomato hybrids, regulating, to a certain extent, the angle of inclination of leaves



and leaflets in relation to the horizontal plane, which in turn, could result in plants with wider adaptation. Higher levels of environmental adaptation would result not only in the increase of the light interception and temperature control, but also by improving the rates of ventilation renewal in the plant canopy. Better ventilation in the plant canopy can influence the CO<sub>2</sub> content as well as the relative humidity of the surrounding air. Another positive crop management consequence of the erectoid leaf trait is to facilitate the distribution of pesticides, especially those targeting the abaxial leaf surface. This leaf surface is the major site of oviposition of important pests such as whiteflies (Silva et al 2014), being also the place of sporulation of a wide array of fungal pathogens. In fact, the erectoid leaf trait may also provide micro environmental conditions that prevent fungal spore germination by maintaining lower humidity levels under the canopy, thus potentially reducing the frequency of fungicide applications. This possible increase in efficiency in pesticide application might reduce both the frequency of sprayings as well as the production costs.

In summary, we found that the genetic factor controlling a peculiar tomato erectoid leaf phenotype (that affects all the aerial organs, especially the young leaf shoots) is located at chromosome 10. We also found that this semi-dominant allele *Erl* co-segregated with a loss-of-function mutation of the gene *Solyc10g009320*. This gene is a strong candidate for the genetic identity of *Erl*. In fact, the protein coded by *Solyc10g009320* has structural features of a TAC1-like proteins of the IGT gene family, which are distributed across a wide array of plant genomes, having a crucial role in the control the vertical growth of the shoots (Dardick et al 2013, Roychoudhry and Kepinski 2015). *TAC1*-like genes regulate auxin polar transport and content in shoots of several species, although it is not possible to discard that *TAC1*-like genes also control auxin transport/content in fruits and roots. From the breeding standpoint, this characterization of the mutant *Erl* will open the possibility for fine-tuning manipulation of tomato architecture and will allow development of more practical co-dominant molecular markers for employment in marker-assistance selection systems of this important trait. In addition, the genetic characteristics (loss-of-function mutation) of *Solyc10g009320* make it a potential target for gene editing strategies (Belhaj et al. 2015; Zsögön et al. 2017).

## REFERENCES

Belhaj K, Chaparro-Garcia A, Kamoun S, Patron NJ, Nekrasov V (2015) Editing plant genomes with CRISPR/Cas9. *Current Opinion in Biotechnology*, 32, 76–84

Boiteux LS, Fonseca MEN, Simon PW (1999) Effects of plant tissue and DNA purification method on randomly amplified polymorphic DNA-based genetic fingerprinting analysis in carrot. *Journal of the American Society for Horticultural Science*, 124, 32–38

Carmo LS, Murad AM, Resende RO, Boiteux LS, Ribeiro SG, Jorrín-Novo JV, Mehta A (2017) Plant responses to tomato chlorotic mottle virus: Proteomic view of the resistance mechanisms to a bipartite begomovirus in tomato. *Journal of Proteomics*, 151, 284–292

Choi Y, Sims GE, Murphy S, Miller JR, Chan AP (2012) Predicting the functional effect of amino acid substitutions and indels. *PLoS One*, 7, e46688

Cowan I (1982) Regulation of water use in relation to carbon gain in higher plants. *Encyclopedia of Plant Physiology*, 12, 589–612

Coyne DP (1980) Modification of plant architecture and crop yield by breeding. *HortScience*, 15, 244–247

Dardick C, Callahan A, Horn R, Ruiz KB, Zhebentyayeva T, Hollender C, Whitaker M, Abbot A, Scorza R (2013) *PpeTAC1* promotes the horizontal growth of branches in peach trees and is a member of a functionally conserved gene family found in diverse plants species. *The Plant Journal*, 75, 618–630

Di Rienzo JA, Casanoves F, Balzarini MG, Gonzalez L, Tablada M, Robledo CW (2013) InfoStat versión 2014. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. URL: <https://www.infostat.com.ar>

Dong H, Zhao H, Xie W, Han Z, Li G, Yao W, Bai X, Hu Y, Guo Z, Lu K, Yang L, Xing Y (2016) A novel tiller angle gene, *TAC3*, together with *TAC1* and *D2* largely determine the natural variation of tiller angle in rice cultivars. *PLoS Genetics*, 12, e1006412

Ehleringer JR, Werk KS (1986) Modifications of solar radiation absorption patterns and implications for carbon gain at the leaf level. In: Givnish T, ed. *On the Economy of Plant Form and Function*, 57–82

Ezcurra E, Montana C, Arizaga S (1991) Architecture light interception and distribution of *Larrea* spp. in the Monte Desert Argentina. *Ecology*, 72, 23–34

Falster DS, Westoby M (2003) Leaf size and angle vary widely across species: What consequences for light interception? *New Phytologist*, 158, 509–525

Ficcadenti N, Sestili S, Pandolfini T, Cirillo C, Rotino GL, Spena A (1999) Genetic engineering of parthenocarpic fruit development in tomato. *Molecular Breeding*, 5, 463–470

Friml J, Wisniewska J, Benkova E, Mendgen K, Palme K (2003) Lateral relocation of auxin efflux transporter *PIN3* mediates tropism in *Arabidopsis*. *Nature*, 415, 806–809

- Georgiev C, Kraptchev B (1992) Genetics investigation of the “erectoid leaf” character in tomato. Report of the Tomato Genetics Cooperative, 42: 14
- Gillaspy G, Ben-David H, Gruissem W (1993) Fruits: A developmental perspective. *Plant Cell*, 5, 1439–1451
- Giordano LB, De Ávila AC, Charchar JM, Ferraz, E, Boiteux LS (2000) ‘Viradoro’: A tospovirus-resistant processing tomato cultivar adapted to tropical environments. *HortScience*, 35, 1368–1370
- Guseman JM, Webb K, Srinivasan C, Dardick C (2017) DRO1 influences root system architecture in *Arabidopsis* and *Prunus* species. *The Plant Journal*, 89, 1093–1105
- Huyghe C (1998) Genetics and genetic modifications of plant architecture in grain legumes: A review. *Agronomie*, 18, 383–411
- Jiao Y, Wang Y, Xue D, Wang J, Yan M, Liu G et al (2010) Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice. *Nature Genetics*, 42, 541–544
- Kinzer S M, Schwager SJ, Mutschler M A (1990) Mapping of ripening-related or-specific cDNA clones of tomato (*Lycopersicon esculentum*). *Theoretical and Applied Genetics*, 79, 489–496
- Ku L, Wei X, Zhang S, Zhang J, Guo S, Chen Y (2011) Cloning and characterization of a putative *tacl* ortholog associated with leaf angle in maize (*Zea mays* L.). *PLoS One*, 6(6), e20621
- Li H, Ruan J, Durbin R (2008) Mapping short DNA sequencing reads and calling variants using mapping quality scores. *Genome Research*, 18, 1851–1858
- Li P, Wang Y, Qian Q, Fu Z, Wang M et al (2007) *LAZY1* controls rice shoot gravitropism through regulating polar auxin transport. *Cell Research*, 17, 402–410
- Lipari V, Paratore A (1988) Parthenocarpy and auxinic treatments in fruiting of tomato in a cold greenhouse. *Acta Horticulturae*, 229, 307– 312
- Pattison RJ, Catalá C (2012) Evaluating auxin distribution in tomato (*Solanum lycopersicum*) through an analysis of the PIN and AUX/LAX gene families. *The Plant Journal*, 70: 585–598
- Pearcy RW, Yang W (1998) The functional morphology of light capture and carbon gain in the Redwood forest understorey plant *Adenocaulon bicolor* Hook. *Functional Ecology*, 12(4), 543–552
- Powell AL, Nguyen CV, Hill T, Cheng KL, Figueroa-Balderas R, Aktas H et al (2012) Uniform ripening encodes a Golden 2-like transcription factor regulating tomato fruit chloroplast development. *Science*, 336(6089), 1711–1715

Reinhardt D, Pesce ER, Stieger P, Mandel T, Baltensperger K, Bennett M, Trass J, Friml J, Kuhlemeier C (2003) Regulation of phyllotaxis by polar auxin transport. *Nature*, 426(6964), 255

Roychoudhry S, Kepinski S (2015) Shoot and root branch growth angle control—the wonderfulness of lateralness. *Current Opinion in Plant Biology*, 23, 124–131

Ryel RJ, Beyschlag W, Caldwell MM. 1993. Foliage orientation and carbon gain in two tussock grasses as assessed with a new whole-plant gas-exchange model. *Functional Ecology*, 7, 115–124

Sarlikioti V, de Visser PHB, Marcelis LFM (2011a). Exploring the spatial distribution of light absorption and photosynthesis of canopies by means of a functional–structural plant model. *Annals of Botany*, 107, 875–883

Sarlikioti V, de Visser PHB, Buck-Sorlin GH, Marcelis LFM (2011b) How plant architecture affects light absorption and photosynthesis in tomato: towards an ideotype for plant architecture using a functional–structural plant model. *Annals of Botany*, 108(6), 1065–1073

Scorza R, Bassi D, Liverani A (2002) Genetic interactions of pillar (columnar), compact, and dwarf peach tree genotypes. *Journal of the American Society for Horticultural Science*, 127(2), 254–261

Scorza R, Lightner GW, Liverani A (1989) The pillar peach tree and growth habit analysis of compact X pillar progeny. *Journal of the American Society for Horticultural Science*, 114, 98–100

Shi J, Dong J, Xue J, Wang H, Yang Z, Jiao Y, Xu L, Huang H (2017) Model for the role of auxin polar transport in patterning of the leaf adaxial–abaxial axis. *The Plant Journal*, 92(3), 469–480

Shi Z, Jiang Y, Han X, Liu X, Cao R, Qi M, Xu T, Li T (2017) SIPIN1 regulates auxin efflux to affect flower abscission process. *Scientific Reports*, 7(1), 14919

Silva KF, Michereff-Filho M, Fonseca, MEN, Silva-Filho JG, Texeira AC, Moita AW, Torres JB, Fernández-Muñoz R, Boiteux LS (2014) Resistance to *Bemisia tabaci* biotype B of *Solanum pimpinellifolium* is associated with higher densities of type IV glandular trichomes and acylsugar accumulation. *Entomologia Experimentalis et Applicata*, 151, 218–230

Silva WB, Vicente MH, Robledo JM, Reartes DS, Ferrari RC, Bianchetti R, Araújo WL, Freschi L, Peres LEP, Zsögön A (2018) SELF-PRUNING acts synergistically with DIAGEOTROPICA to guide auxin responses and proper growth form. *Plant Physiology*, 176(4), 2904–2916

- Sinoquet H, Sonohat G, Phattaralerphong J, Godin C (2005) Foliage randomness and light interception in 3D digitized trees: An analysis from multiscale discretization of the canopy. *Plant, Cell & Environment*, 28(9), 1158–1170
- Srivastava A, Handa, AK (2005) Hormonal Regulation of tomato fruit development: a molecular perspective. *Journal of Plant Growth Regulation*, 24, 67–82
- Teichmann T, Muhr M (2015) Shaping plant architecture. *Frontiers in Plant Science*, 6, 233
- Testa G, Reyneri A, Blandino M (2016) Maize grain yield enhancement through high plant density cultivation with different inter-row and intra-row spacings. *European Journal of Agronomy*, 72, 28–37
- The Tomato Genome Consortium (2012) The tomato genome sequence provides insights into fleshy fruit evolution. *Nature*, 485(7400), 635–641
- Tworokski T, Miller S, Scorza R (2006) Relationship of pruning and growth morphology with hormone ratios in shoots of pillar and standard peach trees. *Journal of Plant Growth Regulation*, 25, 145–155
- Tworokski T, Scorza R (2001) Root and shoot characteristics of peach trees with different growth habits. *Journal of the American Society for Horticultural Science*, 126(6), 785–790
- Tworokski T, Webb K, Callahan A (2015) Auxin levels and MAX1-4 and TAC1 gene expression in different growth habits of peach. *Plant Growth Regulation*, 77(3), 279–288
- Uga Y<sup>1</sup>, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Kitomi Y, Inukai Y, Ono K, Kanno N, Inoue H, Takehisa H, Motoyama R, Nagamura Y, Wu J, Matsumoto T, Takai T, Okuno K, Yano M. (2013) Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. *Nature Genetics*, 45, 1097–1102
- Valladares F, Pugnaire FI (1999). Tradeoffs between irradiance capture and avoidance in semi-arid environments assessed with a crown architecture model. *Annals of Botany*, 83: 459–469
- Vanneste S, Friml J (2009) Auxin: A trigger for change in plant development. *Cell*, 136:1005–1016
- Wang Y, Li J (2008) Molecular basis of plant architecture. *Annual Review of Plant Biology*, 59, 253–279
- Wang B, Smith SM, Li J (2018). Genetic regulation of shoot architecture. *Annual Review of Plant Biology*, 69, 25.1–25.32
- Werner C, Ryel RJ, Correia O, Beyschlag W (2001) Structural and functional variability within the canopy and its relevance for carbon gain and stress avoidance. *Acta Oecologica*, 22, 129–138

Wiśniewska J et al. (2006) Polar PIN localization directs auxin flow in plants. *Science*, 312(5775), 883

Yoshihara T, Iino M (2007) Identification of the gravitropism-related rice gene LAZY1 and elucidation of LAZY1-dependent and –independent gravity signaling pathways. *Plant Cell Physiology*, 48, 678–688

Yoshihara T, Spalding EP, Iino M (2013) AtLAZY1 is a signaling component required for gravitropism of the *Arabidopsis thaliana* inflorescence. *The Plant Journal*, 74, 267–279

Yu B, Lin Z, Li H, Li X, Li J et al. (2007) TAC1, a major quantitative trait locus controlling tiller angle in rice. *The Plant Journal*, 52, 891–899

Zouine M, Maza E, Djari A, Lauvernier M, Frasse P, Smouni A, Pirrello J, Bouzayen M (2017) TomExpress, a unified tomato RNA-Seq platform for visualization of expression data, clustering and correlation networks. *The Plant Journal*, 92, 727–735

Zsögön A, Cermak T, Voytas D, Peres LEP (2017) Genome editing as a tool to achieve the crop ideotype and *de novo* domestication of wild relatives: Case study in tomato. *Plant Science*, 256, 120–130

**Conflict of Interest:** The authors declare that they have no conflict of interest.

## TABLES AND FIGURES

**Table 1** Phenotypic evaluation of an F<sub>2</sub> population (n=274) derived from the cross between ‘LT05’ (erectoid leaf growth; uniform fruit ripening) and ‘TO-937’ with standard (wild-type) leaf growth and green fruit shoulder trait.

	Erectoid leaf growth	No-erectoid leaf growth (intermediate or standard)
Uniform ripening	20	186
Green shoulder	55	13
Fit for expected 3:1 ratio	$\chi^2 > 20$ ( $P < 0.0001$ )	$\chi^2 > 20$ ( $P < 0.0001$ )

**Table 2** Candidate genes located in the genomic region associated with the erectoid leaf growth trait (located at chromosome 10), containing *Solanum lycopersicum* ‘LT05’-specific non-synonymous, single nucleotide polymorphisms (SNPs).

Gene identity	Nucleotide position (SL2.50)	Protein annotation (ITAG 2.40)	SNP position (SL2.50)	Non-conservative amino acid substitution	PROVEAN Prediction Score <sup>1</sup>
<i>Solyc10g005230</i>	184314-188942	Unknown protein	184523	V43M	Neutral (-1.068)
<i>Solyc10g006890</i>	1323124-1333653	WD-repeat protein	1323966	R183C	Neutral (-1.514)
<i>Solyc10g007100</i>	1487998-1492135	Protein detoxification 18	1491582	C56S	Neutral (-0.057)
<i>Solyc10g009320</i>	3390388-3395779	Unknown protein	3394715	Q222*	Deleterious (-61.263)
<i>Solyc10g086620</i>	65405159-65407893	Tropinone reductase homolog At5g06060 isoform X1	65392038	G170*	Deleterious (-248.543)

<sup>1</sup>PROVEAN (Protein Variation Effect Analyzer) is a tool which predicts impact of an amino acid substitution or indel on the biological function of a protein. Variants with a score equal to or below -2.6 are considered ‘deleterious’ and variants with a score above -2.6 are considered ‘neutral’.

**Fig. 1** Vegetative effects of the contrasting erectoid leaf alleles (*Erl*<sup>+</sup> versus *Erl*) in distinct developmental stages of tomato (*Solanum lycopersicum*) plants. **(a)** Plant of the inbred line ‘LT05’, the original homozygous (*Erl*/*Erl*) source of the erectoid leaf phenotype; **(b)** Detail of the foliar insertion angle with respect to the main stem in the ‘LT05’ line. **(c)** Detail of an apical section of a main stem of ‘LT05’ line, showing the effects of the erectoid leaf mutation on the growth of leaflets, leaves, shoots, and trusses. **(d)** Plant of the inbred line ‘LT17’, used as a reference for the standard wild-type phenotype (*Erl*<sup>+</sup>/*Erl*<sup>+</sup>). **(e)** Detail of the foliar insertion angle with respect to the main stem in ‘LT17’ line. **(f)** Plant of ‘IsoL-EL’ isoline (*Erl*/*Erl*). **(g)** Plant of ‘IsoL-WTL’ isoline (*Erl*<sup>+</sup>/*Erl*<sup>+</sup>). **(h)** Heterozygous (*Erl*<sup>+</sup>/*Erl*) F<sub>1</sub> plant derived from the cross of ‘IsoL-EL’ × ‘IsoL-WTL’ with intermediate leaf angle phenotype.

**Fig. 2** Leaf insertion angles in ‘LT05’ × ‘LT17’ in segregating populations. **(a)** Value of the  $\alpha$  angle in ‘LT05’ (erectoid) and ‘LT17’ (wild-type with standard leaf growth) and the reciprocal hybrids generated in both crossing directions F<sub>1</sub> (‘LT05’ × ‘LT17’) and F<sub>1</sub> (‘LT17’ × ‘LT05’). Values shown as mean  $\pm$  standard error (n=13). Treatments with different letters in the bars are significantly different ( $P < 0.0001$ ). **(b)** Phenotypic frequency (mean  $\alpha$  angle of erectoid  $< 100^\circ$ , mean  $\alpha$  angle of intermediate between  $100-125^\circ$  and mean  $\alpha$  angle of wild-type  $> 125^\circ$ ) in three segregating families: (‘LT05’ × F<sub>1</sub>), (‘LT17’ × F<sub>1</sub>) and F<sub>2</sub> (F<sub>1</sub> × F<sub>1</sub>). The Chi-squared test for Mendelian segregation models is shown. **(c) Left:** F<sub>3</sub> families derived from self-pollinating of F<sub>2</sub> plants with erectoid leaf growth phenotype. **Right:** F<sub>3</sub> families derived from self-pollinating of F<sub>2</sub> plants with the standard, wild-type (=non-erectoid) leaf growth phenotype.

**Fig. 3 (a)** Variant density plot of the tomato (*Solanum lycopersicum* L.) chromosome 10. The figure depicts synonymous and non-synonymous variants that were exclusive for the genome of ‘LT05’ (an inbred line with the erectoid leaf growth) in comparison with the genomes of four accessions with standard (wild-type) leaf (viz. ‘Viradoro’, ‘TO-937’, ‘Santa Clara’, and ‘CNPH498’) as well as the reference genome ‘Heinz 1706’; **(b)** Positioning of the single nucleotide polymorphisms – SNPs (black circles) that were selected after applying a filter for non-synonymous variants. The black arrow indicates the position of the linked *SIGLK2* gene (controlling uniform fruit ripening) that was used as phenotypic marker to confirm the chromosome location of the erectoid leaf growth trait; **(c)** Structural features of the *Solyc10g009320* gene, identified as a strong candidate for control of the erectoid growth trait in the tomato line ‘LT05’; **(d)** Genomic sequences of the accessions with standard (= wild-type) and with the erectoid leaf growth. The asterisk marks the position 3,394,715 of chromosome 10, where the ‘LT05’ genome (with the erectoid leaf growth trait) displayed a substitution of a cytosine (C) for a thymine (T) (C>T). This SNP resulted in a change in the CAA codon (=coding for a glutamine) of the wild-type leaf growth to a TAA (= ochre stop-codon) in the ‘LT05’ genome and in F<sub>2</sub> individuals carrying the erectoid leaf growth trait.

**Fig. 4** Multiple alignments of the predicted protein sequences from *TAC1*-like genes of tomato (*Solyc10g009320*), peach (*PpTAC1*), soybean (KRH06413); *Arabidopsis thaliana* (*AtTAC1*), maize (*ZmTAC1*), and rice (*OsTAC1*). The plant species corresponding to each *TAC1* homolog are indicated in left column. Highly conserved residues are highlighted in black, including the conserved G $\phi$ L(A/T)GT domain (at the position 70), which is characteristic of the IGT gene family. The gene *Solyc10g009320* was found to encode a predicted protein of 279 amino acids. The black asterisk highlights the position of the Q222\* on the tomato homolog sequence (corresponding to the position 280 of the consensus alignment) where the stop-codon mutation was found in the *Erectoid leaf* (*Erl*) allele.



a



b



c



d



e



f








g



h





**a****b****c****d**

5' ...AAG CTA <sup>\*</sup>CAA AAA GTG... 3' Standard (wild-type) leaf genome

5' ...AAG CTA TAA AAA GTG... 3' Erectoid leaf genome



	1	10	20	30	40	50	60
Tomato	MK	IFNWVHRKRF	HQKDGKKDIDELKISNEI	-----	IGHH	DTQIL	LDQASF--AHMLDIWNG
Peach	MK	IFNWVHRRL	HQR-VVK--DGFAGNVKK	---	SELETNDK	DTQAF	LKQVGLVNVDGLDGWRD
Soybean	MK	IFNWVHRKRF	HHNTLK--DGFATNMKK	---	TEPTINNT	DSQAL	LKQVAL--TESLGGWED
Arabidopsis	MT	IFNWVQKKL	HQN-VIKEIDGVRKNEKKRN	EGISEIEK	NTKAI	LDQVGL--	VDALDNWFD
Maize	MGLK	VFWLNRMR	HSNTEYCSIHGSKAMEDK	-DSVPHSVVEK	DTREAL	LLHDLV	-----LN
Rice	MALK	VFWLNRKK	HSNVEYCTINENKAMEEKED	SLRASVTEQ	DTREAL	LLRDVL	-----IN
	70	80	90	100	110	120	130
Tomato	IGT	FGFD	-----PLMKNVQDQQSVIDI	ESLEDEEEEILEEEEEYYSV	ENEIQECEIPFNDHEGID	BELY	
Peach	IGT	FGFD	-----PL-KPSTHQNEYFVL	ESEEDDQ---	ESHGFSHSGNDDDDDDDEHY	-DHSVED	BELN
Soybean	IGT	LGYD	-----PL-KSINPHKEYFAL	EIEVEQE--	DGEGENNIMVDDHQEE	EGY---NAAH	BELN
Arabidopsis	IGT	FGFD	-----TL-----	NFKE	DEMDDDECGSGDL	DYVVIDGSI	IKN-----VD
Maize	IGT	LGHVD	SLCPV-ACIEEDDLFIMD	EKAIVEERDDEEPRNDQVKED	VALETEPC-----	BPVW	
Rice	IGT	LGHVN	SLCPE-SCIEQDEPIIMC	EKEVQEKECEEKAE	-----AKQDTPV---	TAPS	EPAS
	140	150	160	170	180	190	200
Tomato	P	IYANIGDEM	IYNEDNIESNN	-----	NSFTDQTNTKMM	-----	KKER
Peach	P	LMFTTTFEHSFED	IGSNFDAIVQKPADVILTV	--DGVPLTPFEGSSEISTKPDQ	SANDQSKNK	--KGQR	
Soybean	P	IYNSFEHNFV	--DVSENNHDGDVVVRSFNDEIVVAPPVISY	EVMESNDVDADAEK	--KK	KGER	
Arabidopsis	P	I-----	SVENKFYDHHEDVGNLYLQADTDHF	-----	GSIKTAETPAVTA	AAAAEIEP	QKKR
Maize	P	I	VEPAKMHSLSMKEDNFTCFVAE	--EILIREVEDG	-----GGANIQERPLLMLE	-----	KVEKVR
Rice	A	I	EPAKMHSSSMKEDNFMCFVKE	--EILMHGMEVE	-----DVPNIQERPLLMLE	-----	KVEKVR
	210	220	230	240	250	260	270
Tomato	I	TLADLF	SADS-D-----HHHNKANRRSVRESEIFTKKSNS	SSPQVKNGLSFA	KK--	LIPRV	KDEPR-
Peach	I	TLADLF	QADVPD-----VGQLKLD--SGKVQPEMEKKMNA	--RTRSGLAFA	KK--	LIPRV	KDDSS-
Soybean	I	TLADLF	LADT-D-----V-KIKLDPYAKVCHLASSEKPNL	--KAKHGLSFA	KK--	LIPRV	KDNPH-
Arabidopsis	T	TLAEL	FMEDH-DKGYDTHWCKKPNPNLDGEEVKYKQNGS	--KLSKRFSFV	KKLVM	SKS	KEEND
Maize	T	TLADLF	FAAEV-----FSSSSPREYSCQDIIDVAVAS	--TSKHASCM	E	MHQK	KPR-KPTPKP
Rice	T	TLADLF	FAAEA-----FSSSDAEDKCYPKIVIVAGAS	-TSKPTSCME	E	MHHK	KPT-KPTSKP
	280	290	300	310	320	330	338
Tomato	--PI	QK	QKLM	T	KVL	K	RKVHP-D
Peach	--PI	KN	QRLM	R	RML	K	RKIHPA
Soybean	--PM	QD	TK	KLM	K	RML	KRKIHP-D
Arabidopsis	LLPI	KK	IR	QMI	K	RML	KRKIHP-D
Maize	LKAT	KK	SRVM	K	RML	G	KKIHP
Rice	LKAT	KK	SRVM	K	RML	G	KKIHP

# A loss-of-function allele of a TAC1-like gene (SITAC1) located on tomato chromosome 10 is a candidate for the Erectoid leaf (Erl) mutation

González-Arcos, Matías

2019-04-16

Attribution-NonCommercial 4.0 International

---

González-Arcos M, de Noronha Fonseca ME, Zandonadi DB, Peres LE, Arruabarrena A, Ferreira DS, Kevei Z, Mohareb F, Thompson AJ and Boiteux LS, A loss-of-function allele of a TAC1-like gene (SITAC1) located on tomato chromosome 10 is a candidate for the Erectoid leaf (Erl) mutation. *Euphytica*, Volume 215, May 2019, Article number 95

<https://doi.org/10.1007/s10681-019-2418-1>

*Downloaded from CERES Research Repository, Cranfield University*