

ARTICLE OPEN



Quantifying non-CO₂ contributions to remaining carbon budgets

Stuart Jenkins¹✉, Michelle Cain^{1,2}, Pierre Friedlingstein^{3,4}, Nathan Gillett⁵, Tristram Walsh¹ and Myles R. Allen^{1,6}

The IPCC Special Report on 1.5 °C concluded that anthropogenic global warming is determined by cumulative anthropogenic CO₂ emissions and the non-CO₂ radiative forcing level in the decades prior to peak warming. We quantify this using CO₂-forcing-equivalent (CO₂-fe) emissions. We produce an observationally constrained estimate of the Transient Climate Response to cumulative carbon Emissions (TCRE), giving a 90% confidence interval of 0.26–0.78 °C/TtCO₂, implying a remaining total CO₂-fe budget from 2020 to 1.5 °C of 350–1040 GtCO₂-fe, where non-CO₂ forcing changes take up 50 to 300 GtCO₂-fe. Using a central non-CO₂ forcing estimate, the remaining CO₂ budgets are 640, 545, 455 GtCO₂ for a 33, 50 or 66% chance of limiting warming to 1.5 °C. We discuss the impact of GMST revisions and the contribution of non-CO₂ mitigation to remaining budgets, determining that reporting budgets in CO₂-fe for alternative definitions of GMST, displaying CO₂ and non-CO₂ contributions using a two-dimensional presentation, offers the most transparent approach.

npj Climate and Atmospheric Science (2021)4:47; <https://doi.org/10.1038/s41612-021-00203-9>

INTRODUCTION

The IPCC's Special Report on Global Warming of 1.5 °C¹ (SR1.5) concluded: 'Reaching and sustaining net zero global anthropogenic CO₂ emissions and declining net non-CO₂ radiative forcing would halt anthropogenic global warming on multi-decadal timescales. The maximum temperature reached is then determined by cumulative net global anthropogenic CO₂ emissions up to the time of net zero CO₂ emissions and the level of non-CO₂ radiative forcing prior to the time that maximum temperatures are reached.' This highlights the importance of future cumulative CO₂ emissions, often termed the 'remaining carbon budget'^{2–5}, together with the increasingly important role of non-CO₂ climate drivers as peak warming is approached. SR1.5 did not, however, give any further scenario-independent quantification of this statement, beyond noting that an increase of 1 W/m² of non-CO₂ radiative forcing and a cumulative emission of 1000 GtCO₂ 'represent approximately equal effects on global mean surface temperature (GMST).' Here we provide this quantification.

The carbon budget framing is helpful because most warming to date has been caused by CO₂^{6–8}, and CO₂, of all major pollutants, has the most permanent impact on the climate system^{8–10}. CO₂-induced warming is approximately proportional to the total quantity of CO₂ emitted over any multi-decade time interval, where the constant of proportionality is termed the Transient Climate Response to cumulative carbon Emissions, or TCRE^{5,11}. There are, however, complications^{1,2,9,12} in the use of TCRE to derive the remaining carbon budget, including the precise definition and estimated current level of global warming; committed warming due to past CO₂ emissions, or the zero emissions commitment (ZEC); possible contributions of Earth System Feedbacks to future warming; uncertainty in the estimated value of the TCRE; and the future contribution of non-CO₂ climate pollutants. Of these, the contribution of non-CO₂ pollutants is

unique in that it depends on future policy decisions, not simply scientific uncertainty.

We focus here on carbon budgets corresponding to increases in GMST because this remains the index used to report observed increases in global temperatures^{1,13,14}, and hence may be used to determine when a temperature threshold is reached. Previous studies have suggested that changing sea-ice cover precludes the use of GMST in projections¹⁵. However, Fig. 3f of ref. ¹⁵ indicates the very limited impact of sea-ice retreat on GMST under ambitious mitigation scenarios, while under sustained warming the impact would correspond to a couple of years of warming at most. Therefore, changing sea-ice cover does not present any fundamental impediment to the use of GMST in projections. Further, while in earlier GMST products the ratio of GSAT to GMST has differed by several percent¹⁶, recent updates to GMST datasets have largely accounted for these differences using statistical infilling of undersampled geographical regions.

Here, headline conclusions are communicated for a global temperature anomaly calculated from a four-dataset mean of these statistically infilled GMST products (similar to the approach taken in ref. ¹⁷) to reduce the impact of any individual dataset, and to ensure that our conclusions are consistent with the estimates of the current level and rate of increase of human-induced global warming (see 3,13,17). Recent updates to these GMST datasets have revised the present decade's warming level up compared to earlier products. This presents a hazard for policymakers: using different indices to report observed warming (e.g. the reference period chosen in the Structured Expert Dialogue¹⁴ informing the Paris Agreement) and to calculate carbon budgets¹⁵ raises the possibility of the carbon budget being exhausted well before a temperature threshold is reached, potentially undermining confidence in the entire construct. Following SR1.5, we focus on budgets consistent with halting warming for a multi-decade period, acknowledging uncertainty in the level of positive or

¹AOPP, Department of Physics, University of Oxford, Oxford, UK. ²Centre for Environmental and Agricultural Informatics, Cranfield University, Cranfield, United Kingdom. ³College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter EX4 4QF, UK. ⁴Laboratoire de Meteorologie Dynamique, Institut Pierre-Simon Laplace, CNRS-ENS-UPMC-X, Departement de Geosciences, Ecole Normale Supérieure, 24 rue Lhomond, 75005 Paris, France. ⁵Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC, Canada. ⁶Environmental Change Institute, School of Geography, University of Oxford, Oxford, UK. ✉email: stuart.jenkins@wadhams.ox.ac.uk

negative emissions that may be required to maintain stable temperatures in the very long term thereafter¹⁸. This allows us to assume the ZEC is negligible and ignore long-term Earth system feedbacks. Most studies find the ZEC contributes at most a small amount to remaining warming under ambitious mitigation scenarios^{10,19–21}.

Most recent estimates of a remaining ‘multi-gas’ carbon budget rely on subtracting a distribution of warming responses to non-CO₂ sources from the target total warming and estimating a CO₂ budget for the remainder^{1,2,22}. This approach relies on: (1) the careful treatment of covarying physical climate response uncertainty to both CO₂ and non-CO₂ contributions and (2) available scenarios from integrated assessment models (IAMs) representing a statistical distribution of possible futures. Guidelines on the use of scenarios in the SR1.5 database make clear they should not be treated as a statistical distribution²³, since they rely on prescriptive, often normative, decisions such that the choice of model has more impact than within-model uncertainties (we show this in Fig. 3). Scenarios representing the most ambitious temperature goals also depend on which IAM set-ups converge at all, an even more arbitrary and opaque constraint. Therefore, percentiles of available scenarios cannot be used to estimate the ‘likely’ non-CO₂ contribution to warming. Further, a predetermined quantity of non-CO₂ warming should not be subtracted from the total remaining warming without considering the accompanying impact of covarying physical climate uncertainty implicit in the choice of TCRC. Refs. ^{1,2,24} all remove a quantity of warming attributed to non-CO₂ pollutants independent of the sampled TCRC percentile. A more transparent treatment of non-CO₂ climate

drivers uses CO₂-forcing-equivalent (CO₂-fe) emissions²⁵, meaning the CO₂ emissions time series that would give precisely the same impact on effective radiative forcing (ERF) and thence GMST. This is similar to the approach of ref. ²⁶, although they use a single representative non-CO₂ forcing scenario. By doing this we can explicitly sample the physical climate response uncertainty for both CO₂ and non-CO₂ contributions identically, and more clearly separate scenario and physical response uncertainties in non-CO₂ contributions. We are therefore here assuming the climate response to effective non-CO₂ forcing is identical to the response to the same level and time-history as CO₂ forcing; while still an assumption, this is clearly preferable to assuming these responses are independent.

RESULTS

CO₂-forcing-equivalent emissions in mitigation scenarios

Originally proposed by Tom Wigley in 1998 under the name of a ‘Forcing-Equivalent Index’, CO₂-fe emissions²⁷ express an emissions time series of any climate pollutant in terms of the time series of CO₂ emissions that would have an identical impact on ERF, and hence GMST on all timescales. They are obtained by converting the ERF associated with that pollutant to a time series of change in CO₂-equivalent concentrations, and then computing the CO₂ emissions required to produce that CO₂ concentration perturbation using a carbon cycle model²⁵ (see Methods for a full explanation of the CO₂-fe methodology along with a simple formula which approximates the full calculation).




Fig. 1 IIAA IAMC database of scenarios in the IPCC Special Report on the Global Warming of 1.5°C. Panel **a** plots the annual CO₂ emissions. Panel **b** (below **a**) shows the running sum (or cumulative) CO₂ emissions from 2018. Panel **c** (bottom right) shows the non-CO₂ radiative forcing for each scenario (dotted lines, right hand axis). Also on panel **c** are the cumulative non-CO₂ CO₂-fe emissions from 2018 corresponding to each non-CO₂ RF line (solid lines, left-hand axis). The axes of panels **b** and **c** are scaled so the cumulative emissions from CO₂ and non-CO₂ are directly comparable. Panel **d** plots the FaIRv2.0-derived temperature response against the diagnosed cumulative CO₂-fe emissions (solid lines) and against the cumulative CO₂-only emissions (dotted lines). For FaIR temperature response TCR = 1.8 °C, ECS = 3.0 °C. Scenarios are coloured by category in the IAMC database: red for 2°C-higher, orange for 2°C-lower and blue for 1.5°C-compatible. Light blue scenarios in panel **c** highlight some example non-CO₂ pathways (lower bound, upper bound and a central scenario from 1.5°C-compatible dark blue plume) with the central light blue scenario (P3 scenario from SR15 SPM.3b) also used in Figs. 2 and 3.

Removing a temperature contribution from non-CO₂ forcing agents would be a valid approach to carbon budget estimation so long as the physical climate response uncertainty were sampled at the same percentile for both CO₂ and non-CO₂ warming contributions. SR1.5, and similar approaches^{2,24}, remove a predetermined range of non-CO₂ warming before the TCRE is sampled. This is inconsistent since a low non-CO₂ warming contribution is significantly more likely if a low TCRE is sampled. A more transparent approach uses CO₂-fe: the scenario uncertainty (fraction of total CO₂-fe budget which is allocated to non-CO₂ forcing agents) is separated from the physical climate behaviour governing the overall size of the cumulative all-pollutant budget.

In contrast to conventional emission metrics, there is no need to specify an arbitrary time-horizon to compute CO₂-fe emissions, since the CO₂ emissions required to produce a particular pathway of concentration anomalies are unambiguously determined by the behaviour of the carbon cycle. IPCC's SR1.5 did not systematically include CO₂-fe in the estimation of remaining budgets²⁸ (it did address alternative metrics which approximate the behaviour of CO₂-fe, namely GWP^{29–31}, but not in the analysis of carbon budgets). There is a clear need for complementary approaches, given that CO₂-fe is no less accurate than the temperature anomaly-based approach defined in SR1.5, and offers greater transparency since non-CO₂ forcing uncertainty is more clearly separated from physical climate uncertainty.

One benefit of the CO₂-fe metric is that it allows direct comparison between the CO₂ and non-CO₂ contributions to warming. In Fig. 1a we plot a number of scenarios for future CO₂ emissions from the IIASA SR1.5 scenario database⁷. They are coloured by ambition according to their label in the database; dark blue corresponds to scenarios tagged as '1.5 °C-compatible', light orange corresponds to 'lower-2 °C-compatible', and dark orange corresponds to 'higher-2 °C-compatible'. Panel b below shows the cumulative CO₂ emissions relative to 2018, which can be translated into the CO₂ warming contributions by multiplying by the TCRE. Panel c shows the corresponding non-CO₂ ERFs for each CO₂ emissions pathway (dotted lines, right axis). We would like to compare these scenarios to the CO₂ emissions, but we cannot simply apply the TCRE as we did for CO₂ as the total warming is non-linear when plotted against cumulative CO₂ emissions alone (dotted lines in panel d). If, instead, we express the non-CO₂ ERFs as cumulative CO₂-fe emissions, they are now physically equivalent quantities and the cumulative CO₂/CO₂-fe

emissions time series in panels b and c (solid lines, left axis) can be directly compared. This would not be possible with CO₂-equivalent emissions calculated using the GWP or GTP metrics, which do not accurately reproduce the warming outcome for a complex multi-gas emissions pathway²⁹.

By converting the full range of pollutants into cumulative CO₂-fe emissions we can use the TCRE in the same way we did for CO₂ alone. Figure 1d shows global temperatures plotted against cumulative total CO₂-fe emissions (solid lines). Cumulative total CO₂-fe emissions multiplied by the TCRE predicts the temperature response, just as in a pure-CO₂ scenario. If non-CO₂ radiative forcing were correlated with cumulative CO₂ emissions in these scenarios, then the latter would also predict the response with a simple scaling factor, or 'effective TCRE', to account for a constant fractional contribution to warming from non-CO₂ drivers^{3,32,33}. Figure 1d shows this is not always the case (nor is there any physical reason for it to be the case in complex multi-gas future scenarios)³⁴; hence the impact of non-CO₂ forcing needs to be treated explicitly.

Having explained the utility of the CO₂-fe metric for assessing the relative CO₂ and non-CO₂ contributions to warming in mitigation scenarios, we now turn to a single scenario and explore the contributions from individual pollutants in greater detail. The three light blue scenarios in Fig. 1 display the range of non-CO₂ RF pathways exhibited in 1.5 °C-compatible scenarios, with the central light blue pathway highlighting the P3 scenario from SR1.5's SPM Fig. 3b¹ (middle of the road scenario which achieves 1.5 °C ambition). Figure 2 shows a breakdown of the total CO₂-fe emissions time series for this central light blue scenario in Fig. 1c. We extend back to pre-industrial using the historical RF time series from ref. ³⁵, and calculate individual CO₂-fe emissions contributions using a differencing approach (see Methods). These individual contributions are stacked and coloured by pollutant, with the panel a showing the annual CO₂-fe emissions, and panel b showing the same cumulatively. In contrast to CO₂-equivalent emissions, whether computed with GWP₁₀₀ or any other conventional metric, CO₂-fe emissions reflect the impact of individual climate drivers on global temperature (panel c), allowing them to be compared objectively. CO₂-fe provides a transparent and easily implementable approach which translates readily to warming: cumulative emissions contributions in panel 2b correspond to warming levels in panel 2c (we have used a TCRE of 0.4 °C/TtCO₂—our best-estimate TCRE found using




Fig. 2 Central light blue 1.5 °C-compatible scenario from Fig. 1 displayed showing component contributions to warming. Panel a plots the annual CO₂ emissions and CO₂-fe emissions for each of the major contributing pollutants (red = CO₂, blue = CH₄ + Ozone + Strat. H₂O, green = N₂O, gold = Other and orange = Aerosols). Panel b shows the corresponding cumulative CO₂ and CO₂-fe emissions time series (stacked by contribution to total). Panel c plots the temperature response for each component. Black solid lines show the total annual (panel a) and cumulative (panels b and c) CO₂-fe emissions, while the total temperature response is shown with a black dotted line (all calculated from total RF). Small backscatter points on panel c show the annual temperature observations using four-dataset mean observations updated from SPM.1, SR1.5. FaIR-derived temperatures (panel c) use thermal parameters chosen to best emulate historical temperatures (TCR = 1.8 °C, ECS = 3.0 °C). RFs before 2020 are taken for individual components from Dessler and Forster (2018) RF dataset, with future component RF rescaled to match the component's best-estimate historical RF in 2020. Methane forcing is scaled by 1.65 to account for Ozone and stratospheric H₂O contributions, following refs. ^{40,30}. Aerosol forcing is ~-0.9 W/m² in 2011, consistent with recent observationally constrained estimates (e.g. Stevens et al., 2015; AR5, 2013). Thin dotted lines show present-day best-estimate anthropogenic warming level (1.23 °C in 2020).

an observationally constrained methodology described in section 3 below).

Methane emissions make a net positive contribution to annual CO₂-fe emissions until they begin to rapidly decline. Thereafter, the short atmospheric residence time of methane³⁰ means that falling methane emissions give a declining radiative forcing, equivalent to negative CO₂-fe emissions. Aerosol cumulative CO₂-fe emissions in Fig. 2b are negative, but show a sharp increase towards zero in the 2020s and 2030s, corresponding to a high associated CO₂-fe emissions rate (orange in Fig. 2a), and consequently a rapid removal of the cooling effect that aerosols have been contributing over history. Long-lived pollutants like nitrous oxide behave like CO₂. Note that the calculation of CO₂-fe emissions is not model specific: with a linear CO₂ impulse response (IR) function (as used for the calculation of GWPs) one can calculate CO₂-fe emissions with a simple matrix inversion and get very similar results (see Methods and ref. ³⁶). The accuracy of CO₂-fe emissions for representing the radiative forcing and therefore temperature impacts of long- and short-lived pollutants gives it clear advantages over GWP₁₀₀ for presenting mitigation scenarios aimed at limiting global warming²⁹.


Total cumulative CO₂-fe emissions and total anthropogenic warming are approximately proportional to the combined warming impact of CO₂ and methane, as indicated by cumulative CO₂-plus-methane CO₂-fe emissions, up to the present-day (red and blue in Fig. 2b, c), but diverge rapidly over the coming decades as aerosol forcing declines (orange). Strikingly, this aerosol decline contributes almost as much to future warming as the remaining CO₂ emissions in this central 1.5 °C-compatible scenario (panel c), highlighting the importance of common and comparable presentations of all climate drivers. Aerosols are often not included in figures showing multi-gas emission scenarios¹⁹ because of the lack of a nonarbitrary way of displaying them on a common axis. This problem is resolved by CO₂-fe. Individual contributions to CO₂-fe emissions from 2020 to the time of peak warming under this scenario are CO₂: 555 GtCO₂-fe; methane: −65 GtCO₂-fe; nitrous oxide: 65 GtCO₂-fe; aerosols: 290 GtCO₂-fe; other forcings: −45 GtCO₂-fe; giving a total CO₂-fe budget of 800 GtCO₂-fe.

Observational constraints on the TCRE

Having demonstrated how the TCRE can be extended to multi-gas scenarios using CO₂-fe emissions, we now consider how CO₂-fe emissions can be used to investigate the TCRE itself. Previous TCRE estimates³⁷ have compared cumulative pure-CO₂ emissions with warming attributable to CO₂ alone, but the fractional uncertainty in the latter is higher than uncertainty in total anthropogenic warming. Comparing total anthropogenic warming with total cumulative CO₂-fe emissions over the historical record presents a useful complementary approach.

To estimate anthropogenic warming over the historical period, we use conventional ‘optimal fingerprinting’ applied to GMST, as is defined in Haustein et al. (2017)³⁸. This uses a two-timescale IR model^{9,39,40} to estimate temperature responses to anthropogenic and natural forcing (for which we use a 1000-member ensemble of representative ERF time series³⁵). A four-dataset mean GMST observation time series is then regressed onto each pair of natural and anthropogenic temperature response time series to determine the most likely contribution from each component, with added CMIP6 PControl simulations included in the regression to account for internal climate variability. For a full discussion see Methods. Estimated anthropogenic warming in 2020 relative to 1850–1900 is 1.23 °C (1.06–1.42 °C) (5–95% confidence interval), higher than ref. ¹ (SR1.5) due to updates in the datasets.

We express the 1000 anthropogenic ERF time series as CO₂-fe emissions pathways²⁵ accounting for uncertainty in cumulative CO₂ airborne fraction to date (0.4 ± 0.04 ⁴¹) in carbon cycle



parameters. Dots in Fig. 3a show estimates of the total anthropogenic warming and cumulative all-pollutant CO₂-fe emissions, coloured by decade. For example, pink dots in Fig. 3a sample the resulting joint distribution of cumulative anthropogenic CO₂-fe emissions 1875 to 2015 inclusive and human-induced warming to the decade 2011–2020 relative to 1850–1900.

Fig. 3 Observational constraints on the TCRE and consequences for the design of Paris Agreement-compatible scenarios. Panel **a** plots attributed human-induced warming against cumulative emissions of CO₂. The space is shaded by the value of the TCRE (Gaussian distribution in best-estimate and likely range in AR5) and the points are coloured by the decade in which the temperature (relative to 1850–1900 baseline) and cumulative CO₂-fe emissions (relative to 1875) are diagnosed. An ellipse is drawn around central 90% of points. Black lines in panel **a** depict the 5th, 50th and 95th percentile of the overall observationally constrained TCRE distribution based on the 2011–2020 decade. Panels **b–d** show the remaining CO₂ and non-CO₂ CO₂-fe budgets for 2020 for each scenario in Fig. 1, coloured by category in the IIASA SR15 database. Panels **b** and **d** show remaining budgets to peak warming in each scenario, while panel **c** shows budgets to 2100 (instead coloured by IAM). In all three panels shading shows budgets compatible with limiting warming to 1.5 °C for the AR5 gaussian TCRE likely range, as in panel **a**. The solid black lines show the corresponding remaining total CO₂-fe budgets instead of using our observationally constrained TCRE 5th–95th percentile range. In panels **b** and **c** the shading, therefore, corresponds to budgets for 0.27 °C remaining warming to 1.5 °C-consistent with 1.23 °C warming in 2020. In panel **d** the shading refers to budget for 0.42 °C remaining warming to 1.5 °C-consistent with 1.08 °C warming in 2020 (re-baselining historical GMST to 0.85 °C for the decade prior to 2015). Pink horizontal box-whisker plots in panels **b** and **d** show estimates of remaining CO₂ budgets for each assumed present-day warming level, assuming a mid-range non-CO₂ budget to peak warming (130 GtCO₂-fe) and plotting the 5th, 33rd, 66th and 95th percentiles (see Methods for information on calculation). In panel **c** IAM abbreviations correspond to: A/C 2.0/2.1 – AIM/CGE 2.0/2.1; C-R 5.005 – C-Roads 5.005; I 3.0.1 – IMAGE 3.0.1; M V.3 – MESSAGE V.3; M-G 1.0 – MESSAGE-GLOBIOM 1.0; R 1.5/1.7 – REMIND 1.5/1.7; R-M 1.7-3 – REMIND-MAqPie 1.7-3.0.

The cumulative anthropogenic CO₂-fe emissions and human-induced warming estimate for each dot correspond to the same ERF time series to account for any covariance, while CO₂ airborne fraction and internal climate variability are sampled independently. Shading shows the AR5 gaussian TCRE distribution, with the likely range and median values highlighted. Ellipses are drawn around each decade's scatter of covarying temperature anomaly and cumulative CO₂-fe emissions, encompassing the central 90% of the distribution, also coloured by decade. The best-fit TCRE is estimated as 0.40 °C/TtCO₂ (0.26–0.78 °C/TtCO₂ 90% confidence interval based on the most recent decade), marked with black lines in panel a. These could be interpreted as median and 5–95% percentiles of a probability distribution if the input ERF pathways are assumed to be equiprobable, but more research characterising the distribution of uncertainty in radiative forcing to date is needed⁴².

For comparison, SR1.5 uses a likely TCRE range of 0.22–0.68 °C/TtCO₂ taken from the assessment in AR5’s WG1 (SR1.5 also includes a 100–200 GtCO₂ budget correction accounting for differences between gaussian and log-normal TCRE distributions), while TCRES derived from the CMIP6 1%/yr CO₂ concentration increase experiment lie in the range 0.36–0.63 °C/TtCO₂ (see supplementary fig. 8, or ref. ⁴³). Other groups have separately diagnosed these (e.g. Williams et al. ⁴⁴), noting additionally that the inter-model spread is strongly affected by cloud feedbacks, particularly in high sensitivity models. Mengis and Matthews (2020) use the CO₂-fe metric to demonstrate the bias introduced by assuming a constant fractional non-CO₂ contribution to warming in TCRE assessments and estimate the TCRE using a single warming pathway (~0.5 °C/TtCO₂)³⁴. Matthews et al. (2021) estimate an observationally constrained TCRE by removing a fractional warming contribution attributed to non-CO₂ pollutants, finding a median TCRE of 0.44 °C/TtCO₂ (0.32–0.62 °C/TtCO₂ 5–95th percentile range)⁴⁵. We argue the contribution from

Table 1. TCRE percentiles and corresponding remaining carbon budgets to 1.5°C.

| Percentile | TCRE (GMST) °C/ TtCO ₂ | 1.5 °C remaining CO ₂ -fe budget (TtCO ₂) (0.27 °C) | 1.5 °C remaining CO ₂ -fe budget (TtCO ₂) (0.42 °C) | 1.5 °C remaining CO ₂ budget (TtCO ₂) (0.27 °C) 130 (50–300) GtCO ₂ -fe non-CO ₂ budget | 1.5 °C remaining CO ₂ budget (TtCO ₂) (0.42 °C) 130 (50–300) GtCO ₂ -fe non-CO ₂ budget | SR1.5 1.5 °C remaining CO ₂ budget (TtCO ₂) (GMST) | SR1.5 1.5 °C remaining CO ₂ budget (TtCO ₂) (GSAT) | Rogelj et al (2019) 1.5 °C remaining CO ₂ budget (TtCO ₂) (GSAT) |
|------------|--------------------------------------|--|--|--|--|---|---|---|
| 5th | 0.26 | 1.04 | 1.62 | 0.91 (0.74–0.99) | 1.49 (1.32–1.57) | – | – | – |
| 17th | 0.31 | 0.87 | 1.35 | 0.74 (0.57–0.82) | 1.22 (1.05–1.30) | – | – | – |
| 33th | 0.35 | 0.77 | 1.20 | 0.64 (0.47–0.72) | 1.07 (0.90–1.15) | 1.08 | 0.84 | 0.74 |
| 50th | 0.40 | 0.68 | 1.06 | 0.55 (0.38–0.63) | 0.93 (0.76–1.01) | 0.77 | 0.58 | 0.48 |
| 66th | 0.46 | 0.59 | 0.92 | 0.46 (0.29–0.54) | 0.79 (0.62–0.87) | 0.57 | 0.42 | 0.32 |
| 83th | 0.57 | 0.47 | 0.73 | 0.34 (0.17–0.42) | 0.60 (0.43–0.68) | – | – | – |
| 95th | 0.78 | 0.35 | 0.54 | 0.22 (0.05–0.30) | 0.41 (0.24–0.49) | – | – | – |

Percentiles of the TCRE found in this study are shown in column 2. CO₂-fe budgets for 0.27 °C additional warming (1.5 °C-compatible using GMST) and 0.42 °C additional warming (1.5 °C-compatible using GMST re-baselined to 0.85 °C over the decade up to 2015) shown in columns 3 and 4, respectively. Corresponding CO₂-only budgets to 1.5 °C are shown in columns 5 and 6, with mid-range non-CO₂ contribution (range shown in brackets for a full range of 1.5-compatible non-CO₂ contributions). For comparison, the headline remaining budget estimates in SR1.5 and Rogelj et al. (2019) are shown in columns 7, 8 and 9.

Percentiles of the TCRE found in this study are shown in column 2. CO₂-fe budgets for 0.27 °C additional warming (1.5 °C-compatible using GMST) and 0.42 °C additional warming (1.5 °C-compatible using GMST) re-baselined to 0.85 °C over the decade up to 2015) shown in columns 3 and 4, respectively. Corresponding CO₂-only budgets to 1.5 °C are shown in columns 5 and 6, with mid-range non-CO₂ contribution (range shown in brackets for a full range of 1.5-compatible non-CO₂ contributions). For comparison, the headline remaining budget estimates in SR1.5 and Rogeli et al. (2019) are shown in columns 7, 8 and 9 (range shown in brackets for a full range of 1.5-compatible non-CO₂ contributions).

non-CO₂ pollutants should be determined explicitly using the CO₂-fe methodology.

The remaining total CO₂-fe emissions budgets for an additional 0.27 °C warming above 2020, corresponding to total warming of 1.5 °C, range between 350–1040 GtCO₂-fe. A detailed breakdown by percentile are shown in Table 1 (see Supplementary Table 1 for equivalent budgets to 2 °C). We calculate the remaining budgets for additional anthropogenic warming relative to the best-estimate current level (1.23 °C) for consistency with Table 2.2 of ref. ¹, reflecting a policy focus on future warming relative to the recent past rather than including uncertainty in pre-industrial temperatures. The wide range of remaining total CO₂-fe budgets we find here are largely a result of the range of present-day RF in our 1000-member total anthropogenic RF ensemble. Reducing RF component uncertainty and accounting for correlations between component RFs would better constrain this range, and is a focus for future research. Here we focus on defining the methodology to estimate the TCRE with CO₂-fe.

Given these total remaining CO₂-fe budgets, the question now becomes what fraction of this budget is used by CO₂ and non-CO₂ pollutants respectively. Figure 3b shows the proportions of the future total CO₂-fe budget allocated to CO₂ and non-CO₂ in these scenarios. Shading indicates the remaining total budget compatible with 0.27 °C additional warming, again using the gaussian TCRE distribution reported in AR5 as in panel a, while solid black lines indicate the total remaining budget using our observationally constrained TCRE distribution from panel a. Scatter points indicate cumulative CO₂ and non-CO₂ CO₂-fe emissions to peak warming in 1.5 °C-compatible, 2 °C-lower and 2 °C-higher scenarios from Fig. 1, with the colours indicating the scenario category as in Fig. 1. In these scenarios, the non-CO₂ contribution to the total remaining budget ranges from 50–300 GtCO₂-fe, exactly the range determined in SR1.5 (where 250 GtCO₂ budget uncertainty was attributed to non-CO₂ scenario uncertainty). Depending on the TCRE, this means non-CO₂ scenario uncertainty contributes between 0.01 and 0.23 °C warming in these scenarios (using the observationally constrained TCRE 5th–95th percentile range). Importantly, dark blue 1.5 °C-compatible scenarios are consistent with their 1.5 °C peak warming categorisation—the scatter of blue dots sits over our best-estimate TCRE.

While individual scenario's CO₂ and non-CO₂ contributions to peak warming are shown in Fig. 3b, panel c instead plots the scenario's budgets out to 2100. Scenario categories are less clear if the budget is based on end of century warming—scenario categorisation appears overly conservative if the remaining budget is allocated to 2100, with around half of the 1.5 °C-consistent scenarios lying outside the likely range. Net-negative CO₂ emissions and declining non-CO₂ radiative forcing after mid-century reduce the cumulative contributions from both CO₂ and non-CO₂ pollutants (see Fig. 1b, c).

Further, Fig. 3c shows how this non-CO₂ contribution is influenced by the IAM choice. Colours indicate the IAM used to generate each scenario, confirming this is not a random distribution (as is discussed above in section 1), while the evident lack of correlation between cumulative CO₂ and non-CO₂-fe emissions to peak warming undermines the use of an 'effective' (multi-gas) TCRE. Since CO₂ and non-CO₂ emissions are affected by different policies, it is potentially misleading to present them using a single index such as percentage aggregate CO₂-equivalent emission reductions by a given date⁴⁶. A two-dimensional presentation, separating CO₂ and non-CO₂ contributions to warming (as in panels 3b, c and d here), is the minimum required to ensure both indicators are on track to achieve a temperature goal. The sum of cumulative CO₂ and non-CO₂ CO₂-fe emissions, multiplied by the TCRE, determines long-term warming.

The scenarios explored here do not represent a random distribution that can be sampled for a particular percentile, so accounting for non-CO₂ contributions to remaining CO₂-only

budget estimates is more challenging. Subtracting a representative mid-range scenario's non-CO₂ contribution from estimated total CO₂-fe budgets gives indicative pure-CO₂ budgets, indicated by horizontal box-whisker symbols in pink on Fig. 3 (using the central non-CO₂ RF scenario highlighted in Fig. 1). The range of non-CO₂ contributions implied by the non-CO₂ forcing scenarios as a whole indicates the potential for trade-offs between CO₂ and non-CO₂ warming.

Using GMST warming of 1.23 °C in 2020 and the mid-range non-CO₂ forcing from Fig. 1 (130 GtCO₂-fe), we find, for a 33, 50 and 66% chance of limiting warming to below 1.5 °C, indicative CO₂-only budgets of 640, 545, 455 GtCO₂ respectively. We re-emphasise this is just one possible pathway for future non-CO₂ forcing, which will be determined by policy choices, some but not all of which also involve trade-offs and synergies with CO₂ policy. Exploring these trade-offs is a matter for policymakers. CO₂-fe emissions, or warming-equivalent emissions³⁰ that are very similar and easier to calculate (see Methods), provide the necessary framework. SR1.5 gave 33rd, 50th and 66th percentile remaining carbon budgets for 1.5 °C GMST warming from 2018 of 1080, 770, 570 GtCO₂, respectively. Our best-estimate remaining carbon budget for 1.5 °C from 2020, 545 GtCO₂, is therefore consistent with SR1.5 having accounted for recent updates to the level of GMST and an additional 2 years of warming. SR1.5's GSAT best-estimate remaining carbon budget from 2018 is also consistent, albeit slightly more conservative (50th percentile remaining GSAT budget to 1.5 °C is 580 GtCO₂ from 2018) confirming that the infilled GMST products approximately correspond to GSAT remaining CO₂ budgets from the present-day. Our range of estimated remaining CO₂-only budgets are given in Table 1, along with comparable budgets from SR1.5.

The definition of present-day anthropogenic warming plays a key role in determining the size of remaining budgets, and therefore in determining the appropriate categories for scenarios. However, re-baselining anthropogenic warming to be consistent with the assessment that 0.85 °C warming occurred up to the 2006–2015 decade relative to 1850–1900 (a statement which is used in the Structured Expert Dialogue to inform the Paris Agreement text¹⁴, and cited explicitly in discussions of 1.5 and 2.0 °C in the Paris Agreement by those familiar with the process⁴⁷) results in a best-estimate remaining CO₂ budget from 2020 of 920 GtCO₂. Figure 3d shows that, then, scenarios currently classified as 'lower-2 °C' are consistent with a peak warming of ~1.5 °C. While revised estimates of present-day GMST used in Fig. 3b mean remaining carbon budgets are consistent with those presented in SR1.5, other interpretations of temperature levels referred to in the Paris Agreement, possibly more consistent with the consensus on the current level of warming at the time the Agreement was signed, may result in significantly larger remaining budgets.

Figure 3b, c, d together show that the current classification of the IAM scenarios are much more consistent with peak warming defined by the increase in statistically infilled (i.e. GSAT-like) GMST datasets relative to pre-industrial levels. Defining temperature this way also restores consistency with most other studies of remaining budgets (see Supplementary Tables 1, 2 of ref. ²), at the expense of consistency with reported present-day levels of warming associated with recent impacts¹⁴, although more closely aligning them with model-based studies of future impacts using GSAT¹⁸.

To complement the assessment of TCRES using observations of the climate system, the TCRES for a subset of CMIP6 GCMs can be calculated over history directly using CO₂-fe emissions to account for the impact of non-CO₂ pollutants. Leach et al. (2020) demonstrate the FaIRv2.0⁴⁸ model can emulate the thermal and carbon cycle properties of CMIP6 GCMs, and provide fitted parameters for several CMIP6 models. RFMIP experiments⁴⁹ allow for the calculation of ERF time series over history (UKESM1-0-LL, NorESM2-LM, GFDL-ESM4) and extended up to 2100 where data is




Fig. 4 TCREs calculated for a range of CMIP6 models. FaIRv2.0 simple climate model used to emulate the carbon cycle and thermal responses of each GCM with parameters from Leach et al. (2021); forced with ERF time series diagnosed from ERF experiments completed as part of RFMIP. UKESM1-0-LL and GFDL-ESM4 calculated GCM diagnosed aerosol ERF time series over 1850–2014; NorESM2-LM, IPSL-CM6A-LR and CanESM5 calculated using transient anthropogenic ERFs from RFMIP runs. Solid lines show individual CMIP6 model TCREs calculated with CO₂-fe over historical experiments (coloured by model), while dashed lines show CMIP6 models if the effective TCRE is plotted (1850–2014) from Lindicoat et al. (2021). Brown plume shows the CMIP6 TCRE range calculated with a 1%/yr concentration increase experiment (from Arora et al., 2020). Black shading shows the AR5 gaussian TCRE range, and black lines shows the observationally constrained TCRE range from Fig. 3.

available (CanESM5, IPSL-CM6A-LR). Using these, we diagnose the TCREs from estimates of the cumulative CO₂-fe emissions budgets and associated warming for each GCM, plotted in Fig. 4 (coloured solid lines; the full description in SI). If total warming is instead plotted against CO₂ emissions alone, nonlinearities are introduced (see Fig. 4, Supplementary Fig. 11, and dotted lines in Fig. 1d). For these five models, we find a CMIP6 TCRE range of 0.35–0.68 °C/TtCO₂, consistent with the range of observationally constrained GSMT TCRE from Fig. 3a (black lines in Fig. 4). CMIP6 ensemble members display TCREs which lie on average above the estimated 50th percentile of the observationally constrained distribution. Even UKESM1-0-LL and CanESM5, both of which have equilibrium climate sensitivities that are above the range consistent with historical observations^{13,50} show high but not out-of-range TCRE estimates, lying around the 83rd percentile of the GSMT TCRE distribution. These estimated TCREs can be compared with TCREs calculated using the 1%/yr CO₂ concentration increase experiment (brown plume in Fig. 4), and calculated in ref.⁴³. CMIP6 TCREs estimated with CO₂-fe are near-identical to the estimates in Arora et al. (2020) using pure-CO₂, indicating the CO₂-fe methodology is indeed identifying the same TCRE parameter as in the 1%/yr idealised experiment. We present all calculated TCREs in supplementary table 2. The CMIP6 ensemble range appears to be consistent with the 5th–95th percentile TCRE found with historical observations, with extremes of the ensemble slightly under-sampling the observationally constrained upper and lower bounds. The mean response (0.48 °C/TtCO₂ in CMIP6 models assessed) is also somewhat higher than the observationally constrained result (0.40 °C/TtCO₂). Simply using the CMIP6 range in isolation as an uncertainty interval, therefore, is potentially problematic, despite the range implying no significant bias. A better approach would use observations to constrain the CMIP6 ensemble TCRE range, such as the yes/no exclusion for models

based on historical temperature gradient reconstruction as suggested in Tokarska et al.²⁴.

DISCUSSION

Here we analyse IAM mitigation scenarios informing the IPCC's SR1.5 report, deconstructing them to highlight the relative contributions from CO₂ and non-CO₂ pollutants (Figs. 1 and 2). CO₂-fe emissions provide a means to quantify non-CO₂ contributions to future scenarios, without relying on traditional metrics which do not translate readily into a corresponding warming response. Further, we demonstrate that a simple scaling factor, or 'effective TCRE', doesn't adequately account for the warming contribution from non-CO₂ drivers^{3,32,33} as future non-CO₂ radiative forcing isn't tightly correlated with cumulative CO₂ emissions in these scenarios. We also use CO₂-fe emissions to constrain the TCRE distribution based on historical temperature observations and radiative forcing estimates, and produce an observationally constrained remaining carbon budget estimate based on central non-CO₂ RF estimates, along with highlighting the covarying uncertainty in the physical climate response to CO₂ and non-CO₂ pollutants. We recommend that a two-dimensional presentation, which separates CO₂ and non-CO₂ contributions to warming (Fig. 3b–d) is the most transparent approach when displaying the physical constraints of remaining budgets.

A remaining carbon budget for 1.5 °C of 420 GtCO₂ from 2018 (the most widely quoted 66th percentile SR1.5 GSAT budget) is consistent with a current level of warming of 1.23 °C (in 2020 relative to 1850–1900)¹⁶, given an observationally constrained TCRE (0.40 °C/TtCO₂; 0.26–0.78 °C/TtCO₂), unless we experience a sudden increase in the TCRE or future non-CO₂ climate forcing above the upper end of the range in the SR1.5 1.5 °C-compatible scenarios (see Fig. 3b and Table 1; although recent updates to datasets of observed CH₄ and N₂O mixing ratios suggest these are tracking a path higher than most future scenarios suggest^{51,52}).

Re-baselining the current level of warming to 0.85 °C in the decade prior to 2015, a figure (based solely on GMST) that was used to contextualise the observed impacts of climate change in the Structured Expert Dialogue used to inform the Paris Agreement^{14,17}, is inconsistent with SR1.5's remaining budget estimates, and means that scenarios conventionally referred to as 'Lower-2.0 °C-consistent' in the IIASA SR1.5 scenario database are in fact 1.5 °C-consistent (panel d). A similar misclassification occurs if CO₂ and non-CO₂ CO₂-fe budgets are considered up to 2100 across the SR1.5 scenarios (panel 3c).

The decision on what index will be used to determine when 1.5 °C is reached has substantial policy implications and hence should not be determined by scientists alone. As long as observed warming continues to be reported in terms of GMST, continuing to report remaining carbon budgets in terms of GMST baselined to several periods (for example, both 0.85 °C over the decade prior to 2015; 1850–1900 pre-industrial baseline) seems the simplest and least policy-prescriptive option available. Regardless, we propose CO₂-fe emissions are the most transparent method to analyse the relative contributions from individual pollutants to remaining warming, particularly in order to disentangle scenario from physical climate uncertainty.

METHODS

Calculating CO₂ forcing-equivalent emissions

CO₂-fe emissions time series are computed with a four-pool carbon cycle model^{9,25,48} based closely on the IR model used for metrics calculations in AR5^{8,53}, but with a minor modification to allow state-dependent time-scales: for these ambitious mitigation scenarios, very similar results are obtained using the AR5 formula itself (see Supplementary Fig. 3). The similarity of the dotted and solid lines in Fig. 1c shows that, over these

scenarios and timescales, a 1 W/m^2 change in ERF is approximately equivalent to $1000 \text{ GtCO}_2\text{-fe}$, consistent with Fig. 8.29 of ref. ⁴⁰.

To calculate $\text{CO}_2\text{-fe}$ emissions we calculate the CO_2 concentration associated with a given pollutant's RF scenario:

$$C_{\text{ref}+\Delta F}(t) = C_0 \exp\left(\frac{\ln(2)(F_{\text{ref}}(t)+\Delta F(t))}{F_{2x}}\right), \quad C_{\text{ref}}(t) = C_0 \exp\left(\frac{\ln(2)F_{\text{ref}}(t)}{F_{2x}}\right), \quad (1)$$

where $F_{\text{ref}}(t)$ is the reference scenario's RF (i.e. the forcing due to all-pollutants other than the one we are considering), and $\Delta F(t)$ are the RF of the given pollutant. F_{2x} is the RF for successive doublings in CO_2 concentration.

Following this, we calculate the CO_2 emissions compatible with each concentration pathway $C_{\text{ref}+\Delta F}(t)$ and $C_{\text{ref}}(t)$:

$$E_{\text{ref}+\Delta F/\text{ref}}(t) = C_{\text{ref}+\Delta F/\text{ref}}(t) - C_0 - \sum_{i=0}^3 R_i(t-1)g_i; \\ \text{with } R_i(t) = a_i E(t) + R_i(t-1)g_i; \\ \text{and } g_i = \exp\left(\frac{-1}{a(t)\tau_i}\right), \quad (2)$$

$$\Delta E(t) = E_{\text{ref}+\Delta F}(t) - E_{\text{ref}}(t), \quad (3)$$

where $E_{\text{ref}+\Delta F}(t)$ and $E_{\text{ref}}(t)$ are the annual emissions at time t resulting in concentrations $C_{\text{ref}+\Delta F}(t)$ and $C_{\text{ref}}(t)$. C_0 is the pre-industrial CO_2 concentration, a_i and τ_i are coefficients defined in the AR5 IR model and $a(t)$ is a scaling factor on response timescales to allow for a changing airborne fraction over time, as detailed in ref. ³. Finally, we find the $\text{CO}_2\text{-fe}$ emissions attributable to pollutants with RF time series $\Delta F(t)$ by differencing the calculated annual emissions from the all forcing case, and the annual emissions for the case where we remove the RF from the pollutant: $(\text{ref} + \Delta F) - (\text{ref})$.

Figures 1, 2, 3 and 4 all rely on $\text{CO}_2\text{-fe}$ emissions calculated as above by inverting the carbon cycle of FaIR. A differencing approach is taken when calculating individual pollutant $\text{CO}_2\text{-fe}$ emissions time series, as this is suggested in Jenkins et al. (2018) to best account for nonlinearities in the carbon cycle response to under high RF perturbations²⁵.

For ambitious mitigation scenarios, we can linearise the system to this calculation further. Now, a non- CO_2 ERF time series $\Delta F(t)$ due to a particular climate forcing agent is converted to a perturbation CO_2 concentration, accounting for the nonlinearity in CO_2 forcing, as follows:

$$\Delta C(t) = C_0 \exp\left(\frac{\ln(2)(F_{\text{ref}}(t))}{F_{2x}}\right) \left(\exp\left(\frac{\ln(2)(\Delta F(t))}{F_{2x}}\right) - 1 \right) \approx \frac{\Delta F(t)}{RE} \quad (4)$$

where $F_{\text{ref}}(t)$ is the forcing due to all other agents, F_{2x} is the forcing due to a CO_2 doubling and C_0 is the pre-industrial CO_2 concentration, and RE is the radiative efficiency of CO_2 given present-day concentrations. The $\text{CO}_2\text{-fe}$ emission time series is then found using the following iterative formula:

$$\Delta E(t) = \Delta C(t) - \sum_{i=0}^3 R_i(t-1)g_i; \\ R_i(t) = a_i \Delta E(t) + R_i(t-1)g_i; \quad g_i = \exp\left(\frac{-1}{a(t)\tau_i}\right) \quad (5)$$

For ambitious mitigation scenarios, the linearization $a(t) = 1$ and constant RE provides a very close approximation to the full carbon cycle inversion. Setting $a(t) = 1$ in this way reproduces the linear IR model used in metric calculations of Chapter 8, AR5⁴⁰.

Approximating the $\text{CO}_2\text{-fe}$ calculation

The calculation of $\text{CO}_2\text{-fe}$ emissions in Figs. 1 and 2 does not require a specific model. In the main text, we invert the carbon cycle in the FaIRv1.0 simple climate model to calculate compatible $\text{CO}_2\text{-fe}$ emissions with a given forcing input. In Supplementary Figs. 3 and 4 we reproduce these figures by calculating the $\text{CO}_2\text{-fe}$ emissions with the AR5 IR model². This model is also employed for metric calculations in IPCC's AR5. The AR5 model is linear, meaning it can be inverted as a matrix to calculate $\text{CO}_2\text{-fe}$ emissions⁵⁴.

Converting all climate forcing agents to $\text{CO}_2\text{-fe}$ emissions provides the most accurate and physically justified definition of an 'all-pollutants CO_2 budget' but requires full forcing histories and an invertible carbon cycle model. On decade-to-century timescales, however, $\text{CO}_2\text{-fe}$ emissions associated with any individual forcing agent may be approximated by 'warming equivalent' emissions, $\text{CO}_2\text{-we}$, a linear combination of the components RF level and trend over a recent time interval, Δt .

First, considering how CO_2 emissions alone impact the global temperature anomaly. For CO_2 the change in radiative forcing, ΔF_{CO_2} ,

due to CO_2 emissions over an interval Δt depends on both the cumulative CO_2 emissions over that period, G , and the average- CO_2 induced forcing, \bar{F}_{CO_2} :

$$\Delta F_{\text{CO}_2} = \beta G - \rho \bar{F}_{\text{CO}_2} \Delta t \quad (6)$$

where the coefficient β is the additional radiative forcing per tonne CO_2 emitted (around 1 W/m^2 per 1000 GtCO_2), and ρ is the rate at CO_2 -induced RF declines after CO_2 emissions reach zero. We can rearrange this formula to consider how to treat non- CO_2 radiative forcing contributions in terms of a cumulative $\text{CO}_2\text{-we}$ budget, G_* :

$$G_* = (\Delta F + \rho \bar{F} \Delta t) / \beta \quad (7)$$

Therefore, to calculate the total human-induced warming associated with a CO_2 emissions time series, $E(t)$, and a non- CO_2 forcing time series, $F(t)$, over a time period Δt we can write:

$$\Delta T = \kappa(G + G_*) = \kappa \left(G + \frac{\Delta F}{\beta} + \frac{\rho \bar{F} \Delta t}{\beta} \right) \quad (8)$$

where the first term in brackets are the cumulative CO_2 emissions, the second is the CO_2 budget associated with a change in non- CO_2 RF ΔF , and the third is the CO_2 emissions budget associated with the average global energy imbalance due to non- CO_2 sources, \bar{F} . κ is the TCRE. Values of β and ρ can be related back to the AGWP _{CO_2} ; typical values are $\beta = 1 \text{ W/m}^2$ per 1000 GtCO_2 , and $\rho = 0.3\%$ per year. The AR5 likely range for TCRE is $0.45 \pm 0.23^\circ\text{C}$ per 1000 GtCO_2 (and in Fig. 3 we estimate a TCRE likely range between $0.26\text{--}0.78^\circ\text{C/TtCO}_2$) which far outweighs uncertainty in other coefficients. The third term is relatively small in almost all scenarios but is retained to emphasise that halting warming due to non- CO_2 climate drivers requires declining, not constant, non- CO_2 forcing, although the required rate of decline is small: of the order of $1\text{--}3\%$ per decade. For comparison of this formula with the full $\text{CO}_2\text{-fe}$ calculation see Supplementary Fig. 2.

These options, along with simplifications such as GWP*, mean there are a number of alternatives to plotting the less physically representative GWP₁₀₀ emissions time series for non- CO_2 pollutants, even if in a given setting the full $\text{CO}_2\text{-fe}$ calculation isn't appropriate.

Observationally constrained TCRE estimate

We estimate the TCRE in Fig. 3 using an observationally constrained methodology. We first estimate the present-day temperature anomaly, using an optimal fingerprinting approach outlined in Haustein et al. (2017)³⁸, and based on methodologies in refs. ^{55,56}.

The approach uses ordinary least-squares (OLS) regression onto observed GMST to find the residual-minimising combination of anthropogenic and natural temperature responses. For GMST we use a four-dataset-mean of HadCRUT5 (statistically infilled)⁵⁷, NOAA⁵⁸, Berkeley⁵⁹ and GISTEMP⁶⁰, similar to Chapter 1, SR15¹⁷. The optimal fingerprinting methodology additionally samples uncertainty from internal variability (by sampling detrended CMIP6 PIControl experiment GSAT time series (104 members)⁶¹); along with a range of physical climate response parameters in the FaIR model when deriving temperature response shapes (18 members); uncertainty in the observed temperature anomaly using a 200-member ensemble of observational uncertainty from HadCRUT5. The derived present-day warming level (1.23°C ($1.06\text{--}1.42^\circ\text{C}$) in 2020; 1.12°C ($0.90\text{--}1.33^\circ\text{C}$) average over decade 2010–2019) is in agreement with other recent assessments of the present-day warming level (e.g. Gillett et al., 2021).

With this derived warming level, we now must determine the corresponding all-pollutant cumulative $\text{CO}_2\text{-fe}$ emissions. We do this by inverting the FaIRv1.0 carbon cycle as derived above, with carbon cycle response parameters determined by best-estimate fits the historical relationship between carbon emissions and concentrations (from Jenkins et al., 2018), using historical temperature observations to inform likely historical carbon cycle temperature feedback behaviour. A 1000-member all-pollutant total anthropogenic ERF ensemble from ref. ³⁵ is used to calculate a 1000-member ensemble of cumulative $\text{CO}_2\text{-fe}$ emissions over history, which we plot against the decadal warming levels to produce Fig. 3a. Ellipses are drawn around the central 90% of the underlying decade scatterpoint distributions, using a similar approach to that used in Fig. 1 in ref. ⁶². TCRE estimates are found based on the decade (2011–2020). The observationally constrained TCRE distribution has a log-normal shape and is plotted in Supplementary Fig. 6.

Defining remaining budgets to 1.5 °C

To define the remaining CO₂ budgets from present-day we must remove from the total CO₂-fe budget the contribution from non-CO₂ sources of warming. Using the highlighted scenarios in Fig. 1c we find compatible non-CO₂ contributions of between 50–300 GtCO₂ to peak warming in 1.5 °C-compatible scenarios in the IIASA SR15 database, with a central estimate of 130 GtCO₂.

The observationally constrained range of TCRES found in Fig. 3 of the main text is 0.26–0.78 °C/TtCO₂ (central 90 percent of distribution; 0.95–2.86 °C/TtC) with a best-estimate value of 0.40 °C/TtCO₂ (1.46 °C/TtC). Using the methodology of Haustein et al. (2017)³⁸ we find an estimated 2020 anthropogenic temperature anomaly = 1.23 °C, (5th–95th percentile = 1.06–1.42 °C) (see Supplementary Fig. 5d).

Therefore, the range of total CO₂-fe remaining budgets to 1.5 °C assuming best-estimate warming to date is:

Max: $(1.5 - 1.23) \times 1000 / 0.26 = 1040$ GtCO₂

Min: $(1.5 - 1.23) \times 1000 / 0.78 = 350$ GtCO₂

Best-estimate: $(1.5 - 1.23) \times 1000 / 0.40 = 680$ GtCO₂

Assuming a central estimate of the 1.5 °C-compatible non-CO₂ budget remaining from 2020 to peak warming (130 GtCO₂-fe in a central light blue scenario from Fig. 1, range 50–300 GtCO₂-fe) we find the remaining carbon budget to 1.5 °C corresponding to different TCRES percentiles as follows:

5th percentile: $(0.27 \times 1000 / 0.26) - 130 = 1040 - 130 = 910$ GtCO₂
(range: 740–990 GtCO₂)

33rd percentile: $(0.27 \times 1000 / 0.35) - 130 = 770 - 130 = 640$ GtCO₂
(range: 470–720 GtCO₂)

50th percentile: $(0.27 \times 1000 / 0.40) - 130 = 680 - 130 = 550$ GtCO₂
(range: 380–630 GtCO₂)

66th percentile: $(0.27 \times 1000 / 0.46) - 130 = 590 - 130 = 460$ GtCO₂
(range: 290–540 GtCO₂)

95th percentile: $(0.27 \times 1000 / 0.78) - 130 = 350 - 130 = 220$ GtCO₂
(range: 50–300 GtCO₂)

This methodology is followed to find remaining CO₂-fe budgets in Table 1 and to find CO₂-only budgets at the 33rd, 50th and 66th percentile in the main text and for pink error bars in Fig. 3b, d.

CO₂-fe breakdown of an example 1.5 °C-compatible scenario

Figure 2 uses the same best-estimate historical ERF from ref. ³⁵ which is the basis for the observationally constrained TCRES in Fig. 3. Over the remainder of the 21st century (after 2020) we use the central non-CO₂ forcing scenario from Fig. 1c (highlighted in light blue). Components of ERF are rescaled to match 2020 ERF in the historical ERF time series. Aerosol ERF is consistent with -0.9 W/m^2 in 2011 (the value reported in Chapter 8, AR5; refs. ^{42,63}).

Component CO₂-fe budgets to peak warming are reported in the main text. Component 2100 budgets are: CO₂: 140 GtCO₂; methane: -455 GtCO₂-fe; nitrous oxide: 90 GtCO₂-fe; aerosols: 500 GtCO₂-fe; other forcings: -220 GtCO₂-fe; giving a total CO₂-fe budget of 55 GtCO₂-fe. Figure 2 reports a slightly different non-CO₂ cumulative CO₂-fe budget to peak warming than is used in Fig. 3. This is a result of scalings which are used in Fig. 2 to match IAM output to historical ERF estimates (the difference is equivalent to around a year of warming).

CMIP6 TCRES in Fig. 4

Figure 4 compares CMIP6 model TCRES to the observationally constrained distribution from Fig. 3. We use Arora et al. (2020)'s TCRES estimates using 1%/yr experiments, and compare to TCRES derived with a full CO₂-fe methodology where available historical ERF data is available from RFMIP⁴⁹. Parameters derived using FaIRv2.0 to reproduce carbon cycle behaviour in individual CMIP6 models are given in Leach et al.⁴⁸. We use these to estimate the TCRES for the five models [CanESM5, UKESM1-0-LL, IPSL-CM6A-LR, NorESM2-LM, GFDL-ESM4] where ERF data is available. These TCRES (reproduced for individual models in Supplementary Table 2) agree well with 1%/yr concentration increase TCRES estimates in Arora et al. (2020). Historical CO₂ emissions from Lindicoat et al. (2020) are also used to show CO₂-only effective TCRES (dashed lines).

DATA AVAILABILITY

All data required to reproduce the figures in this paper are available freely online, and relevant databases and repositories are referenced in the text.

CODE AVAILABILITY

All codes required to reproduce the figures in this paper are available from the corresponding author.

Received: 19 January 2021; Accepted: 13 September 2021;

Published online: 14 October 2021

REFERENCES

- IPCC. *Summary for Policymakers of the Special Report on the Global Warming of 1.5 °C* (IPCC, 2018).
- Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J. & Séférián, R. Estimating and tracking the remaining carbon budget for stringent climate targets. *Nature* **571**, 335 (2019).
- Leach, N. J. et al. Current level and rate of warming determine emissions budgets under ambitious mitigation. *Nat. Geosci.* **11**, 574 (2018).
- Millar, R. J. et al. Emission budgets and pathways consistent with limiting warming to 1.5 °C. *Nat. Geosci.* **10**, 741–747 (2017).
- Matthews, H. D., Gillett, N. P., Stott, P. A. & Zickfeld, K. The proportionality of global warming to cumulative carbon emissions. *Nature* **459**, 829–832 (2009).
- Quere, C. L. et al. Global Carbon Budget 2018. *Earth Syst. Sci. Data* **10**, 2141–2194 (2018).
- Huppmann, D. et al. *IAMC 1.5 °C Scenario Explorer and Data hosted by IIASA*. (Integrated Assessment Modeling Consortium & International Institute for Applied Systems Analysis, 2018). <https://doi.org/10.22022/SR15/08-2018.15429>
- Joos, F. et al. Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.* **13**, 2793–2825 (2013).
- Millar, R. J., Nicholls, Z. R., Friedlingstein, P. & Allen, M. R. A modified impulse-response representation of the global near-surface air temperature and atmospheric concentration response to carbon dioxide emissions. *Atmos. Chem. Phys.* **17**, 7213–7228 (2017).
- Ehlert, D. & Zickfeld, K. What determines the warming commitment after cessation of CO₂ emissions? *Environ. Res. Lett.* **12**, 015002 (2017).
- Allen, M. R. et al. Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature* **458**, 1163–1166 (2009).
- Matthews, H. D., Zickfeld, K., Knutti, R. & Allen, M. R. Focus on cumulative emissions, global carbon budgets and the implications for climate mitigation targets. *Environ. Res. Lett.* **13**, 010201 (2018).
- WMO. *WMO Provisional Statement on the State of the Global Climate in 2019* (WMO, 2019).
- UNFCCC. *Report on the Structured Expert Dialogue on the 2013–2015 Review* (UNFCCC, 2015).
- Tokarska, K. B. et al. Recommended temperature metrics for carbon budget estimates, model evaluation and climate policy. *Nat. Geosci.* **12**, 964–971 (2019).
- Gillett, N. P. et al. Constraining human contributions to observed warming since the pre-industrial period. *Nat. Clim. Change* **11**, 1–6 (2021).
- Allen, M. R. et al. Framing and Context. In: V. Masson-Delmott et al. (eds) *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. (2018).
- Collins, M. et al. *Long-term Climate Change: Projections, Commitments and Irreversibility* (IPCC AR5, 2013).
- Matthews, H. D. & Zickfeld, K. Climate response to zeroed emissions of greenhouse gases and aerosols. *Nat. Clim. Change* **2**, 338–341 (2012).
- MacDougall, A. H. et al. Is there warming in the pipeline? A multi-model analysis of the zero emissions commitment from CO₂. *Biogeosciences* **17**, 2987–3016 (2020).
- Jones, C. D. et al. The zero emissions commitment model intercomparison project (ZECMIP) contribution to C4MIP: quantifying committed climate changes following zero carbon emissions. *Geosci. Model Dev.* **12**, 4375–4385 (2019).
- IPCC. *AR5 Synthesis Report: Climate Change 2014*. (IPCC, 2013). <https://www.ipcc.ch/report/ar5/syr/>.
- Huppmann, D., Rogelj, J., Kriegler, E., Krey, V. & Riahi, K. A new scenario resource for integrated 1.5 °C research. *Nat. Clim. Change* **8**, 1027–1030 (2018).
- Tokarska, K. B. et al. Uncertainty in carbon budget estimates due to internal climate variability. *Environ. Res. Lett.* **15**, 104064 (2020).
- Jenkins, S., Millar, R. J., Leach, N. & Allen, M. R. Framing climate goals in terms of cumulative CO₂-forcing-equivalent emissions. *Geophys. Res. Lett.* **45**, 2795–2804 (2018).

26. Mengis, N., Partanen, A.-I., Jalbert, J. & Matthews, H. D. 1.5 °C carbon budget dependent on carbon cycle uncertainty and future non-CO₂ forcing. *Sci. Rep.* **8**, 1–7 (2018).
27. Wigley, T. M. L. The Kyoto Protocol: CO₂ CH₄ and climate implications. *Geophys. Res. Lett.* **25**, 2285–2288 (1998).
28. Forster, P. D. et al. *IPCC Special Report on the Global Warming of 1.5°C* (IPCC, 2018).
29. Allen, M. R. et al. A solution to the misrepresentations of CO₂ -equivalent emissions of short-lived climate pollutants under ambitious mitigation. *Npj Clim. Atmos. Sci.* **1**, 16 (2018).
30. Cain, M. et al. Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *Npj Clim. Atmos. Sci.* **2**, 1–7 (2019).
31. Lynch, J., Cain, M., Pierrehumbert, R. & Allen, M. Demonstrating GWP_{ast}: a means of reporting warming-equivalent emissions that captures the contrasting impacts of short- and long-lived climate pollutants. *Environ. Res. Lett.* **15**, 044023 (2020).
32. Matthews, H. D. et al. Estimating carbon budgets for ambitious climate targets. *Curr. Clim. Change Rep.* **3**, 69–77 (2017).
33. Millar, R. J. & Friedlingstein, P. The utility of the historical record for assessing the transient climate response to cumulative emissions. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* **376**, 20160449 (2018).
34. Mengis, N. & Matthews, H. D. Non-CO₂ forcing changes will likely decrease the remaining carbon budget for 1.5 °C. *Npj Clim. Atmos. Sci.* **3**, 1–7 (2020).
35. Dessler, A. E. & Forster, P. M. An estimate of equilibrium climate sensitivity from interannual variability. *J. Geophys. Res. Atmos.* **123**, 8634–8645 (2018).
36. Allen, M., Jenkins, S., Sha, F. & Macey, A. Defining carbon neutrality, climate neutrality and net zero emissions. *Climate Policy* (2021, In Review).
37. Gillett, N. P., Arora, V. K., Matthews, D. & Allen, M. R. Constraining the ratio of global warming to cumulative CO₂ emissions using CMIP5 simulations. *J. Clim.* **26**, 6844–6858 (2013).
38. Hausteine, K. et al. A real-time global warming index. *Sci. Rep.* **7**, 15417 (2017).
39. Geoffroy, O. et al. Transient climate response in a two-layer energy-balance model. Part I: analytical solution and parameter calibration using CMIP5 AOGCM experiments. *J. Clim.* **26**, 1841–1857 (2012).
40. Myhre, G. et al. Anthropogenic and Natural Radiative Forcing. In: Stocker, T. F. et al. (eds) *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2013).
41. Friedlingstein, P. et al. Global carbon budget 2019. *Earth Syst. Sci. Data* **11**, 1783–1838 (2019).
42. Bellouin, N. et al. Bounding global aerosol radiative forcing of climate change. *Rev. Geophys.* **58**, e2019RG000660 (2020).
43. Arora, V. K. et al. Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models. *Biogeosciences* **17**, 4173–4222 (2020).
44. Williams, R. G., Ceppi, P. & Katavouta, A. Controls of the transient climate response to emissions by physical feedbacks, heat uptake and carbon cycling. *Environ. Res. Lett.* **15**, 0940c1 (2020).
45. Damon Matthews, H. et al. An integrated approach to quantifying uncertainties in the remaining carbon budget. *Commun. Earth Environ.* **2**, 1–11 (2021).
46. United Nations Environment Programme (2019). Emissions Gap Report 2019. *UNEP, Nairobi*.
47. Mace, M. J. Mitigation commitments under the Paris Agreement and the Way Forward. *Clim. Law* **6**, 21–39 (2016).
48. Leach, N. J. et al. FalRv2.0.0: a generalised impulse-response model for climate uncertainty and future scenario exploration. *Geosci. Model Dev.* **14**, 1–29 (2020).
49. Pincus, R., Forster, P. M. & Stevens, B. The radiative forcing model intercomparison project (RFMIP): experimental protocol for CMIP6. *Geosci. Model Dev.* **9**, 3447–3460 (2016).
50. Tsutsui, J. Diagnosing transient response to CO₂ forcing in coupled atmosphere-ocean model experiments using a climate model emulator. *Geophys. Res. Lett.* **47**, e2019GL085844 (2020).
51. Tian, H. et al. A comprehensive quantification of global nitrous oxide sources and sinks. *Nature* **586**, 248–256 (2020).
52. Ganesan, A. L. et al. Advancing scientific understanding of the global methane budget in support of the Paris Agreement. *Glob. Biogeochem. Cycles* **33**, 1475–1512 (2019).
53. Held, I. M. et al. Probing the fast and slow components of global warming by returning abruptly to preindustrial forcing. *J. Clim.* **23**, 2418–2427 (2010).
54. Smith, M. A., Cain, M. & Allen, M. R. Further improvement of warming-equivalent emissions calculation. *Npj Clim. Atmos. Sci.* **4**, 1–3 (2021).
55. Hegerl, G. C. et al. Detecting greenhouse-gas-induced climate change with an optimal fingerprint method. *J. Clim.* **9**, 2281–2306 (1996).
56. Hasselmann, K. Multi-pattern fingerprint method for detection and attribution of climate change. *Clim. Dyn.* **13**, 601–611 (1997).
57. Morice, C. P. et al. An updated assessment of near-surface temperature change from 1850: the HadCRUT5 data set. *J. Geophys. Res. Atmos.* **126**, e2019JD032361 (2021).
58. Smith, T. M., Reynolds, R. W., Peterson, T. C. & Lawrimore, J. Improvements to NOAA's historical merged land–ocean surface temperature analysis (1880–2006). *J. Clim.* **21**, 2283–2296 (2008).
59. Rohde, R. A. & Hausfather, Z. The Berkeley Earth land/ocean temperature record. *Earth Syst. Sci. Data* **12**, 3469–3479 (2020).
60. Lenssen, N. J. L. et al. Improvements in the GISTEMP uncertainty model. *J. Geophys. Res. Atmospheres* **124**, 6307–6326 (2019).
61. Eyring, V. et al. Overview of the coupled model intercomparison project phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* **9**, 1937–1958 (2016).
62. Otto, A. et al. Energy budget constraints on climate response. *Nat. Geosci.* **6**, 415–416 (2013).
63. Stevens, B. Rethinking the lower bound on aerosol radiative forcing. *J. Clim.* **28**, 4794–4819 (2015).

ACKNOWLEDGEMENTS

S.J., M.R.A., T.W. and P.F. acknowledge funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 821003 (4C project). S.J. is supported by the NERC Doctoral Training Partnership NE/L002612/1. The authors thank Chris Smith for contributing transient ERF time series calculated as part of RFMIP experiments for use in the production of Fig. 4 and Spencer Lindicoat for providing CMIP6 historical CO₂ emissions time series in Fig. 4.

AUTHOR CONTRIBUTIONS

S.J. and M.R.A. designed the study. M.C. and S.J. designed the simple formula approximating CO₂-fe emissions in Methods. P.F. contributed CMIP6 model diagnosed CO₂ emissions and temperatures from 1%/yr runs for TCRE estimates and NG provided CMIP6 GSAT and GMST time series for comparisons of budgets in Fig. 3. T.W. and S.J. completed analysis of CMIP6 historical runs to estimate GCM TCRES in Fig. 4. All authors contributed to writing.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41612-021-00203-9>.

Correspondence and requests for materials should be addressed to Stuart Jenkins.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

Quantifying non-CO2 contributions to remaining carbon budgets

Jenkins, Stuart

2021-10-14

Attribution 4.0 International

Jenkins S, Cain M, Friedlingstein P, et al., (2021) Quantifying non-CO2 contributions to remaining carbon budgets, npj Climate and Atmospheric Science, Volume 4, October 2021, Article number 47

<https://doi.org/10.1038/s41612-021-00203-9>

Downloaded from CERES Research Repository, Cranfield University