1 Climate Change

2 Carbon losses in the Alps 3

Soil carbon stocks depend on inputs from decomposing vegetation and returns to the atmosphere as CO₂. Monitoring of carbon stocks in German alpine soils has shown large losses linked to climate change and a possible positive feedback loop.

8 Guy Kirk

4

5

6

7

9 10 The response of the terrestrial carbon cycle to global change is one of the main uncertainties in current 11 climate change predictions¹. Most terrestrial carbon is held in soils as organic matter derived from the 12 decay of plants. Soil organic matter accounts for roughly three times more carbon than living 13 vegetation, and for more carbon than vegetation and the atmosphere combined. Because elevated 14 atmospheric CO_2 concentrations have a fertilizing effect on plant growth, anthropogenic CO_2 15 emissions have triggered increases in the land carbon sink². However, models predict that other 16 factors – such as water and nutrients – will eventually become limiting to plant growth, and hence to 17 the land carbon sink. In contrast, the turnover of soil organic matter producing CO_2 is expected to 18 increase as the Earth warms. As a result, simulations using coupled carbon-climate models predict 19 that the land surface will become a net source of CO_2 before the end of the century, leading to a 20 feedback loop between climate and soil carbon losses: increased emissions of CO₂ from soil organic 21 matter will lead to enhanced warming, which may then feedback to further soil organic matter losses. 22 Prietzel and colleagues³, writing in *Nature Geoscience*, provide evidence that warming has already 23 caused a decline in soil organic matter in the German Alps. 24

25 The umbrella term "soil carbon" covers a diverse array of molecules, which are stabilised to varying 26 extents by binding to soil minerals and by physical protection within soil aggregates⁴. Its turnover is 27 mediated by soil microbes, with half-times ranging from weeks to millennia. Physical protection 28 within aggregates can render soil carbon inaccessible to microbes and the enzymes they produce to 29 cleave complex carbon molecules into easily digestible compounds. Like other metabolic and 30 enzymatic processes, rates of turnover are sensitive to temperature and moisture. Increasing 31 temperatures associated with climate change are expected to increase soil carbon turnover rates¹, but 32 direct evidence that this is currently taking place is limited. Since high elevation ecosystems are 33 expected to warm more quickly than other parts of the planet, with associated changes in snow cover 34 and freeze-thaw cycles, they are a good place to determine whether warming is already causing soil 35 carbon losses.

36 37 Prietzel and colleagues³ measured losses of carbon from forest soils in the German Alps over the past 38 30 years, which they link to higher summer temperatures. They did this with repeated sampling of 39 soils at 35 sites across both forests and pastures. They found no change in soil carbon in the pasture 40 areas, but large losses in absolute and percentage terms in soils with high carbon stocks in lower 41 elevation forests (< 1100 m). Changes in management weren't responsible for the carbon losses, 42 because forests across the German Alps have been under constant, low intensity management for 43 more than 50 years. Instead, the losses matched changes in mean summer temperatures at the low 44 elevation sites: temperatures at these elevations increased by 0.5 °C per decade over the survey period,

45 even though mean winter temperatures at these elevations decreased by 0.3 °C per decade.

46

47 Measuring climate-induced changes in soil carbon is difficult. Measurable changes occur on decadal 48 time scales, and it can be challenging to detect the changes against large background concentrations 49 and the tremendous spatial variability inherent in soils. Further, the effects of climate must be 49 disentangled from the effects of past changes in land use or management, which may persist for many 49 decades. It is therefore crucial to have long-term soil monitoring schemes that are properly designed, 50 managed and resourced to account for these factors, such as the scheme used by Prietzel et al. 53

Various repeated soil inventories have been made in different parts of the world over the past two or
 three decades, and some have shown losses of carbon but others gains. In most cases the changes

- could be attributed to changes in land use and management. For example, large losses of carbon from soils across England and Wales during the 1980s and 90s⁵ were largely due to land use changes, but up to 10 % could have been due to climate change⁶. The losses of the order of 100 g C m⁻² yr⁻¹
 reported by Prietzel and colleagues are somewhat greater than those found in England and Wales⁵ for forest soils with similar carbon contents but at mostly lower elevations.
- 61

62 To understand how the sources and sinks of greenhouse gases will change with the climate, it is 63 essential to have reliable mathematical models. But simulating soil carbon processes is a particularly 64 thorny challenge. In addition to the inherent complexity of soil carbon and its interactions with soil 65 physical and chemical conditions, a number of other factors complicate the picture. The availability of 66 readily decomposed substrates, such as root secretions, may fuel the turnover of otherwise more-67 stable components in so-called priming effects. Soil microbes may adapt to warmer temperatures, and 68 thereby reduce or in some cases increase the magnitude of soil carbon responses to climate change⁷. 69 Further, the carbon cycle is inextricably linked to other nutrient cycles, and changes in these other 70 cycles, such as changes in atmospheric nitrogen deposition, can have cascading effects on plant productivity and soil carbon dynamics⁸. Lastly, extreme weather events, such as drought or storms, 71 72 can alter the turnover rates of terrestrial carbon pools leading to prolonged release of CO₂ to the 73 atmosphere⁹.

74

Models must be under-pinned by field studies, and ultimately by long-term soil monitoring at regional and national scales, of which the study by Prietzel and colleagues³ is exemplary. Their evidence that climate change has already started depleting soil carbon in the German Alps raises the possibility that a positive feedback between climate and ecosystems is beginning. Combined with the expected decline in the fertilization effect from elevated atmospheric CO₂ concentrations, soils could become

80 important contributors to accelerating climate change.

81

82 *Guy Kirk is in the School of Water, Energy and Environment, Cranfield University, Cranfield,*83 *Bedford MK43 0AL, UK.*

84 *e-mail* g.kirk@cranfield.ac.uk85

86 References

- Ciais, P., *et al.* in *Climate Change 2013: The Physical Science Basis* (ed. Stocker, T. F. *et al.*)
 465–570 (Cambridge Univ. Press, 2013).
- 89 2. Zhu, Z., et al. Nature Clim. Change doi:10.1038/nclimate3004 (2016).
- 90 3. Prietzel, J., Zimmermann, L., Schubert, A. & Christophel, D. Nature Geosci. ... (2016)
- 91 4. Schmidt, M. W. *et al. Nature* **478**, 49–56 (2011)
- 92 5. Bellamy, P. H., Loveland, P. J., Bradley, R. I., Lark, R. M. & Kirk, G. J. D. *Nature* 437, 245–248
 93 (2005).
- 94 6. Kirk, G. J. D. & Bellamy, P. H. Eur. J. Soil Sci. 61, 406–411 (2010).
- 95 7. Karhu, K. et al. Nature 513, 81–84 (2014).
- 96 8. Fernandez-Martinez, M. et al. Nature Clim. Change 4, 471-476 (2014).
- 97 9. Reichstein, M. et al. Nature 500, 287–295 (2013).
- 98 99
- 100
- 101
- 102
- 103 104



Figure 1 | Soil carbon dynamics. Plant productivity-and plant carbon stocks-can be stimulated by increased atmospheric CO₂ concentrations. That plant carbon is then decomposed and converted to soil carbon, where it can remain immobilized for weeks to thousands of years before being mineralized to CO₂ and released back to the atmosphere. However, warming can stimulate the decomposition and mineralization of soil carbon, leading to increased emissions of CO₂ to the atmosphere. Prietzel et al.³ used repeated soil measurements in the German Alps over the last 30 years to show that soil carbon has decreased in managed forests concomitantly with an increase in temperatures.

CERES Research Repository

School of Water, Energy and Environment (SWEE)

Staff publications (SWEE)

Climate change: carbon losses in the Alps

Kirk, Guy J. D.

2016-06-13 Attribution-NonCommercial 4.0 International

Guy Kirk. Climate change: carbon losses in the Alps. Nature Geoscience Volume 9, pp478-479 (2016) http://dspace.lib.cranfield.ac.uk/handle/1826/10070 Downloaded from CERES Research Repository, Cranfield University