

Erratum

Alternaria in Food: Ecophysiology, Mycotoxin Production and Toxicology

Hyang Burm Lee^{1*}, Andrea Patriarca² and Naresh Magan³

¹Environmental Microbiology Lab., Division of Food Technology, Biotechnology & Agrochemistry, College of Agriculture & Life Sciences, Chonnam National University, Gwangju 500-757, Korea

²Departamento de Química Orgánica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Pabellón II, 3o Piso, 1428 Buenos Aires, Argentina

³Applied Mycology Group, AgriFood Theme, Cranfield University, Cranfield, Bedford, MK43 0AL, UK

Volume 43(2): 93-106, 2015

Table 1 contained errors in direction of arrowheads.

Table 1. Summary of the range of environmental conditions which allow germination, growth and toxin production (based on data from several studies) by *Alternaria alternata* and *A. tenuissima* (psychrotolerant)

Factors	Germination	Growth	Toxin production
Temperature (°C)	1~35	< 1 and > 35	< 10 and > 35
Water activity (a _w)	0.84~0.995	< 0.85	< 0.90
pH	2.5~10	< 2.5 and > 10	< 2.5 and > 9

The corrected Table 1 follows below (direction of arrowheads).

Table 1. Summary of the range of environmental conditions which allow germination, growth and toxin production (based on data from several studies) by *Alternaria alternata* and *A. tenuissima* (psychrotolerant)

Factors	Germination	Growth	Toxin production
Temperature (°C)	1~35	> 1 and < 35	> 10 and < 35
Water activity (a _w)	0.84~0.995	> 0.85	> 0.90
pH	2.5~10	> 2.5 and < 10	> 2.5 and < 9