

## Detecting Recovery of the Stratospheric Ozone Layer

Martyn P. Chipperfield<sup>1,2</sup>, Slimane Bekki<sup>3</sup>, Sandip Dhomse<sup>1</sup>, Neil R.P. Harris<sup>4</sup>, Birgit Hassler<sup>5</sup>,  
Ryan Hossaini<sup>6</sup>, Wolfgang Steinbrecht<sup>7</sup>, Rémi Thiéblemont<sup>3</sup> and Mark Weber<sup>8</sup>

- 5 1. School of Earth and Environment, University of Leeds, UK.  
2. National Centre for Earth Observation, University of Leeds, U.K.  
3. LATMOS-IPSL, Paris, France.  
4. Centre for Atmospheric Informatics and Emissions Technology, Cranfield University, UK.  
5. Bodeker Scientific, New Zealand.  
10 6. Lancaster Environment Centre, Lancaster University, UK.  
7. Deutscher Wetterdienst, Hohenpeissenberg, Germany.  
8. Institute of Environmental Physics, University of Bremen, Germany.

Revised version 2 July 28, 2017

15

**As a result of the 1987 Montreal Protocol and its amendments, the atmospheric loading of anthropogenic ozone-depleting substances is decreasing. Accordingly, recovery of the stratospheric ozone layer is expected. However, short data records and atmospheric variability confound the search for early signs of recovery. Moreover, climate change is masking ozone recovery from ozone-depleting substances in some regions and will increasingly affect the extent of recovery. Here we discuss the nature and timescales of ozone recovery, and explore the extent to which it can be currently detected in different atmospheric regions.**

20

Depletion of the stratospheric ozone layer has been one of the most prominent environmental issues of the past 40 years. The layer prevents biologically damaging solar ultraviolet (UV) radiation (wavelengths below about 300 nm) from reaching the surface and was thus a key factor in creating the conditions for the evolution of life on earth<sup>1</sup>. Ozone is also an efficient absorber of terrestrial infra-red radiation and therefore a 'greenhouse' gas, albeit one that is produced in the atmosphere rather than emitted at the surface. In recent years it has become clear that there is a strong coupling between stratospheric ozone and climate change.

25

Serious concern over ozone depletion started in the 1970s when it was realised that the break-down of man-made compounds such as chlorofluorocarbons (CFCs) in the middle stratosphere would release chlorine atoms which could catalytically destroy ozone<sup>2,3</sup>. Research activity increased greatly following the discovery in 1985 of large and unexpected depletion in the Antarctic lower stratosphere during spring<sup>4</sup>, the so-called Antarctic ozone hole. This depletion was caused by increasing levels of chlorine and bromine in the atmosphere but, crucially, also involved the conversion of stable chlorine reservoir species into active ozone-destroying forms on the surface of polar stratospheric clouds which form in winter and spring<sup>5</sup>. Atmospheric scientists failed to predict the ozone hole in advance as the models used to forecast the evolution of the ozone layer did not include such processes<sup>6</sup>.

35

By the time of the discovery of the Antarctic ozone hole the process for international protection of ozone layer had already been initiated and the framework for its implementation

45 set up by the signing of the Vienna Convention in 1985. The Montreal Protocol on  
Substances which Deplete the Ozone Layer was signed in September 1987 and ratified by  
January 1989. The protocol, with its 30<sup>th</sup> anniversary this year, was a major achievement in  
terms of global environmental protection although it initially placed only modest limits on the  
production and consumption of major ozone-depleting substances (ODSs) such as CFCs  
and bromine-containing halons. Importantly, the protocol allowed for further strengthening  
50 through later amendments and adjustments, and over time these revisions have led to an  
almost complete ban on major classes of ODSs including CFCs, replacement  
hydrochlorofluorocarbons (HCFCs) and related compounds such as methyl chloroform and  
carbon tetrachloride. These compounds have atmospheric lifetimes of the order of 10-100  
years<sup>7</sup>, and so the response of the atmospheric chlorine and bromine loading to changes in  
emissions is slow. Nevertheless, observations show<sup>5</sup> that the abundance of these gases in  
55 the lower atmosphere is largely responding to the Montreal Protocol limits as expected and  
most major ODSs are decreasing (**Figure 1**). The sum of tropospheric chlorine peaked in  
1993 and the sum of tropospheric bromine peaked a few years later in 1997. The  
stratospheric abundance of chlorine (which largely resides as HCl in the upper  
60 stratosphere<sup>8</sup>) and bromine, derived from these ODSs, followed these tropospheric  
variations but with a delay of ~3 to 7 years (depending on the region) due to the slow  
timescale for transport and degradation of ODSs through the stratosphere<sup>9</sup>. Accordingly, we  
expect stratospheric ozone depletion due to chlorine and bromine to follow this behaviour of  
increasing depletion through the late 1990s, followed by a turn-around and slow 'recovery'.  
65 Unfortunately, variability in other factors affecting ozone, such as stratospheric dynamics  
(i.e. wind, temperature), aerosol loading, and solar irradiance, all complicate this simple  
picture and mask the small signal of ozone recovery from ODSs – expected to be around a  
few percent per decade globally. An important question is therefore to what degree and  
where can we detect ozone recovery. A further important question is the ultimate extent to  
70 which the ozone layer will recover, given the increasing impact of climate change on  
atmospheric structure and composition. In this perspective, we analyse what observations  
and models can already tell us about the current status of recovery is and where ozone is  
heading. We show that ozone recovery is proceeding consistent with our understanding and  
argue that it should not be seen as a single detectable event, but rather a slow direction of  
75 travel for which the evidence will become gradually clearer as our data records increase.

### Recovery of the Ozone Layer

Although the concept of an 'ozone layer on the mend' may appear simple, discussions on  
this topic within the atmospheric science community are often heated and become very  
80 nuanced. So, why is detecting the recovery of the stratospheric ozone layer a tricky subject?  
Can't we just define recovery as when ozone levels return to their former values before  
significant ozone depletion occurred? At first that appears conceptually trivial, but  
complications immediately arise when we consider that the rest of the atmosphere is also  
changing. The continuing rise in CO<sub>2</sub> is altering the physical structure of the atmosphere: the  
85 tropopause is rising so the stratosphere is getting thinner; its temperature structure is  
changing, and the Brewer-Dobson Circulation in which air is transported into and through the  
stratosphere (**Figure 1**) may accelerate in the future<sup>10</sup>. As a result, there may be stronger  
upwelling in the tropics and faster descent over the middle latitudes and polar regions.  
Because ozone abundance increases with altitude in the lower stratosphere (where most of

90 the ozone resides), this would lead to less ozone in the tropics and more ozone at higher  
latitudes. Cooling in the upper stratosphere is already increasing ozone in that region by  
slowing gas-phase ozone destruction cycles<sup>11,12</sup>. All these changes are driven fundamentally  
by increasing greenhouse gases (GHGs), especially CO<sub>2</sub>. At the same time, N<sub>2</sub>O and CH<sub>4</sub>  
95 are also currently increasing, with CH<sub>4</sub> in particular very sensitive to variations in emissions  
from its natural and anthropogenic sources<sup>13</sup>. The balance between the various catalytic  
cycles which result from ODS, CH<sub>4</sub> and N<sub>2</sub>O degradation and drive ozone loss will therefore  
change as well. What is clear is that the chemistry and dynamics of the stratosphere will  
have changed sufficiently that recovery to ozone levels prior to ozone depletion is not a  
100 sensible concept. The picture is complicated further when we consider how recovery occurs  
in different parts of the atmosphere or in different seasons.

If the atmosphere is not returning to its former state, then should we not look to changes in  
surface UVB radiation (or even the consequences) as the best measure for recovery? After  
all, the motivation for the Montreal Protocol was to avoid the risk of health and other impacts  
of increased UVB. However, possible changes in non-stratospheric factors such as  
105 tropospheric ozone, clouds, aerosols and the earth's albedo, coupled with their large  
variability, all make it hard to draw meaningful conclusions about recovery of surface UVB<sup>14</sup>.  
It is a more complex picture even than for ozone.

From a regulatory point of view, as control of production and consumption is the tool that  
policy-makers can employ, the success of the Montreal Protocol is judged primarily by the  
110 changes in atmospheric ODS concentrations. From this perspective, the Montreal Protocol is  
already undoubtedly a success; ODS levels are decreasing<sup>15</sup> with expected benefits for  
ozone and UVB radiation and also for climate. However, the impact of these ODS declines  
on ozone levels has proved much harder to detect.

Definitions of ozone recovery have tended to be based on the concept of a state or stage  
115 being reached. Since recovery is often defined with respect to the effect of ODSs (the key  
factor in the Montreal Protocol), each stage requires a clear attribution of ozone changes to  
the decline and ultimately return of ODSs to their pre-industrial levels<sup>16</sup>. The following stages  
(or fingerprints) of recovery have been defined<sup>16</sup>: 1) a significant slowing down of  
stratospheric ozone decline; then 2) the onset of a significant increase; and finally 3) the full  
120 recovery of ozone from ODSs, when ozone is no longer significantly affected by them.  
However, it is more beneficial to think of recovery as the direction of travel rather than the  
destination. Indeed, full recovery does not necessarily imply a return of stratospheric ozone  
to pre-1980 levels because the influence of other factors, notably increasing GHG levels, is  
growing. The ODS levels in the atmosphere are clearly decreasing (**Figure 1**) and the first  
125 stage (or 'fingerprint') of the ozone response, the end of the ozone decline, has been  
observed<sup>5,17</sup>. However, it has been difficult to establish the occurrence of the next stage, i.e.  
a general upward trend in ozone due to declining ODSs. This may be surprising, since ODS  
levels have now been in decline for 15 to 20 years. However, due to the long atmospheric  
lifetimes of ODSs (typically many decades<sup>7</sup>), this decline is about three times slower than  
130 their rapid increase before the Montreal Protocol came into effect. Back then, it took 10 to 15  
years to clearly detect the significant decrease in global ozone. Everything else being equal,  
we might expect 30 to 40 years before detecting a significant upward trend in global ozone  
due to declining ODS levels becomes possible<sup>18</sup>.

### 135 **Diagnosing Ozone Recovery from Current Observations**

Regular stratospheric ozone observations started with ground-based Dobson spectrophotometers in the mid-1920s<sup>19,20</sup>. Continuous measurements from space started in 1978 with the Solar Backscatter UltraViolet (SBUV) instrument<sup>21</sup> and so global ozone observations now span a time period of nearly forty years (see **Figure 2**). This includes  
140 about 20 years of observations after the global stratospheric ODS peak around 1997 (or 16 years after the later ODS peak in polar regions<sup>9</sup>). The length of the observable recovery period thus covers about two decades which is short to identify uniquely ODS-related recovery among other sources of variability that operate on multi annual or decadal timescales. However, given the importance of this topic there have been many studies which  
145 have investigated observationally based ozone trends over this period<sup>22–28</sup>.

Satellite and ground-based data revealed a dramatic decline in the total ozone column (i.e. the total number of ozone molecules in the whole depth of the atmosphere, per unit area) of about 3%/decade until the mid-1990s, caused by ODS increases (**Figure 2**). In the NH, the lowest annual mean total ozone columns occurred in 1992, resulting from enhanced ozone  
150 destruction linked to heterogeneous chemistry on volcanic aerosols and transport changes<sup>29–31</sup>, after the major volcanic eruption of Mt Pinatubo in 1991, a few years before the peak in stratospheric ODS. In the late 1990s, annual mean total ozone increased rapidly, faster than expected from the slow decrease in ODS. This is related to variability in atmospheric dynamics, notably for ozone transport, exemplified by the Brewer-Dobson  
155 circulation<sup>32,33</sup>. All long observational time series show such decadal variability (e.g. the record at Arosa Switzerland since 1925<sup>34</sup>).

Since 2000, stratospheric ODS levels have been decreasing slowly (at only 1/3 of the rate of the previous ODS increase), and extratropical total ozone has levelled off. **Figure 2** also shows results of a multiple linear regression (MLR, see **Supplementary Information**) with  
160 independent linear trends before and after the ODS peak in 1996 (2000 for the polar regions<sup>9</sup>) applied to annual or September mean total ozone. The MLR approach is used here to linearly decompose the ozone variability into components of a long-term trend and shorter term variability. The regression accounts for well-established sources of shorter term ozone variability (solar variations, stratospheric aerosols, variations in the Brewer-Dobson  
165 circulation, modes of climate variability such as the quasi-biennial oscillation (QBO) and El Nino Southern Oscillation). In the tropics and at mid-latitudes, linear trends after 1997 are generally small and not statistically significant. In the tropics, observed total ozone has not changed much at all over the entire time period since 1979. The only region with a possibly significant positive trend in total ozone in the last decade is the Antarctic in September (the  
170 month when the ozone hole reaches its maximum areal extent), see **Figure 2** or results from<sup>35–38</sup>; however, results are sufficiently sensitive to uncertainties in the MLR and to the inclusion or not of a specific year in the time series, that formal identification of Antarctic ozone recovery remains uncertain (see<sup>37</sup> and **Figure S1** in **Supplementary Information**). Antarctic October time series shows no significant trend, nor does the Arctic in February and  
175 March (not shown).

The interannual variability of ozone is largest in the lower stratosphere (which dominates the behaviour of the total column). It is much smaller in the upper stratosphere where the end of the ozone decline (the first sign of ozone recovery) was first detected<sup>17</sup>. Vertically resolved ozone measurements (from multiple satellite and ground-based instruments) have now  
180 reported a possible signal of the next stage of recovery in the upper stratosphere, with ozone increases of 2-4% /decade<sup>5,39</sup>, although the statistical significance of this recovery remains

unclear<sup>39</sup>. **Figure 3** shows the corresponding updated ozone time series at 40 km (2 hPa). Nearly all the different observational datasets closely follow the inverted curve of stratospheric chlorine loading. However, model results show that both ODS and greenhouse gases have contributed equally to the recent upper stratospheric increase<sup>5</sup>.

Key issues in the detection of ozone increases from observations are the availability of several independent long-term records (to quantify uncertainty in the data) and the uncertainties of the trend estimates. Optimally, the total uncertainty should include any uncertainties based on: 1) the measurement technique<sup>40</sup>, 2) data sampling<sup>41,42</sup>, 3) the uncertainties introduced by the regression, 4) uncertainties from data preparation, especially merging data from different satellites to create a full time series, and 5) differences of trends from different observation systems. While the first two sources of uncertainty are relatively well understood<sup>41</sup>, the last three have only started to be addressed in recent years. Consistent multi-decadal, vertically resolved ozone measurements from a single instrument are available only from a few sparse ground-based stations<sup>43-45</sup>. Satellite measurements do provide global coverage, but no single satellite instrument spans the several decades necessary for a robust analysis of both the ozone decline and the expected subsequent increase. Therefore, measurements from different satellite instruments are being combined<sup>46-51</sup> but individual satellite instrument characteristics (offsets and drifts) are difficult to determine<sup>52-55</sup>. At this point robust records with comprehensive uncertainties have not been achieved and a full uncertainty calculation for vertically resolved ozone trends is still missing (see<sup>39,5</sup>).

### Diagnosing Ozone Recovery from Atmospheric Models

Attribution of ozone trends to specific factors (e.g. evolution of ODS levels) requires the use of atmospheric chemical-dynamical models. These models, although they are by no means perfect and often show significant differences compared to observations and other models, encapsulate our best understanding of the fundamental physics and chemistry that control ozone and its variations. Those most suited for diagnosing ozone recovery are chemical transport models (CTMs) and chemistry-climate models (CCMs). Both include relevant internal and external drivers, especially changing concentrations of ODSs, variations in solar forcing, effects of volcanic eruptions, and changing surface conditions. CTMs are well suited to comparisons with specific observations because dynamics (wind and temperature fields) are prescribed from meteorological re-analyses such as ERA-Interim<sup>56</sup> or MERRA<sup>57</sup>, thus ignoring feedbacks of chemistry on temperature and dynamics via radiatively driven heating. CCMs normally calculate their own “random” realizations of meteorology, including the feedbacks of changing trace gases on temperature and transport, although they can be nudged to follow prescribed meteorology and therefore perform like a CTM<sup>36</sup>.

In contrast to observations, models allow us to compare various scenarios with different assumptions of factors affecting ozone. Using different model runs, in combination with observed time series, allows the attribution of ozone changes and thus the diagnosis of ozone recovery without relying on an MLR (as in **Figure 2**).

We can use CTM simulations to quantify the expected ozone change in different regions due to separate forcings in the past, notably ODS changes. **Figure 4** presents such results from the TOMCAT CTM<sup>58</sup> which quantify ozone depletion since 1960, when anthropogenic ODS levels were very low, and since 1980, near the start of the global ozone record (see

**Supplementary Information**). The past accumulating emissions of ODSs between 1960 and 1980 contribute more to ozone depletion than the emissions after 1980 (for which the signal only appears in the stratosphere a few years later), illustrating the difference in taking these two baselines as a reference. In any case, the model-predicted signal of recovery from ODS (light green shaded region following peak ODSs in 1995) has clearly only reached a small fraction of this past depletion by 2015. Also shown in **Figure 4** is the estimated impact of dynamical variability, mainly through transport, on ozone (orange line). This impact is larger in the lower stratosphere and thus it is important for the total column, since ozone density is generally largest at the higher pressures of the lower stratosphere. The resulting year-to-year variations are clearly larger than the predicted signal in recovery.

Shepherd et al.<sup>59</sup> compared column ozone observations with simulations from a CCM nudged with ERA-Interim reanalyses. The difference between a run with 1980 ODS levels and a run with realistically changing ODS indicates that global ozone columns should already have benefited from declining ODS levels. The model results in **Figure 4** confirm the findings of Shepherd et al. that by 2010 ODS-related midlatitude column ozone loss had already declined by 10% since the peak ODS loading in the late 1990s. Shepherd et al. also noted that tropospheric ozone increases may have compensated for ozone depletion in the tropical lower stratosphere, explaining the lack of long-term column trend there (**Figure 2**).

As the largest impact of ODSs occurs in the springtime Antarctic lower stratosphere<sup>4</sup>, it is reasonable to suppose that this may be the region with the earliest and/or clearest signs of ozone recovery<sup>17,60,61</sup>. Solomon et al.<sup>36</sup> analysed Antarctic September column ozone observations through 2014 and found signs of recovery since 2000, at 90% confidence. They did not include the year 2002, with unusual polar vortex behaviour and small ozone loss<sup>62-64</sup>, in their analysis (see also **Figure 2d**). Their nudged CCM simulations indicate that about one half of the ozone increase observed in September over the period 2000 to 2014 could be attributed to declining ODS. They attributed the other 50% of the increase to transport changes and the very large ozone holes of 2011 and 2015 to additional chemical losses triggered by aerosol enhancements from relatively small volcanic eruptions in Chile<sup>65</sup>. However, considerable uncertainties pertain to the simulated effects of transport and aerosol changes. It cannot be ruled out that both have contributed a larger fraction to the observed ozone increase from 2000 to 2014, implying a small impact of decreasing ODS over the period.

Separating small ozone changes due to slowly decreasing ODS (less than a few percent per decade) from large dynamical (transport and temperature) related variations is difficult. This is shown in **Figure S2** in the **Supplementary Information**, which compares the different TOMCAT simulations shown in **Figure 4**. All time series show large similar inter-annual variations, up to  $\pm 10$  DU. These are mostly due to meteorological variability, and are not seen with repeating 1980 meteorological conditions every year. The recovery signal due to declining ODS does not exceed +6 DU at mid-latitudes, which is much smaller than the large year-to-year variations. The largest modelled recovery signal is in the Antarctic in September at over 20 DU. **Figure S2** also shows that the model captures the large Arctic ozone depletion observed in March 2011<sup>66,67</sup>. This large depletion was caused by exceptionally persistent cold stratospheric conditions and is consistent with our understanding of stratospheric ozone and is within the range of variability expected in the Arctic. Importantly, it does not, therefore, undermine our expectation of long-term ozone recovery as ODS decline. A similar argument applies to the large Antarctic depletion of 2015<sup>65</sup>.

275 In the upper stratosphere ozone concentrations are increased by GHG-induced cooling. The  
increases of observed ozone in the upper stratosphere at northern mid-latitudes after 2000  
discussed above<sup>5,39</sup> are of the same magnitude as those calculated by CCM simulations<sup>68</sup>,  
giving some confidence in both observations and simulations. Comparison of these different  
simulations shows that declining ODS and stratospheric cooling have contributed about  
equally to the observed increase of ozone in the upper stratosphere since 2000; neither  
factor alone is sufficient to explain the observations. WMO/UNEP<sup>5</sup> therefore concluded that  
280 declining ODS play a role, and that the signal of ozone recovery from ODS is seen in the  
upper stratosphere.

The TOMCAT CTM simulations in **Figure 4** (and **Figure S2**) clearly reiterate that the main  
obstacle to detecting and estimating the small ozone recovery from ODS for the 2000-2015  
period is the large year-to-year variations, mostly of dynamical origin. The most common  
285 approach to account for this high frequency variability in trend analysis is the MLR, although  
some of its assumptions may be questioned. In order to test the linear decomposition of  
ozone variability into independent contributions from different factors and increase the level  
of confidence in the MLR results, we carry out a trend analysis of observational and model  
time series. The external factors considered in the MLR (solar, aerosol, detrended heat  
290 fluxes (used a proxy for the variations in the Brewer Dobson Circulation), quasi-biennial  
oscillation (QBO), see also **Figure 2**) mostly generate high frequency (e.g. year-to-year)  
variability in ozone. The long-term trend in ozone should be overwhelmingly caused by the  
decline in ODS and the rise in GHGs. Since the MLR cannot discriminate between these  
effects, it contains only one trend term.

295 The values of the 2000-2015 ozone trends derived from a simple linear regression (i.e.  
regression where only a linear trend is fit) and MLR are generally in good agreement for both  
observations and the model simulations for the tropics, mid-latitudes (NH, SH), and Antarctic  
(**Figure 5**). In addition, the MLR trend values for the observations and the control simulation  
are also consistent within error bars. As expected, the MLR trend representing the effect of  
300 declining ODS ('fixed dynamics' simulation) is clearly positive in all regions, and most  
pronounced in the Antarctic. In contrast, the MLR trend attributed to dynamical variability  
('fixed ODS' simulation) varies strongly from one region to another. For total ozone, the trend  
is negative at mid-latitudes but null in Antarctica; in the upper stratosphere, it is positive in  
the tropics and at northern mid-latitudes but negative at southern mid-latitudes. As  
305 mentioned before, low frequency natural dynamical variability, e.g. in the Brewer-Dobson  
circulation, can randomly generate significant trends over 10-15 year periods<sup>69</sup>. They can  
mask ozone recovery from ODS. Interestingly, the value of the MLR trend for the control  
simulation is approximately equal to the sum of those of the 'fixed dynamics' and 'fixed ODS'  
simulations, suggesting that the effects of changes in ODS and dynamics are approximately  
310 additive and that an attribution analysis based on the simulations is justified<sup>70</sup>. The  
uncertainty in the overall trend is dominated by the dynamical variability and large errors  
bars indicate that the dynamical proxies (QBO, detrended heat flux) in the MLR are not able  
to account for all the interannual variability. Note that as no ozone-dynamics feedbacks are  
included in the 'fixed dynamics' simulation, our estimation of the declining ODS contribution  
315 (inferred from this simulation) is possibly an upper limit, particularly in the upper  
stratosphere<sup>17</sup>.

Taken together, **Figures 4, 5, and S2** demonstrate how small the expected ozone recovery  
currently is compared to the confounding interannual dynamical variations. As expected, the

320 most certain signs for a beginning recovery can now be seen in regions where the ozone  
layer is most sensitive to ODS and the dynamical variability is not too large. These regions  
are the upper stratosphere<sup>5</sup>, and, to a lesser extent, the Antarctic in September<sup>36</sup>, where our  
model results indicate that statistically significant recovery could be detectable with a  
suitable observing system. For other regions, variability is too large (e.g. Arctic, extra-tropical  
325 lower stratosphere), ODS effects are too small (e.g. tropical stratosphere), negative  
dynamical trends mask an ODS-driven positive trend (extratropical column ozone) or the  
observational record is too short or not accurate enough (e.g. tropical lower stratosphere).  
The discrepancies between observed and simulated total ozone (black and dark blue lines in  
**Figure S2**), notably in the NH midlatitudes in the 1980s and in the Antarctic, indicate that  
simulations are subject to additional uncertainties, such as missing processes, uncertain rate  
330 constants and incomplete parametrizations, which pose additional challenges for attributing  
ozone recovery from models.

### **Expectations for Recovery in the 21<sup>st</sup> Century**

335 As ODS levels continue to decline throughout this century, the associated ozone recovery  
signal will become stronger and hence easier to extract, especially with gradually longer data  
records<sup>71</sup>. Although ODS changes have been the main driver of ozone evolution throughout  
most of the stratosphere since the 1970s and will still be important in the future, the influence  
of increasing GHG concentrations (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>) on global ozone has been growing and  
will become dominant in the second-half of the century<sup>5</sup>. Future low frequency variability  
340 (trends) in stratospheric ozone will likely be driven by competing effects (e.g. effects of  
decreasing ODS versus those of increasing GHG) and the complex chemical-dynamical-  
radiative couplings will not be unravelled easily, again making attribution uncertain.

Our best tools for future ozone projections are 3-D chemistry-climate models, which  
encompass all relevant knowledge of key processes, but they need to be evaluated against  
345 observations and require significant computer resources to run a wide range of scenarios  
over century timescales. That work is ongoing<sup>68,72</sup> but here we show some illustrative  
results from a simpler zonal mean latitude-height model (**Figure 6**). Driven by ODS  
concentrations declining throughout this century, global ozone is projected to recover its  
1980 level by ~2030 and its 1960 level by mid-century for a standard scenario for ODS and  
350 GHG emissions (Representative Concentration Pathway (RCP) 6.0, a scenario wherein the  
combined increase of greenhouse gases produces an increase in radiative forcing of 6 W/m<sup>2</sup>  
by 2100). Note that the projected dates of return to specific historical levels, especially 1960,  
can vary greatly from model to model. For instance, some models predict a return of global  
ozone to 1960 level as early as 2030 whereas, according to other models, global ozone  
355 barely recovers the 1960 level by the end of century<sup>68</sup>. Interestingly, the model-projected  
return of global ozone to historical levels is faster than the return of ODS to their natural  
levels which is likely to occur towards the end of the century if current control measures of  
the Montreal Protocol are adhered to in the future. This future accelerated ozone recovery,  
the so-called 'super-recovery', is mainly due to the positive influence of increasing CO<sub>2</sub>,  
360 which cools the upper stratosphere, and therefore reduces the rate of ozone loss, and CH<sub>4</sub>  
which influences chlorine partitioning in the upper stratosphere and is a source of odd-  
hydrogen species which catalytically destroy ozone. In contrast the increase in N<sub>2</sub>O tends to  
decrease ozone<sup>73</sup> through increased NO<sub>x</sub>-catalysed loss. Note that the predicted recovery  
rate for global ozone is strongly sensitive to the assumed future GHG emissions which,



365 unlike ODS emissions, are uncertain, especially in the second half of the century. This, along  
with model uncertainties, makes model-projected ozone recovery extent and associated  
return dates to historical levels uncertain.

The combined influence of these drivers (ODS and GHG changes) is predicted to result in a  
long-term evolution of stratospheric ozone more diverse regionally than would be expected  
370 from ODS decline alone. Ozone recovery is projected to be faster in the NH than in the SH<sup>5</sup>.  
Although the Antarctic is a region where recovery may be starting to be definitively  
detectable, it is also where the return of column ozone to 1960 levels is projected to occur  
relatively late, towards the end of the century, which is about the same time as the return of  
ODS concentrations to their near-natural 1960 levels<sup>68</sup>. Although the signal is more difficult  
375 to detect<sup>74</sup>, the return of Arctic ozone to 1960 levels is projected to occur earlier, around  
2030, which is indicative of the strong influence of rising GHG concentrations<sup>75</sup>. In contrast  
to the rest of the globe, the future evolution of tropical ozone column is expected to be  
essentially driven by changes in GHG levels in the stratosphere and possibly ozone changes  
in the troposphere. Stratospheric ozone projections in the tropics vary widely with the  
380 assumed GHG future emission scenarios of CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>. Models predict that the  
tropical stratospheric partial column ozone may experience either a continued decline (no  
recovery) with column values lower at the end of the century than the present-day due to  
enhanced upwelling, or a super recovery with column values higher than the 1960 levels. A  
future decrease in tropical total (stratosphere + troposphere) column ozone, which models  
385 suggest could be a few percent, could have serious consequences as it would increase  
surface UVB radiation in a region where about 40% of the world population lives and  
population growth is large. Finally, we note that in some regions future ozone recovery might  
be affected in the short-term by sporadic volcanic eruptions or by the occurrence of  
persistently cold Arctic winters such as in 2010/11. They are part of the natural variability  
390 and would not impact the long-term ozone recovery<sup>76</sup>, although there are suggestions that  
this variability may increase due to climate change, for example, cold winters in the Arctic  
stratosphere may be getting colder<sup>77</sup>, though the evidence for this is not conclusive<sup>5</sup>.

### **Outlook for the Ozone Layer**

395 Our best understanding indicates that the measures taken to date through the Montreal  
Protocol have started to remediate ozone depletion and will carry on doing so globally. The  
measurements and the models are imperfect, but studies have used evidence from both to  
show there is a consistent picture, in accord with our current understanding, that ozone  
recovery in the observations is still largely masked by natural variability in most regions but  
400 with clear evidence of it in the upper stratosphere and early signs of it in the Antarctic. We  
can therefore say that we are on track for ozone 'recovery'. However, we emphasise that  
'recovery' is not a single event but a continuing journey; as time passes we will have more  
confidence that it is occurring and a better estimate of its extent. Uncertainty will gradually  
decrease as measurement records become longer and the growing signal emerges from the  
405 underlying natural variability. It is a matter of waiting and ensuring that high quality and  
consistent observations continue. The quadrennial WMO/UNEP ozone assessment, with the  
next one due in 2018, is the forum and the milestone for community-wide scientific updates  
on how far we are on the road to ozone recovery.

410 We need to be vigilant against other factors which may perturb the ozone layer (e.g. extreme  
dynamical events<sup>78,79</sup>, volcanic eruptions<sup>31,65,80</sup>, irregular solar flux variations<sup>81</sup>, uncontrolled  
very short-lived halogenated substances<sup>82–85</sup>, deliberate climate intervention through  
geoengineering<sup>86,87</sup>) to ensure that any observed changes are consistent with our  
415 understanding and do not change our expectation of long-term recovery. Provided ODS  
decline continues in the future as mandated by the Montreal Protocol, we will be moving  
from a stratospheric ozone layer mainly perturbed by anthropogenic chlorine and bromine, to  
one where the impact of increasing greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O)<sup>88</sup> and associated on-  
going climate change become more important and dominate after 2050. Safe-guarding the  
420 ozone layer in the 2<sup>nd</sup> half of this century, therefore, will require continued measurements of  
ozone and ODS as well as of CO<sub>2</sub> and the atmospheric temperature structure, but  
increasingly also measurements of trace gases arising from increased emissions of CH<sub>4</sub> and  
N<sub>2</sub>O. The fundamental underlying processes are understood and are represented  
reasonably well in current atmospheric models. Nevertheless, since more factors with  
425 competing effects are coming into play, further improvements in these computationally  
expensive models (and in the machines on which they run) will be necessary, to understand  
better the measurements, to untangle the more complex interplay of processes controlling  
stratospheric ozone in the future, and to ascertain that ozone has indeed recovered from  
anthropogenic ODS by the end of the century. Future knowledge about ozone recovery will  
430 require the commissioning and launch of new satellite instruments and the continuation of  
the ground-based network. In particular, these instruments must measure at a high enough  
vertical resolution in the lower and mid-stratosphere.

**Acknowledgements:** We are grateful to John Pyle and Paul Newman for helpful comments  
on this work. We thank Anja Schmidt for comments on an early version of the manuscript,  
435 Eric Fleming for providing Figure 6, and Wuhu Feng for help with TOMCAT. The TOMCAT  
modelling work was supported by the NERC National Centre for Atmospheric Science  
(NCAS) and the simulations were performed on the Archer and Leeds ARC/N8 computers.  
MPC acknowledges support of a Royal Society Wolfson Merit Award. MW acknowledges  
440 partial support from the Deutsche Forschungsgemeinschaft (DFG) Research Unit SHARP  
(Stratospheric Change and its Role for Climate Prediction) and the ESA CCI-Ozone project.  
RT acknowledges his funding by the LABEX L-IPSL project (grant ANR-10-LABX-18-01). SB  
and NRPH have been partially supported by the European project StratoClim (603557 under  
programme FP7-ENV.2013.6.1-2). NRPH also acknowledges support from NERC CAST  
(NE/I030051/1). We thank the three anonymous reviewers for their comments which helped  
445 to improve the manuscript significantly.

**Author Contribution:** MPC initiated the study and recruited the author team. All authors  
contributed to the writing of the paper. MPC and SD ran and analysed the TOMCAT  
450 simulations. SB, RT and MW performed the MLR studies. RH, MW, WS, SD, RT and MPC  
produced the figures.

**Author Information:** Requests for information should be sent to Martyn Chipperfield.

## References

455

1. Wayne, R. P. *Chemistry of Atmospheres*. (Oxford University Press, 2000).
2. Molina, M. & Rowland, F. Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone. *Nature* **249**, 810–812 (1974).
3. Stolarski, R. S. & Cicerone, R. J. Stratospheric Chlorine: a Possible Sink for Ozone. *Can. J. Chem.* **52**, 1610–1615 (1974).
4. Farman, J. C., Gardiner, B. G. & Shanklin, J. D. Large losses of total ozone in Antarctica reveal seasonal ClO<sub>x</sub>/NO<sub>x</sub> interaction. *Nature* **315**, 207–210 (1985).
5. World Meteorological Organization (WMO). *Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and Monitoring Project - Report No. 55*. (2014).
6. Solomon, S., Garcia, R. R., Rowland, F. S. & Wuebbles, D. J. On the depletion of Antarctic ozone. *Nature* **321**, 755–758 (1986).
7. SPARC. *Lifetimes of Stratospheric Ozone-Depleting Substances, Their Replacements, and Related Species*. (World Climate Research Programme, WCRP-15/2013, 2013).
8. Michelsen, H. A. *et al.* Stratospheric chlorine partitioning: Constraints from shuttle-borne measurements of [HCl], [ClNO<sub>2</sub>], and [ClO]. *Geophys. Res. Lett.* **23**, 2361–2364 (1996).
9. Newman, P. A., Daniel, J. S., Waugh, D. W. & Nash, E. R. A new formulation of equivalent effective stratospheric chlorine (EESC). *Atmos. Chem. Phys.* **7**, 4537–4552 (2007).
10. Butchart, N. *et al.* Chemistry-climate model simulations of twenty-first century stratospheric climate and circulation changes. *J. Clim.* **23**, 5349–5374 (2010).
11. Haigh, J. D. & Pyle, J. A. Ozone perturbation experiments in a two-dimensional circulation model. *Q. J. R. Meteorol. Soc.* **108**, 551–574 (1982).
12. Jonsson, A. I., Fomichev, V. I. & Shepherd, T. G. The effect of nonlinearity in CO<sub>2</sub> heating rates on the attribution of stratospheric ozone and temperature changes. *Atmos. Chem. Phys.* **9**, 8447–8452 (2009).
13. Nisbet, E. G. *et al.* Rising atmospheric methane : 2007-14 growth and isotopic shift . *Global Biogeochem. Cycles* **30**, 1356–1370 (2016).
14. McKenzie, R. L., Aucamp, P. J., Bais, A. F., Björn, L. O. & Ilyas, M. Changes in biologically-active ultraviolet radiation reaching the Earth's surface. *Photochem. Photobiol. Sci.* **6**, 218–231 (2007).
15. Montzka, S. A. *et al.* Present and future trends in the atmospheric burden of ozone-depleting halogens. *Nature* **398**, 690–694 (1999).
16. World Meteorological Organization (WMO). *Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project - Report No. 50*. (2007).
17. Newchurch, M. J. *et al.* Evidence for slowdown in stratospheric ozone loss: First stage of ozone recovery. *J. Geophys. Res.* **108**, 4507 (2003).
18. Weatherhead, E. C. *et al.* Detecting the recovery of total column ozone. *J. Geophys. Res.* **105**, 22201–22210 (2000).

465

470

475

480

485

490

495

19. Dobson, G. Forty years' research on atmospheric ozone at Oxford: A history. *Appl. Opt.* **7**, 387 (1968).
- 500 20. Staehelin, J. *et al.* Total ozone series at Arosa (Switzerland): Homogenization and data comparison. *J. Geophys. Res.* **103**, 5827–5841 (1998).
21. McPeters, R. D., Bhartia, P. K., Haffner, D., Labow, G. J. & Flynn, L. The version 8.6 SBUV ozone data record: An overview. *J. Geophys. Res.* **118**, 8032–8039 (2013).
- 505 22. Reinsel, G. C. *et al.* Trend analysis of total ozone data for turnaround and dynamical contributions. *J. Geophys. Res.* **110**, D16306 (2005).
23. Zanis, P. *et al.* On the turnaround of stratospheric ozone trends deduced from the reevaluated Umkehr record of Arosa, Switzerland. *J. Geophys. Res.* **111**, D22307 (2006).
- 510 24. Dhomse, S., Weber, M., Wohltmann, I., Rex, M. & Burrows, J. P. On the possible causes of recent increases in northern hemispheric total ozone from a statistical analysis of satellite data from 1979 to 2003. *Atmos. Chem. Phys.* **6**, 1165–1180 (2006).
25. Angell, J. K. & Free, M. Ground-based observations of the slowdown in ozone decline and onset of ozone increase. *J. Geophys. Res.* **114**, D07303 (2009).
- 515 26. Chehade, W., Weber, M. & Burrows, J. P. Total ozone trends and variability during 1979–2012 from merged data sets of various satellites. *Atmos. Chem. Phys.* **14**, 7059–7074 (2014).
27. Nair, P. J. *et al.* Subtropical and midlatitude ozone trends in the stratosphere: Implications for recovery. *J. Geophys. Res.* **120**, 7247–7257 (2015).
- 520 28. Yang, E.-S. *et al.* Attribution of recovery in lower-stratospheric ozone. *J. Geophys. Res.* **111**, D17309 (2006).
29. Poberaj, C., Staehelin, J. & Brunner, D. Missing Stratospheric Ozone Decrease at Southern Hemisphere Middle Latitudes after Mt. Pinatubo: A Dynamical Perspective. *J. Atmos. Sci.* **68**, 1922–1945 (2011).
- 525 30. Aquila, V., Oman, L. D., Stolarski, R., Douglass, A. R. & Newman, P. A. The Response of Ozone and Nitrogen Dioxide to the Eruption of Mt. Pinatubo at Southern and Northern Midlatitudes. *J. Atmos. Sci.* **70**, 894–900 (2013).
31. Dhomse, S. S. *et al.* Revisiting the hemispheric asymmetry in mid-latitude ozone changes following the Mount Pinatubo eruption: A 3-D model study. *Geophys. Res. Lett.* **42**, 3038–3047 (2015).
- 530 32. Fusco, A. C. & Salby, M. L. Interannual Variations of Total Ozone and Their Relationship to Variations of Planetary Wave Activity. *J. Clim.* **12**, 1619–1629 (1999).
33. Weber, M. *et al.* The Brewer-Dobson circulation and total ozone from seasonal to decadal time scales. *Atmos. Chem. Phys.* (2011). doi:10.5194/acp-11-11221-2011
- 535 34. Brönnimann, S., Luterbacher, J., Staehelin, J. & Svendby, T. M. An extreme anomaly in stratospheric ozone over Europe in 1940–1942. *Geophys. Res. Lett.* **31**, L08101 (2004).
35. Salby, M., Titova, E. & Deschamps, L. Rebound of Antarctic ozone. *Geophys. Res. Lett.* **38**, L09702 (2011).
- 540 36. Solomon, S. *et al.* Emergence of healing in the Antarctic ozone layer. *Science* **353**, 269–274 (2016).
37. de Laat, A. T. J., van der A, R. J. & van Weele, M. Tracing the second stage of ozone recovery in the Antarctic ozone-hole with a 'big data' approach to multivariate

- regressions. *Atmos. Chem. Phys.* **15**, 79–97 (2015).
- 545 38. Kuttippurath, J. & Nair, P. J. The signs of Antarctic ozone hole recovery. *Sci. Rep.* **7**, 585 (2017).
39. Harris, N. R. P. *et al.* Past changes in the vertical distribution of ozone – Part 3: Analysis and interpretation of trends. *Atmos. Chem. Phys.* **15**, 9965–9982 (2015).
- 550 40. Millan, L. F. *et al.* Case studies of the impact of orbital sampling on stratospheric trend detection and derivation of tropical vertical velocities: Solar occultation vs. limb emission sounding. *Atmos. Chem. Phys.* **16**, 11521–11534 (2016).
41. Toohey, M. & von Clarmann, T. Climatologies from satellite measurements: the impact of orbital sampling on the standard error of the mean. *Atmos. Meas. Tech.* **6**, 937–948 (2013).
- 555 42. Toohey, M. *et al.* Characterizing sampling biases in the trace gas climatologies of the SPARC Data Initiative. *J. Geophys. Res.* **118**, 11847–11862 (2013).
43. Kirgis, G., Leblanc, T., McDermid, I. S. & Walsh, T. D. Stratospheric ozone interannual variability (1995–2011) as observed by lidar and satellite at Mauna Loa Observatory, HI and Table Mountain Facility, CA. *Atmos. Chem. Phys.* **13**, 5033–5047
- 560 (2013).
44. Nair, P. J. *et al.* Ozone trends derived from the total column and vertical profiles at a northern mid-latitude station. *Atmos. Chem. Phys.* **13**, 10373–10384 (2013).
45. Vigouroux, C. *et al.* Trends of ozone total columns and vertical distribution from FTIR observations at eight NDACC stations around the globe. *Atmos. Chem. Phys.* **15**, 2915–2933 (2015).
- 565 46. Froidevaux, L. *et al.* Global Ozone Chemistry And Related trace gas Data records for the Stratosphere (GOZCARDS): methodology and sample results with a focus on HCl, H<sub>2</sub>O, and O<sub>3</sub>. *Atmos. Chem. Phys.* **15**, 10471–10507 (2015).
47. Davis, S. M. *et al.* The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) database: a long-term database for climate studies. *Earth Syst. Sci. Data* **8**, 461–490 (2016).
- 570 48. Sofieva, V. F. *et al.* Harmonized dataset of ozone profiles from satellite limb and occultation measurements. *Earth Syst. Sci. Data* **5**, 349–363 (2013).
49. Sioris, C. E. *et al.* Trend and variability in ozone in the tropical lower stratosphere over 2.5 solar cycles observed by SAGE II and OSIRIS. *Atmos. Chem. Phys.* **14**, 3479–3496 (2014).
- 575 50. Kyrölä, E. *et al.* Combined SAGE II–GOMOS ozone profile data set for 1984–2011 and trend analysis of the vertical distribution of ozone. *Atmos. Chem. Phys.* **13**, 10645–10658 (2013).
- 580 51. Frith, S. M. *et al.* Recent changes in total column ozone based on the SBUV Version 8.6 Merged Ozone Data Set. *J. Geophys. Res.* **119**, 9735–9751 (2014).
52. Hubert, D. *et al.* Ground-based assessment of the bias and long-term stability of 14 limb and occultation ozone profile data records. *Atmos. Meas. Tech.* **9**, 2497–2534 (2016).
- 585 53. Tegtmeier, S. *et al.* SPARC Data Initiative: A comparison of ozone climatologies from international satellite limb sounders. *J. Geophys. Res.* **118**, 12,229–12,247 (2013).
54. Rahpoe, N. *et al.* Relative drifts and biases between six ozone limb satellite measurements from the last decade. *Atmos. Meas. Tech.* **8**, 4369–4381 (2015).
55. Eckert, E. *et al.* Drift-corrected trends and periodic variations in MIPAS IMK/IAA

- 590 ozone measurements. *Atmos. Chem. Phys.* **14**, 2571–2589 (2014).
56. Dee, D. P. *et al.* The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Q. J. R. Meteorol. Soc.* **137**, 553–597 (2011).
57. Rienecker, M. M. *et al.* MERRA: NASA’s Modern-Era Retrospective Analysis for Research and Applications. *J. Clim.* **24**, 3624–3648 (2011).
- 595 58. Chipperfield, M. P. New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer experiments. *Q. J. R. Meteorol. Soc.* **132**, 1179–1203 (2006).
59. Shepherd, T. G. *et al.* Reconciliation of halogen-induced ozone loss with the total-column ozone record. *Nat. Geosci.* **7**, 443–449 (2014).
- 600 60. Yang, E.-S. *et al.* First stage of Antarctic ozone recovery. *J. Geophys. Res.* **113**, D20308 (2008).
61. Charlton-Perez, A. J. *et al.* The potential to narrow uncertainty in projections of stratospheric ozone over the 21st century. *Atmos. Chem. Phys.* **10**, 9473–9486 (2010).
- 605 62. Simmons, A. *et al.* ECMWF Analyses and Forecasts of Stratospheric Winter Polar Vortex Breakup: September 2002 in the Southern Hemisphere and Related Events. *J. Atmos. Sci.* **62**, 668–689 (2005).
63. Feng, W. *et al.* Three-dimensional model study of the Antarctic ozone hole in 2002 and comparison with 2000. *J. Atmos. Sci.* 822–837 (2005). doi:10.1175/JAS-3335.1
- 610 64. Stolarski, R. S., McPeters, R. D. & Newman, P. A. The Ozone Hole of 2002 as measured by TOMS. *J. Atmos. Sci.* **62**, 716–720 (2005).
65. Ivy, D. J. *et al.* The influence of the Calbuco eruption on the 2015 Antarctic ozone hole in a fully coupled chemistry-climate model. *Geophys. Res. Lett.* **44**, 2556–2561 (2017).
- 615 66. Manney, G. L. *et al.* Unprecedented Arctic ozone loss in 2011. *Nature* **478**, 469–475 (2011).
67. Chipperfield, M. P. *et al.* Quantifying the ozone and ultraviolet benefits already achieved by the Montreal Protocol. *Nat. Commun.* **6**, 7233 (2015).
- 620 68. Eyring, V. *et al.* Multi-model assessment of stratospheric ozone return dates and ozone recovery in CCMVal-2 models. *Atmos. Chem. Phys.* **10**, 9451–9472 (2010).
69. Abalos, M., Legras, B., Ploeger, F. & Randel, W. J. Evaluating the advective Brewer-Dobson circulation in three reanalyses for the period 1979-2012. *J. Geophys. Res.* **120**, 7534–7554 (2015).
- 625 70. World Meteorological Organization (WMO). *Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project - Report No. 52.* (2011).
71. Weatherhead, E. C. & Andersen, S. B. The search for signs of recovery of the ozone layer. *Nature* **441**, 39–45 (2006).
- 630 72. Morgenstern, O. *et al.* Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI). *Geosci. Model Dev.* **10**, 639–671 (2017).
73. Revell, L. E., Bodeker, G. E., Huck, P. E., Williamson, B. E. & Rozanov, E. The sensitivity of stratospheric ozone changes through the 21st century to N<sub>2</sub>O and CH<sub>4</sub>. *Atmos. Chem. Phys.* **12**, 11309–11317 (2012).
- 635 74. Kreher, K., Bodeker, G. E. & Sigmund, M. An objective determination of optimal site locations for detecting expected trends in upper-air temperature and total column

- ozone. *Atmos. Chem. Phys.* **15**, 7653–7665 (2015).
75. Eyring, V. *et al.* Long-term ozone changes and associated climate impacts in CMIP5 simulations. *J. Geophys. Res.* **118**, 5029–5060 (2013).
- 640 76. Bednarz, E. M. *et al.* Future Arctic ozone recovery: the importance of chemistry and dynamics. *Atmos. Chem. Phys.* **16**, 12159–12176 (2016).
77. Rex, M. *et al.* Arctic ozone loss and climate change. *Geophys. Res. Lett.* **31**, L04116 (2004).
78. Newman, P. A., Coy, L., Pawson, S. & Lait, L. R. The anomalous change in the QBO in 2015-16. *Geophys. Res. Lett.* **43**, 8791–8797 (2016).
- 645 79. Osprey, S. M. *et al.* An unexpected disruption of the atmospheric quasi-biennial oscillation. *Science (80-. )*. **353**, 1424–1427 (2016).
80. McCormick, M. P., Thomason, L. W. & Trepte, C. R. Atmospheric effects of the Mt Pinatubo eruption. *Nature* **373**, 399–404 (1995).
- 650 81. Dhomse, S. S. *et al.* On the ambiguous nature of the 11 year solar cycle signal in upper stratospheric ozone. *Geophys. Res. Lett.* **43**, 7241–7249 (2016).
82. Hossaini, R. *et al.* Growth in stratospheric chlorine from short-lived chemicals not controlled by the Montreal Protocol. *Geophys. Res. Lett.* **42**, 4573–4580 (2015).
83. Leedham Elvidge, E. C. *et al.* Increasing concentrations of dichloromethane, CH<sub>2</sub>Cl<sub>2</sub>, inferred from CARIBIC air samples collected 1998–2012. *Atmos. Chem. Phys.* **15**, 1939–1958 (2015).
- 655 84. Sinnhuber, B.-M. & Meul, S. Simulating the impact of emissions of brominated very short lived substances on past stratospheric ozone trends. *Geophys. Res. Lett.* **42**, 2449–2456 (2015).
85. Hossaini, R. *et al.* The increasing threat to stratospheric ozone from dichloromethane. *Nat. Commun.* **8**, 15962 (2017).
- 660 86. Tilmes, S., Garcia, R. R., Kinnison, D. E., Gettelman, A. & Rasch, P. J. Impact of geoengineered aerosols on the troposphere and stratosphere. *J. Geophys. Res.* **114**, D12305 (2009).
87. Nowack, P. J., Abraham, N. L., Braesicke, P. & Pyle, J. A. Stratospheric ozone changes under solar geoengineering : implications for UV exposure and air quality. *Atmos. Chem. Phys.* **16**, 4191–4203 (2016).
- 665 88. Fleming, E. L., Jackman, C. H., Stolarski, R. S. & Douglass, A. R. A model study of the impact of source gas changes on the stratosphere for 1850–2100. *Atmos. Chem. Phys.* **11**, 8515–8541 (2011).
- 670 89. Froidevaux, L. *et al.* Validation of Aura Microwave Limb Sounder stratospheric ozone measurements. *J. Geophys. Res.* **113**, D15S20 (2008).
90. Lawrence, M. G., Jöckel, P. & von Kuhlmann, R. What does the global mean OH concentration tell us? *Atmos. Chem. Phys.* **1**, 37–49 (2001).
91. Montzka, S. A. *et al.* Recent trends in global emissions of hydrofluorocarbons and hydrofluorocarbons: Reflecting on the 2007 Adjustments to the Montreal Protocol. *J. Phys. Chem. A* **119**, 4439–4449 (2015).
- 675 92. Wild, J. D., Long, C. S., Barthia, P. K. & McPeters, R. D. Constructing a long-term ozone climate data set (1979–2010) from V8.6 SBUV/2 profiles. in *Quadrennial Ozone Symposium* (2012).
- 680 93. Coldewey-Egbers, M. *et al.* The GOME-type Total Ozone Essential Climate Variable (GTO-ECV) data record from the ESA Climate Change Initiative. *Atmos. Meas. Tech.*

**8**, 3923–3940 (2015).


- 685 94. Bojkov, R. D. *et al.* Vertical ozone distribution characteristics deduced from similar to 44,000 re-evaluated Umkehr profiles (1957-2000). *Meteorol. Atmos. Phys.* **79**, 127–158 (2002).
95. Blunden, J. & Arndt, Derek S., (Eds.). *State of the Climate in 2015. Bulletin of the American Meteorological Society* **97**, (2016).
- 690 96. Newman, P. A. *et al.* What would have happened to the ozone layer if chlorofluorocarbons (CFCs) had not been regulated? *Atmos. Chem. Phys.* **9**, 2113–2128 (2009).
97. Garcia, R. R., Kinnison, D. E. & Marsh, D. R. World avoided simulations with the Whole Atmosphere Community Climate Model. *J. Geophys. Res.* **117**, 1–16 (2012).
98. Egorova, T., Rozanov, E., Gröbner, J., Hauser, M. & Schmutz, W. Montreal protocol benefits simulated with CCM SOCOL. *Atmos. Chem. Phys.* **13**, 3811–3823 (2013).

695




## Top Ten References (Numbers refer to list above)


- 700 2. Molina, M.J. and Rowland, F.S.: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone, *Nature*, 249, 810–812, doi:10.1038/249810a0, 1974.  
**This paper first reported that CFCs could release chlorine in the stratosphere which would lead to ozone depletion.**
- 705 3. Stolarski, R. S. and Cicerone, R. J.: Stratospheric chlorine: a possible Sink for Ozone, *Canadian Journal of Chemistry*, 52, 1610-1615, doi:10.1139/v74-233,1974.  
**This paper first reported that chlorine chemistry could catalytically destroy ozone.**
- 710 4. Farman, J. C., B. G. Gardiner & J. D. Shanklin, Large losses of total ozone in Antarctica reveal seasonal ClO<sub>x</sub>/NO<sub>x</sub> interaction, *Nature*, 315, 207–210, doi:10.1038/315207a0, 1985.  
**This paper reports the discovery of the Antarctic ozone hole.**
- 715 5. WMO/UNEP, Scientific Assessment of Ozone Depletion, 2014.  
**This is the most recent of the quadrennial ozone assessments. It is detailed summary of the state of the ozone layer.**
17. Newchurch, M.J., et al., Evidence for slowdown in stratospheric ozone loss: First stage of ozone recovery, *J. Geophys. Res.*, 108, 4507, 2003.  
**First paper to report evidence for slowdown in ozone loss in the upper stratosphere.**
- 720 36. Solomon, S., D. J. Ivy, D. Kinnison, M. J. Mills, R. R Neely, A. Schmidt, Emergence of healing in the Antarctic ozone layer, *Science*, 353 (6296), 269–274.  
doi:10.1126/science.aae0061, 2016.  
**This paper reported a decrease in Antarctic ozone depletion attributed to chlorine and bromine chemistry.**
- 725 39. Harris, N.R.P., et al., Past changes in the vertical distribution of ozone – Part 3: Analysis and interpretation of trends, *Atmos. Chem. Phys.*, 15, 9965-9982, 2015.  
**Recent analysis of vertically resolved stratospheric ozone trends.**
- 730 59. Shepherd, T.G., et al., Reconciliation of halogen-induced ozone loss with the total ozone column record, *Nat. Geosci.*, 7, 443-449, doi:10.1038/ngeo2155.  
**This paper demonstrated the overall global agreement between long-term ozone changes and variations in stratospheric chlorine and bromine.**
- 735 60. Yang, E.-S., D. M. Cunnold, M. J. Newchurch, R. J. Salawitch, M. P. McCormick, J. M. Russell, J. M. Zawodny, and S. J. Oltmans. 2008. First Stage of Antarctic Ozone Recovery, *J. Geophys. Res.*, 113, D20308, doi:10.1029/2007JD009675.  
**First paper to report on decreasing ozone loss in the Antarctic ozone hole.**
- 740 71. Weatherhead, E. C., and S.B. Andersen, The search for signs of recovery of the ozone layer, *Nature* 441 (7089), 39–45, doi:10.1038/nature04746, 2006.  
**Paper which established the long timescales needed to detect ozone recovery in different regions.**



**Figure 1.** Latitude-height cross section of stratospheric ozone and time series of chlorine in the troposphere and stratosphere. **(a)** October mean (2004-2016) ozone concentration ( $10^{12}$  molecules/cm<sup>3</sup>) from the Microwave Limb Sounder (MLS) instrument aboard the Aura satellite<sup>89</sup>. The thick black lines denotes the location of the climatological tropopause<sup>90</sup>. Annotated text shows the main regions where ozone is most severely destroyed by halogens. The black arrows indicate the Brewer-Dobson circulation which transports air upwards in the tropics, polewards and downwards at high latitudes, with stronger transport towards the winter pole. **(b)** Observed monthly mean surface mole fraction (ppt) of selected ozone-depleting substances (left axis, see legend for colour coding) from the National Oceanic and Atmospheric Administration (NOAA) long-term monitoring program<sup>91</sup>. The thick grey line shows the evolution of total tropospheric chlorine (includes contributions from other halocarbons, e.g. HCFCs) from the World Meteorological Organization A1 scenario (right axis). **(c)** Time series of monthly mean HCl (ppt) in the tropical upper stratosphere (4.6 hPa, ~40km, pink) and lower mesosphere (0.46 hPa, ~55km, black) from the GOZCARDS satellite measurement compilation<sup>46</sup>. HCl is a degradation product of chlorine-containing ODSs and increases with altitude.




**Figure 2.** Time series of observed total (column) ozone (DU) for (a) SH mid-latitudes, (b) tropics, (c) NH mid-latitudes and (d) Antarctica in September. Shown are timeseries of the merged SBUV v8.6 data from NOAA<sup>92</sup> (light blue) and NASA<sup>21,51</sup> (dark blue), merged GOME/SCIAMACHY/GOME-2 (GSG, light green)<sup>33</sup> and the GOME/SCIAMACHY/GOME-2/OMI (GTO, dark green) datasets<sup>93</sup> as well as zonal mean data derived from ground-based data collected at the World Ozone and UV Data Center (WOUDC, pink) (updated from<sup>94</sup>). Total ozone trends are derived from a multiple linear regression (MLR) applied to the NASA SBUV data and the regression model timeseries is shown as the orange line. The **Supplementary Information** gives more details of the MLR approach used here. Linear trends (black long-dashed line) and  $2\sigma$  uncertainties (black short-dashed lines) as derived from the MLR are indicated for the periods before and after the ODS peak, estimated to be in 1996 (middle latitudes and tropics) and 2000 (Antarctic)<sup>9</sup>.




785 **Figure 3.** Time series of observed upper stratospheric annual mean ozone anomalies at 2 hPa (~40 km) in three zonal latitude bands. Data are from the merged SAGE II/OSIRIS<sup>49</sup> (light blue) and GOZCARDS<sup>46</sup> (green) records and from the BUV/SBUV/SBUV2 v8.6 merged products from NASA<sup>21,51</sup> (red) and NOAA<sup>92</sup> (pink, base period: 1998–2008). The orange curves represent effective equivalent stratospheric chlorine (EESC<sup>9</sup>), scaled to reflect the expected ozone variation due to stratospheric halogens. Updated from<sup>95</sup>.

790




**Figure 4** TOMCAT 3-D model calculations of the percent change in ozone at 40, 30 and 20 km altitude and in the column (DU) since 1960 for different assumptions in the ODS scenarios. Panel (a) shows the annual mean ozone variation at southern mid-latitudes. Panel (b) shows September mean ozone in the Antarctic. The control model simulation is shown by the black line. The dark blue and combined light and dark blue shading quantify the ozone depletion due to ODS emissions since 1980 and 1960, respectively. The light green shading shows the difference between the control simulation and one which used fixed 1996 ODS levels after 1996 and therefore quantifies the expected recovery signal. The red shading illustrates how severe ozone depletion may have become under a ‘world avoided’ (continuous ODS increase) scenario, which is a measure of success of the Montreal Protocol<sup>96–98</sup>. The orange line shows the difference in ozone between the control simulation and one with repeating 1980 meteorology, and therefore illustrates the impact of interannual dynamical variability.



810 **Figure 5.** Observed and modelled 2000-2015 ozone trends (%/decade) from simple (Reg, see text) and multiple linear regressions (MLR) for different regions. Trend results are for (top row) ozone in the upper stratosphere (2 hPa, 40 km) at (left-right) SH mid-latitudes, the tropics and NH mid-latitudes, (middle row) column ozone at (left) SH and (right) NH mid-latitudes and (bottom row) Antarctic ozone in September for (left) column and (right) 50 hPa.

815 Each panel shows on the left side (red shaded) the observed trends derived from both simple linear regression (Reg) and MLR. The right hand side (blue shaded) shows trends derived from the simple linear regression for the model control (varying ODS and dynamics) simulation, and MLR for the control simulation, the fixed dynamics (varying ODS) simulation and the fixed ODS (varying dynamics) simulation. The error bars indicate the  $2\sigma$

820 uncertainties of the trend estimate. The observational dataset (SBUV v8.6) is the same as in **Figure 2**. Note there are no observed September trends for 50hPa at  $90^{\circ}\text{S}$ - $60^{\circ}\text{S}$  because there are no SBUV measurements available for this month over the period considered.



825

830

**Figure 6.** Simulated global annual averaged total ozone response to the changes in CO<sub>2</sub> (red line), CH<sub>4</sub> (brown line), N<sub>2</sub>O (green line), and ODSs (blue line) from the GSFC two-dimensional chemistry-climate model. The total response to ODSs and GHGs combined is shown as the black line. The responses are taken relative to 1960 values. Future GHG concentrations are based on the IPCC SRES A1B (medium) scenario. Ground-based total ozone observations (base-lined to the mid-1960s) are shown as magenta cross symbols. Also shown are results from the GEOSCCM (grey line). Adapted from<sup>88</sup>.