

Assessing bioavailability of complex chemical mixtures in contaminated soils: progress made and research needs

Cipullo S., Prpich G¹., Campo P., Coulon F.*

Cranfield University, School of Water, Energy and Environment, Cranfield, MK43 0AL, UK

¹ University of Virginia, Department of Chemical Engineering

*Corresponding author: f.coulon@cranfield.ac.uk, +44 (0)1234 754981

Abstract: Understanding the distribution, behaviour and interactions of complex chemical mixtures is key for providing the evidence necessary to make informed decisions and implement robust remediation strategies. Much of the current risk assessment frameworks applied to manage land contamination are based on total contaminant concentrations and the exposure assessments embedded within them do not explicitly address the partitioning and bioavailability of chemical mixtures. These oversights may contribute to an overestimation of both the eco-toxicological effects of the fractions and the mobility of contaminants. In turn, this may limit the efficacy of risk frameworks to inform targeted and proportionate remediation strategies. In this review we analyse the science surrounding bioavailability, its regulatory inclusion and the challenges of incorporating bioavailability in decision making process. While a number of physical and chemical techniques have proven to be valuable tools for estimating bioavailability of organic and inorganic contaminants in soils, doubts have been cast on its implementation into risk management soil frameworks mainly due to a general disagreement on the interchangeable use of bioavailability and bioaccessibility, and the associated methods which are still not standardised.

This review focuses on the role of biotic and abiotic factors affecting bioavailability along with soil physicochemical properties and contaminant composition. We also included advantages and disadvantages of different extraction techniques and their implications for bioavailability quantitative estimation. In order to move forward the integration of bioavailability into site-specific risk assessments we should (1) account for soil and contaminant physicochemical characteristics and their effect on bioavailability; (2) evaluate receptor's potential exposure and uptake based on mild-extraction; (3) adopt a combined approach where chemical-techniques are used along with biological methods; (4) consider a simplified and cost-effective methodology to apply at regulatory and industry setting; (5) use single-contaminant exposure assessments to inform and predict complex chemical mixture behaviour and bioavailability.

KEYWORDS: bioavailability, partitioning, contaminated land, risk assessment, sorption, ageing

1 Introduction

Contaminated sites are often impacted by a wide range of organic and inorganic chemical mixtures, for example, heavy metals, polycyclic aromatic hydrocarbons (PAHs), phenols or chlorinated hydrocarbons (CHC) (European Environment Agency, 2012). These contaminants will often form complex mixtures in soil that complicate the assessment of risk and the achievement of site-specific remediation objectives (Renoux et al. 2013). Unlike single contaminants, the physico-chemical interactions of mixed contaminants are not well understood as the additive, synergistic or antagonistic effects of mixtures will often yield bioavailability values that differ from those of individual contaminants (Ramakrishnan et al. 2011). Poor understanding about the fate and behaviour of contaminant mixtures in soil limits the effectiveness of risk-based contaminated land management decisions.

Risk assessment is an established methodology that is employed to assess the potential impacts of contaminants on human and ecological health (Vegter et al. 2002). Reflecting regional legislation, expertise, and socio-economic issues, several risk-based contaminated land management frameworks have been published to support environmental management decisions (Brassington et al., 2010, Kabari et al., 2016). Typically, these frameworks use tiered assessment approaches. A limitation of such frameworks, similar to that of exposure assessment methods, is the inability to assess the risk posed by complex chemical mixtures. In fact, these frameworks are based on conservative risk screening levels and therefore tend to overestimate the risk, as they do not take into account the amounts of chemicals potentially bioavailable in soil and bioaccessible to organisms. Determining appropriate site specific measures and remedial objectives depends on our understanding of contaminant partitioning and interaction with the soil matrix over time. Measuring the total concentration of contaminants in soil does not provide a useful basis for the evaluation of the potential risks to humans and the environment. The variety of physical–chemical properties, and thus differences in the migration and fate of individual compounds, as well as the toxicity of different fractions and compounds must be taken into account in risk assessments.

Over the last 30 years, accounting for the bioavailable nature of soil contaminants has received increasing attention. As a result, a great amount of scientific literatures have reported on the development of methods to estimate the bioavailable fraction of these contaminants. Despite this progress, implementation of these methods into contaminated land decision-making processes has not yet been statutorily defined, and uncertainties remain on how bioavailability should be assessed and integrated into existing risk based management frameworks (Ortega-Calvo et al. 2015; Wu et al., 2013; Harmsen & Naidu 2013; Naidu et al. 2015)

In this review we will highlight the factors that influence the bioavailability of chemical in soil and will discuss the challenges that complex chemical mixtures pose. We will critically review the extant literature to assess the use of bioavailability in contaminated land risk assessments. Finally, we will offer suggestions for how bioavailability could be integrated into existing contaminated land risk assessment frameworks.

2 Bioavailability concept

Defined from a chemical perspective, bioavailability is the fraction of freely available (not sorbed or sequestered) contaminant in the environment that is mobile and thus most likely to lead to human exposure (Dean and Scott, 2004; Ruby et al., 1996). Similarly, Semple et al. (2004) defined bioavailability as the contaminant fraction “freely available” in a medium and able to reach the cellular membrane of an organism over a given time. Thus, for a contaminant to be bioavailable it must be mobile and there must be a likelihood for exposure with a biological membrane.

2.1 Factors affecting bioavailability

Managing risk associated with chemical mixture in the environment requires an understanding of how contaminants are released, transported, and taken up by a target receptor. The different transportation and uptake pathways that affect the quantitative estimation of bioavailable fractions of metals and oil-derived products in soil depends on both the physicochemical characteristics (Table 1), the receptors (Table 2), and other additional factors (Table 3). Among the physicochemical factors, soil characteristics (pH, soil composition, organic carbon percentage, and salinity), compound properties (hydrophobicity, aqueous solubility, and acid dissociation constant) and transformation/degradation processes are generally responsible for interactions occurring between the soil matrix and the chemical compounds (Table 1). Biological processes (e.g. bioaccumulation, biotransformation) whereby contaminants are transported into an organism, are highly dependent on the type of organism and its biology (Table 2). It is important to recognize that any combination of individual physicochemical and biological processes will affect contaminant bioavailability and exposure of receptors.

Soil matrix heterogeneity will also affect bioavailability (Farmer 1997). Among physicochemical factors, sorption is the main factor influencing the biotic and abiotic transformations happening over time (i.e ageing) in solid environmental matrices, which normally yield to a more stable solid-

associated compound and therefore a decrease in bioavailability (Zhang et al., 2014; Moyo et al., 2014; Dube et al., 2001; Kleber et al., 2007) .

Sorption, which includes absorption and adsorption, is the process whereby a chemical compound adheres (reversibly or otherwise) to the surface of a soil particle (Olu-Owolabi et al., 2014).

The sorbed substance is referred to as the *sorbate* (compound) and the material that it is sorbed to is referred to as the *sorbent* (solid phase). When contaminants are released in the soil, the chemistry of the particles and the equilibrium between phases will influence the pathways and interactions between sorbate and sorbent.

Contaminants will interact with both the mineral and organic content of soil, either sorbing to surfaces, or migrating within the porous structure of soil compartments (Reid et al. 2000). Contaminants can also dissolve into the pore water of a soil matrix, making it available for biodegradation **Figure 1**). The interaction between contaminant and soil particle will lead to different degrees of desorption (1) rapid — compounds can easily desorb and return to the pore water; (2) slow — reversible but over a longer timeframe (Ren et al. 2018); (3) non reversible — rate of contaminant removal is low and compounds are bound (sequestered) to the soil (Kuppusamy et al. 2017). The non-reversible fraction is generally believed not to be relevant for bioavailability assessment.

Partitioning of a contaminant at the solid-water interface will depend on the chemical structure of the contaminant. For example, small organic contaminants and low molecular weight PAHs could dissolve into the soil pore water, or could be rapidly sorbed onto the particle surface (Vicent et al. 2013; Abdel-Shafy & Mansour 2016). PAHs with high molecular and larger organic molecules with non-polar structures, on the other hand, tend to sorb onto the non-polar, condensed organic domains of soils (Loibner et al., 2000). These fraction resist degradation and will persistent. However, even small molecules can become persistent environmental pollutants, for example, chloro-organic compounds show a great stability and recalcitrance due to their C–Cl bond (Nikel et al., 2013). Metals also sorb to soil particles, particularly iron hydroxides, clays, and carbonate minerals, and can form solid stable compounds with oxygen and sulphur, becoming irreversibly enclosed and thus, not bioavailable.

1 **Table 1** : Physicochemical factors influencing bioavailability of metals and oil-derived products in soil (similarities and differences)

Factors	Metals	Implication for element behaviour and bioavailability	Reference of special interest	Oil derived product	Implication for oil compounds behaviour and bioavailability	Reference of special interest
Contaminant characteristics	Present in different elemental forms (metal speciation)	Metals' bioavailability can increase or decreased depending on the chemical form. Formation of sulphide cause low solubility (low bioavailability).	(Violante et al. 2010) (Rinklebe et al. 2016) (Liu et al. 2016) (Shahid et al. 2017)	Molecular weight , polarity hydrophobicity , solubility octanol partitioning coefficient (K_{OW}) sorption coefficients (K_{OC} , K_d), acid dissociation constant (pK_a)	Highly complex chemical mixture and concentration constantly changing due to transformation and interactions with environmental media over time	(Duan & Naidu 2013) (National Research Council 2014) (Trellu et al. 2017)
Soil characteristic and Sorption Desorption	Influenced by both geochemical processes (e.g., redox/pH) and soil characteristics (e.g particle size, organic content).	High pH form insoluble metal (decrease in bioavailability), low pH form free ionic species or organo-metals (increase in bioavailability). The presence of mineral phosphates and carbonates decreases bioavailability.	(Lomaglio et al. 2017) (Wang et al. 2016) (Pauget et al. 2011) (Pan et al. 2016) (Tahervand & Jalali 2016)	Quantity and type/quality of organic carbon, clay content, organic content (condensed humic material, soot particles), and soil organic matter can influence bioavailability.	Binding of PAHs to condensed organic domains rend these compounds less bioavailable. Adsorption can be also influenced by pH, depending on the presence or absence of intrinsic positive or negative charges on the compounds functional groups.	(Cecchin et al. 2016) (Lukić et al. 2016) (M. Zhang et al. 2014) (Sabljic & Nakagawa 2014) (Wu et al. 2013) (Chen et al. 2017) (Yu et al. 2018)
Transformation Degradation (biological/chemical)	No degradation	Metal cannot be degraded; inorganic contaminants can only be bio accumulated or sequestrated (Olaniran et al., 2013). Their bioavailability depends on the partitioning and distribution across soil substrates. However presence of metals (if bioavailable) can inhibit organic compound	(H.-Y. Yu et al. 2016) (Palleiro et al. 2016) (Young 2013)	Both biotic (microbial degradation), and abiotic degradation (volatilisation leaching, photodegradation) can lead to transformation and degradation of organic compounds in soil.	Bioavailability of organic compounds over time tends to decrease due to diffusion into soil particles, formation of stabile complexes, and to microbial degradation.	(Sihag et al. 2014) (Vila et al. 2015) (Ghosal et al. 2016) (Marquès et al. 2016) (Alegbeye et al. 2017)

degradation interfering with microbial processes.

Oxidation/Reduction Cation exchange capacity, and soil pH	Influenced by the presence of organo-mineral colloids (adsorption). Complexation with humus, precipitation in presence of clay mineral and Fe, Mn, Al oxides and carbonates.	Reducing conditions due to a high content of organic carbon and/or sulphide can cause formation of less soluble species e.g Cr(III)	(Ashraf et al. 2012) (H. Y. Yu et al. 2016) (An et al. 2015) (Schneider et al. 2016) (Kunhikrishnan et al. 2016)	Changes in pH can influence mostly ionizable organic compounds, impacting sorption and removal of organic solutes from solution (Naidu, 2011). Changes in redox potential and pH can accelerate oxidation of organic contaminants (Eggleton and Thomas, 2004).	Both mineral and humic substances can impact bioavailability of organic pollutants via oxidative and reductive transformation processes.	(Ling et al. 2015) (Xiao et al. 2014) (Zhang & Fan 2016) (Zhang et al. 2015)
--	--	---	--	--	--	---


2 **Table 2** : Biological factors influencing bioavailability of metals and oil-derived products in soil (similarities and differences)

Factors	Metals	Implication for element behaviour and bioavailability	Reference of special interest	Oil derived product	Implication for oil compounds behaviour and bioavailability	Reference of special interest
Uptake	Metals uptake is typically based upon bioassay exposures to a dissolved chemical, therefore highly dependent on the metals solubility in solution, and oxidation states.	Highly dependent on the system considered for example in plants the bioavailability of a certain metal in the water phase, depends on root structure, but also presence/absence of organic acids exudate (such as citrate and oxalate). For aquatic species bioavailability depends on both ingestion of metal-enriched sediment during feeding, and uptake of metal suspended particles from solution (du Bray 1995).	(Wyszkowska et al. 2012) (Seshadri et al. 2015) (Tangahu et al. 2011) (Rüdel et al. 2015)	Depend on multiple factors such as concentration in soil, its chemical form, soil pH, biological species, and uptake pathways of specific species.	Depend on where and how an organism lives and feeds in the soil or sediment.	(Peters et al. 2016) (Juhasz et al. 2014) (Beriro et al. 2016) (Lal et al. 2015) (Rostami & Juhasz 2011)
Bio concentration, bioaccumulation, and biotransformation	Metal bioaccumulation (in bacteria, fungi, and plants) can happen through biosorption or absorption and uptake. Metal can potentially interact and affect function of enzymes involved in biodegradation of chlorinated organic compounds.	Depend on uptake, levels of fats (lipids) within the organism, metabolism, age, growth life stage, and gender.	(Berthelot et al. 2008) (Jaishankar et al. 2014) (Tchounwou et al. 2012) (Khan et al. 2015)	Strong correlations between the bio-concentration factor, bioaccumulation factor and the octanol: water partition coefficient (K_{ow})	Depend on uptake, levels of fats (lipids) within the organism, metabolism, age, growth life stage, and gender.	(Fantke et al. 2016) (McLachlan et al. 2011) (Pampanin 2017) (Vasseur & Bonnard 2014)

5 **Table 3** : Additional factors influencing bioavailability of metals and oil-derived products in soil (similarities and differences)

Factors	Metals	Implication for element behaviour and bioavailability	Reference of special interest	Oil derived product	Implication for oil compounds behaviour and bioavailability	Reference of special interest
Ageing	A rapid uptake via electrostatic adsorption is usually followed by a secondary transformation that form a more stable complex.	Ageing can have an effect on inorganic contaminant, where metal precipitation can occur rapidly causing a decrease in bioavailability and toxicity. However is less clear how stable/reversible is the process (Hamon, McLaughlin, and Lombi 2006).	(Wijayawardena et al. 2015) (Liang et al. 2014) (Wang et al. 2017) (Romero-Freire et al. 2017) (Jiang et al. 2017)	Different processes might occur: incorporation into natural organic matter (absorption), slow diffusion into small pores (soil intraparticle).	Overall a decrease in bioavailability has been observed during time due to different factors (dilution, dispersion biodegradation, volatilisation, and irreversible sorption).	(Duan et al. 2014) (Duan et al. 2015) (An et al. 2017) (An et al. 2017) (Smith et al. 2011) (Liu & Haderlein 2013)
Co-contaminant interaction	Metal-metal interaction is mostly competitive affecting affinity for soil-surface and sorption. For instance Zn is a competitor for Cd and Pb sorption sites . Metal-organic joint interaction such as: Cu-pyrene (Chigbo et al. 2013) can significantly enhance heavy metals concentration in shoot and roots. Heavy metals transport can be enhanced by the presence of organic contaminants due to : association with mobile colloids, formation of metal-organic	Metals that compete for the same sorption sites can trigger the release of the competitor metal, enhancing its bioavailability. Necessity of addressing contaminant as a mixture	(Sun & Zhou 2010) (Van Genderen et al. 2015) (Meyer et al. 2015) (Zhao et al. 2016) (Chigbo et al. 2013)	Organic-metal interaction: high concentration of inorganic might influence mobility of PAHs. Organic-organic interaction: competitive displacement, co-solvency. Molecules with similar structure are highly competitive for sorption sites because of their interchangeability.	As for metals, the organic-organic interactions might challenge bioavailability predication. The sorption/desorption rate can be altered by competitive displacement processes and therefore manifest a non-linear behaviour. In addition some natural compounds might also share a similar structure and therefore displace anthropogenic compounds increasing their bioavailability. Necessity of addressing contaminant as a mixture	(Olaniran et al. 2013) (Gauthier et al. 2014) (Biswas et al. 2015) (Wuana et al. 2014)

complexes that are not
sorbed onto the surface
(competitive sorption).



7

8 **Figure 1:** Process of sequestration of the compounds in soil, adapted from (Reid et al., 2000), bold
 9 textbox indicates the non-desorbing fraction (irreversible processes), dashed textbox indicates the
 10 rapidly-desorbing fraction (reversible processes), and the highlighted textbox represent the
 11 dissolved fraction (bioavailable).

12

13 Contaminant retention is largely regulated by soil particle size distribution (Table 1). Smaller
 14 particle sizes provide a greater surface for interactions with hydrophobic organic chemicals (Capri
 15 et al., 2004). Clays and fine-grained sediments have the greatest surface area and therefore a high
 16 capacity to retain organic and inorganic compounds. Further to this, the presence of oxides (Fe and
 17 Al oxides, hydroxides, and oxyhydroxides) along with reactive calcium carbonate (CaCO_3) can
 18 enhance organic and inorganic contaminants retention (Loibner et al., 2006) and therefore favour
 19 the biological stabilization of organic carbon (encapsulation) (Heng et al., 2010). Such mechanisms
 20 will however hinder microbial degradation of the compounds of concern due to reduced
 21 accessibility (Krull et al., 2001).

22

23 Hard- and soft-organic matter are also associated with retention and ageing processes (Table 3).
 24 Soil organic matter (SOM) is formed from natural organic matter (e.g. vegetal decomposed
 25 material), animal residues at various stages of decomposition, fulvic acids, and humic acids
 26 (Sharma et al., 2010). Generally, SOM is thought to be composed of “soft carbon” (amorphous or
 27 hydrolysable carbon), and “hard carbon” (condensed or non-hydrolysable carbon) constitute of

28 kerogen, black carbon, and lignin (Weber et al., 1992). Black carbon and kerogen, in particular, can
29 bind tightly the organic contaminants reducing their solubility and/or dissolution rate, and thus
30 bioavailability (Stroud et al. 2007; van Elsas et al. 2006; Berkowitz et al. 2008). Large amounts of
31 organic matter in the soil have also been shown to effect the residence time of organic matter-
32 associated metals. For example, when organic matter is oxidized, the associated metals are likely to
33 be released, becoming more bioavailable. Conversely, soil with low organic matter content will
34 often accumulate oxide minerals (e.g. clay) that favour the complexation of both metals and
35 metalloids, thus reducing the bioavailable fraction (John & Leventhal 1995). Soil properties are site
36 specific and will vary from one site to another, therefore, if two sites contain equivalent amounts of
37 a certain contaminant, their bioavailability may still vary significantly, depending on how tightly
38 the chemical is bound to the soil.


39

40 The ratio between bioavailable and non-bioavailable fractions is shown in Figure 2. Over time, the
41 proportion of bioavailable contaminant will decrease, relative to the non-bioavailable fraction.
42 Pollutants in soil and sediment do not disperse quickly and the desorption and remobilization of
43 metals and oil-derived products in soil are considered long-term processes. For example, heavy
44 metals associated with fluvial sediments can display a residence time from days to years, on the
45 order of 100 – 1000 years (depending on stream-flow dynamics) (Ciszewski & Grygar 2016;
46 Coulthard & Macklin 2003). Heavy metals also have a long residence time in soil (Sayadi et al.
47 2017), ranging from 500 to 3000 years (Lepp 2012; Jørgensen 2000; Ayres 1992; Alloway 1995).
48 The process of aging can enhance the amount of absorbed heavy metals in soil, where a
49 redistribution from weakly bound fractions to more strongly bound fractions can be observed
50 (Settimio et al. 2014; Wang et al. 2015). Recent metal contamination tends to be more reactive and
51 prone to dissolution compared to older contamination where the elements might be in crystalline
52 forms (Lynch et al. 2014), which presents a lower environmental risk (Baran et al. 2015;
53 Environmental Agency 2008).

54

55 Sediment and soil-bound organic contaminants can persist over decades without significant
56 concentration reductions. Chlorinated or hydrophobic contaminants, in particular, tend to desorb
57 very slowly over time with a desorption rate on the order of years, due to their hydrophobicity
58 (Eggleton & Thomas 2004). Though oil derived products tend to persist in soil, the PAH
59 bioavailable fraction will decline exponentially over time (Yang et al. 2016). Weathered
60 hydrocarbon residues pose negligible risks to human health, and this is reflected in post-treatment
61 remedial objectives (Coulon et al., 2010; Jiang et al., 2016). When bioavailable fractions are high
62 (despite being below risk-based clean-up levels) further treatment and more stringent clean-up

63 levels should be mandated to reduce the elevated risk of exposure that is present (Cipullo et al.,
64 2017).
65



66
67 **Figure 2:** Relationship between the percentage of bioavailable and non-bioavailable contaminants
68 in soil, exposure risks and risk based clean-up level (adapted from Reid et al., 2000; Tri-Service
69 Ecological Risk Assessment Workgroup, 2003).

70
71 The necessity of collecting case-specific parameters can challenge the development of a unified
72 methodology to assess the bioavailable fraction, and to determine the risks to human and
73 environment in a straightforward way. A number of analytical methods to assess readily (bio)
74 available compounds in soil and sediments are available, and we discuss these in the following
75 sections.

76 77 **2.2 Methods for estimating bioavailability of heavy metals**

78 Metals can be present in soil as either free metal ions, forming various soluble complexes with
79 inorganic or organic ligands; or associated with colloidal and mineral materials (McLean and
80 Bledsoe, 1992) becoming strongly incorporated with the soil matrix (inert). Most of the divalent

81 heavy metal cations (e.g., Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} and Zn^{2+}) share a relatively similar structure
82 and therefore display similar behaviour in soil (Olaniran et al., 2013).

83 Total metal concentration estimated with acid digestion (e.g. aqua regia) mobilizes all forms of
84 metal in soils and sediment. This measure refers to both metal content in particulate (sorbed +
85 precipitated), and dissolved (inorganic complexes + organic complexes + free ionic forms)
86 fractions. Particulate metals do not contribute to the solid-solution distribution (potentially
87 bioavailable), and may only become available through very slow desorption processes. This fraction
88 does not provide appropriate basis for expressing metal bioavailable (labile) concentration in soil,
89 thus presumably not readily available for receptor uptake.

90
91 Several approaches are used to determine the pool of labile metals (concentration and distribution)
92 in soils and sediments; including exchange resins, diffusive gradient in thin films (DGT),
93 conventional single-step extractions, and sequential extractions (Table 4). Passive samplers (e.g
94 exchange resins) act as ion sinks and are used to quantify free ion activities, solution fractions, and
95 labile pool concentrations of metals in soils (Qian and Schoenau, 2002; Ge et al., 2005). Free ion
96 activity in solution represents the most relevant parameter for assessing bioavailability and toxicity
97 of metals in contaminated soils. The use of exchange resins has been successfully applied to predict
98 uptake of metal in the environment (e.g. in plants (Peijnenburg et al., 2007)).

99 To date, no standard technique has been validated. Diffusive gradients in thin films (DGT), is a
100 relatively cost-effective technique, based on a layer of hydrogel and resin gel. The DGT devices
101 allow for the passive accumulation of labile species from a solution (in real time), and have been
102 used to assess in situ the fraction of metals dissolved in water (Zhang 1998; Hooda et al., 1999;
103 Parker et al., 2016).

104
105 As an in situ sampling technique, DGT can be also used in soil to determine the labile (bioavailable)
106 fraction of elements, and to understand how this fraction changes in the environment. However,
107 continuous depletion of metals from pore water can limit the diffusion of analytes to the DGT
108 sampler, affecting the estimation of the available pool of metals (Peijnenburg et al. 2014). In
109 addition, development and validation of a DGT method to establish accumulation in biological
110 endpoints remains a challenge, due to the complexity of the uptake by model organisms and
111 inconsistent results (Menegário et al. 2017).

112
113 Single-step extraction (Table 4) uses a wide range of extractants including: salt solutions ($CaCl_2$,
114 $NaNO_3$, NH_4NO_3 , $Ca(NO_3)_2$, $AlCl_3$, $MgCl_2$), acid solutions (HNO_3 , CH_3COOH , HCl) and chelating
115 agents (EDTA, DTPA). These tests were initially designed to predict nutrient deficiency in soil, and

116 generally contain organic chelates and acids in order to mimic plant metals uptake (National
117 Research Council 2003). Single-step extraction were lately adapted to measure the labile
118 concentration of metals in soil and sediments, potentially available for ecological receptors uptake
119 (Alvarez et al. 2011).

Method	Advantages	Disadvantages		Reference
Passive sampler	<ul style="list-style-type: none"> Passive samplers are relatively low-cost, can be used for insitu measurements (Menegário et al. 2017), can achieve low detection limits (Peijnenburg et al. 2014), and allow long-term trends assessments. DGT was found to be a good tool for measuring in situ metal bioavailability in sediments (Ren et al. 2015). DMG is a relevant tool for <i>in situ</i> assessment of environmental risks posed by metals (Perez et al. 2016) Passive sampler measurement can be translated into fugacity models and equilibrium studies to understand chemical potential activity and estimate potential risk (Amiard-Triquet et al. 2015). 	<ul style="list-style-type: none"> Little attention is given to the effects that water chemistry and method of field deployment may have on uptake kinetics (Mills et al. 2014). In some cases require a time-consuming elution step, with acids, in order to retrieve the analyte from the solid sorbent phase (Almeida et al. 2014). DGT and DMT not yet considered suitable for routine analysis due to poor detection limit, time-consuming procedures, and a lack of validation (Brand et al. 2009) Deployment of DGT samplers where nanoparticles are high may result in an overestimation of dissolved metal concentrations (Pham et al. 2015) 	Exchange resins	(Cantwell et al. 1982) (Holm et al. 1995) (Lorenz et al. 1997) (Christensen & Christensen 1999)
			Diffusive gradient thin film (DGT) and Donnan membrane technique (DMT) Diffusive Milli-Gels (DMG)	(Davison & Zhang 1994) (Agbenin & Welp 2012) (Koopmans et al. 2008) (Pampura et al. 2006) (Weng et al. 2005) (Perez et al. 2015)
Stripping voltammetry	<ul style="list-style-type: none"> High sensitivity, high reproducibility, and mainly used for the detection of trace levels of heavy metal ions (Almeida et al. 2014). Found to be suitable for assessment of heavy metals bioavailability to plants (Dyrtrova et al. 2008). 	<ul style="list-style-type: none"> Limitations for on field measurements, sample perturbations due to sample handling and storage (Rurikova & Kudrava 2006) Adsorption effects of humic and fulvic acids in soil accompanied by the metal complexation can limit the success of this method (Rurikova & Kudrava 2006) 	Anodic (or cathodic) stripping voltammetry	(Sauvé et al. 1997) (Zima & van den Berg 1994) (Davidson & Smyth 1979)
Competitive chelation	<ul style="list-style-type: none"> Method has good sensitivity and can provides reliable estimates of ion activities (Amacher 1984) 	<ul style="list-style-type: none"> Equilibrium between chelate and soil can be take a long time to be attained (Norvell & Lindsay 1972; Norvell & Lindsay 1969) The success of the metod depend on abundancy of metal of interest and the selected competing metal(Workman & Lindsay 1990) 		(Xue et al. 1995)
Ion exchange	<ul style="list-style-type: none"> Ion exchange results were comparable to ion selective electrode, and anodic stripping voltammetry (Ge et al. 2005). Simple, cost-effective, relatively easy to 	<ul style="list-style-type: none"> Requires a characterization of the resin adsorption properties (Weng et al. 2005). Soil solution composition need to be 	Cation exchange resin	(Sunda 1984) (Apte & Batley 1995) (Qian & Schoenau 2002) (Ge et al. 2005)

	<p>use, and applicable to different soil type (Qian & Schoenau 2002)</p> <ul style="list-style-type: none"> • Possibility of simultaneous multimetal measurement (Weng et al. 2005) 	<p>considered during speciation analysis (Fotovat & Naidu 1997)</p>		
Single extraction salt solutions	<ul style="list-style-type: none"> • Single extractions can be used to perform fast screening analysis of the labile pool of elements in soils and sediments (Sakan et al. 2016). • Significant positive correlations between the single extractions methods results and lettuce shoot content were obtained for several metals (Pinto et al. 2015). • CaCl₂ extraction has been reported as being a good proxy for bioavailability of metals in soils to plants (Houben et al. 2013) and was correlated with concentrations of potential harmful element in plant (Qasim et al. 2015) 	<ul style="list-style-type: none"> • Chemical extractions can hardly account for the complex processes involved in metals uptake by plants therefore not sufficient for estimating soil metal bioavailability to plants (Krishnamurti et al. 2015). • Complexing and chelating extractants (EDTA and DTPA) showed poor correlation with phes concentrations in plant (Qasim et al. 2015) • No consensus on best single step extraction conditions to extract maintain integrity of arsenic species (Sun et al. 2015) • At low reactive element to organic matter ratios, dilute nitric acid extraction (0.43 M) can underestimates concentration of geochemically reactive elements with a particularly high affinity for organic matter or oxides (Groenenberg et al. 2017) 	0.01-1 M CaCl ₂	(Houba et al. 2000) (Young et al. 2000) (Novozamsky et al. 1993) (Ure 1996)
			0.1 M NaNO ₃	(Gupta & Aten 1993)
			NH ₄ NO ₃	(Novozamsky et al. 1993)
			0.1 M Ca(NO ₃) ₂	(Meers et al. 2007)
			0.3M AlCl ₃	(Hughes & Noble 1991)
			0.02-0.1 M MgCl ₂	(Makino et al. 2006)
Single extraction acid solutions	<ul style="list-style-type: none"> • Leaching test employing a neutral salt solution (CaCl₂ or NH₄NO₃) is considered to be sufficient to measure the bioavailable fraction of mobile metals (in particular Cd, Ni, and Zn) (Kim et al. 2015). 	<ul style="list-style-type: none"> • The single extraction method involving EDTA presented good precision (Sahito et al. 2015). 	different concentrations of HNO ₃	(Tipping et al. 2003) (Almås et al. 2007) (Novozamsky et al. 1993)
			CH ₃ COOH	(Ure 1996)
			HCl	(Novozamsky et al. 1993)
Single extraction Chelating agents:	<ul style="list-style-type: none"> • The single extraction method involving EDTA presented good precision (Sahito et al. 2015). • Using the single-step extraction (EDTA or acetic acid) allows to predict metal extractable content (bioavailable) in vineyard soil-grapevine system (Vázquez Vázquez et al. 2016) 	<ul style="list-style-type: none"> • Using the single-step extraction (EDTA or acetic acid) allows to predict metal extractable content (bioavailable) in vineyard soil-grapevine system (Vázquez Vázquez et al. 2016) 	EDTA	(Ure 1996) (Cappuyns 2012)
			DTPA	(Leggett & Argyle 1983) (Lindsay & Norvell 1978)
Sequential extractions	<ul style="list-style-type: none"> • These methods are simple, low cost, applicable to different soil type, and results are often comparable (Rosado et al. 2016). • BCR method provide relevant information on the relationships between soil characteristics and metal potential fractions for uptake by the plant (Sungur et al. 2014). 	<ul style="list-style-type: none"> • Sequential extraction are inadequate for determining the extraction kinetics, and subjected to high risk of bias due to re-adsorption processes (Rosas-Castor et al. 2015) • BCR method drawbacks includes lacks of specificity and difficulty in interpreting results (Huang et al. 2014), • BCR method results tedious and time- 	BCR extraction	(Ure et al. 1993)
			modified BCR extraction	(Rauret, López-Sánchez, Sahuquillo, Barahona, et al. 2000)
			four-step chemical fractionation procedure	(Tessier et al. 1979)
			four-step chemical fractionation	(Elliott et al. 1990)

<ul style="list-style-type: none"> • BCR method showed correlations between available metal and the plant uptake (Fernández-Ondoño et al. 2017). • The modified version of the three-step procedure proposed and validated by the BCR (Community Bureau of Reference) could potentially be accepted as the standard method(Ahmadipour et al. 2014) • CISED method can provide a powerful tool for understanding metal fractionation in soils (Cave et al. 2015) • CISED is a valuable methodology for studying the solid-phase fractionation of potential harmful element in soil and potential bioavailability (Reis et al. 2014; Palumbo-Roe et al. 2013; Cox et al. 2013) • Sequential leaching studies can help understanding leachability, solubility, and mobility of heavy metal,;therefore allowing to make assumptions on metal bioavailability for risk assessment (Kaakinen et al. 2015) 	<p>consuming due to long shaking time and filtration requirements (Matong et al. 2016).</p> <ul style="list-style-type: none"> • Interpretation of data from the CISED extraction may more challenging than selective chemical extractions (Reis et al. 2014) • Main limitation associated with sequential extraction procedures is the long time associated extraction (Khan et al. 2013). • Limitations associated with sequential extraction methods includes; redistribution of analytes among phases, incomplete extraction, non-selectivity of reagents, and precipitation of other minerals (Selim 2015) • Laborious and time-consuming techniques, results often of difficult interpretation (Domini et al., 2011). 	procedure	
		five-step chemical fractionation procedure	(McLaren and Crawford 1973)
		six-step chemical fractionation procedure	(Miller et al. 1986)
		seven-step chemical fractionation procedure	(Krishnamurti et al. 1995)
		non-specific sequential extraction (CISED)	(Cave et al. 2004)

121 Complex organic reagents (EDTA, DPTA) are also used to mimic the organic exudates produced by
122 plants, and have been positively correlated with metal concentration found in plants (Domínguez
123 2008) and further exploited to mimic the bioavailable fraction. Chelating agents, due to their high
124 affinity for metal ions, are used to enhance the solubilization of metal(oids) through the formation
125 of soluble chelates (Bolan et al. 2014). Caution is needed, however, because studies have shown
126 that results are not consistent and robust across different types of soil (National Research Council
127 2003), and are highly dependent on extraction conditions and trace elements speciation (Cappuyns
128 2012).

129

130 Single-step extraction techniques widely vary in type of extractant used, its concentration, soil:
131 solution ratio, and extraction time. They can partially dissolve trace elements associated with
132 different fractions (e.g pore water and exchangeable), which provides useful information about
133 metals behaviour (Alvarez et al. 2011). Though relatively simple to use (minor sample handling), in
134 single-step extraction mode the non-labile metal fraction might also become solubilized, which
135 might cause an overestimation of the labile pool of several metals (Moreno et al. 2005; Qasim et al.,
136 2015).

137

138 While passive samplers and single-step sequential extraction have been used to measure labile
139 metals, these methods do not provide information on the fate and behaviour of contaminants over
140 time. Sequential extraction, however, can be used to quantify the distribution of metals and assess
141 the mobility of potentially harmful elements over time (Sungur et al. 2015). The procedure involves
142 leaching successive fractions of metal by increasing the strength of an acid solution (HNO₃, HF-
143 HClO₄, HClO₄, HCl, CH₃COOH) or other reagents (such as Na₄P₂O₇, NH₂OH) used for each
144 phase association (Zimmerman and Weindorf, 2010). The number of step may vary from 3 to 7
145 (Table 4). Sequential extraction has been used for over 30 years (Tessier et al., 1979) and has been
146 modified to create the community Bureau of reference method (BCR) (Rauret, López-Sánchez,
147 Sahuquillo, Muntau, et al. 2000) and the Chemometric Identification of Substrates and Element
148 Distribution (CISED) method (Cave, et al., 2004), which overcome non-selectivity and
149 redistribution of trace elements. The BCR method has been validated against a sediment reference
150 material (BCR-701) and provides extractable concentrations for several metals (Rodgers et al,
151 2015). The non-specific sequential extraction CISED has been validated against a sediment
152 reference material (BGS-102) and uses chemometric data processing to provide mechanistic
153 information about metal-soil phase associations (Gál et al., 2006). Results from CISED could be a
154 powerful tool to understand how mineralogical forms might affect bioavailability, yet considerably
155 more chemometric and geochemistry information need to be applied (Giacomino et al., 2011).

156 Despite the large amount of information that sequential approaches can provide the standardisation
157 and application of these technique in commercial laboratories is limited because of the laborious
158 and time-consuming nature of these techniques and the difficulty of interpreting the results (Alvarez
159 et al., 2011).

161 **2.3 Methods for estimating bioavailability of oil-derived products**

162 Estimating the bioavailability of organic compounds and integration into decision processes
163 remains a scientific and regulatory challenge. Research into the bioavailability of oil-derived
164 contaminants has received considerable attention in the last 20 years (Thompson 2016). In the
165 context of implementing bioavailability into regulatory frameworks it is important to both quantify
166 the (potentially) bioavailable fraction, but to also understand the mobility and behaviour of
167 contaminants in soil in order to assess potential effects of complex contaminations on receptors.
168 Empirical approaches are often used to predict contaminant toxicity or assess the effectiveness of
169 remediation treatments (Environment Agency 2006; Environment Agency 2010) Computational
170 methods to integrate the multitude of compounds and molecular structures has proven most
171 challenging, for example, crude oil and its composition of several hundred different organic
172 compounds, each of which possesses chemical characteristics that influence the rate and efficacy of
173 degradation (Weng et al., 2015). Molecular weights have been shown to most strongly affect a
174 compound's persistence in soil (Atlas, 1995). Moreover, complex physico-chemical interactions
175 between different compounds can lead to unexpected or poorly understood reactions (e.g. co-
176 solvency), which might alter the bioavailable nature of a compound in mixture.

177 A wide range of analytical procedures have been used to estimate the bioavailable fractions of
178 organic contaminants in mixtures (e.g. oil constituents) (see for review Brand et al., 2012; Ortega-
179 Calvo et al., 2015). These procedures can measure different fractions: freely dissolved fraction
180 (passive samplers), which measure the dissolved (actual) concentration in a matrix, and rapidly
181 desorbed fraction (non-exhaustive techniques), which uses extractants to recover compounds from
182 soil (Table 5).

183

Type	Method	advantages	disadvantages
Passive sampler (solid phase)	<ul style="list-style-type: none"> • Polydimethylsiloxane(PDMS) • Solid-Phase Microextraction (SPME) • Polyoxymethylene solid phase extraction (POM-SPE) • Triolein embedded cellulose acetate membrane (TECAM) • C18 membrane disks • Diffusive gradients in thin films (DGT) • Semipermeable membrane devices 	<ul style="list-style-type: none"> • Consistent relationship between chlorobenzenes levels in biota and in the PDMS-SPME fibers (van der Wal et al., 2004) • SPME can accurately measure freely dissolved pore water concentrations to estimates earthworms uptake (Van Der Wal et al. 2004) • PDMS is very sensitive and able to detect PAHs freely dissolved pore water concentrations (ng/L) (Laak et al., 2006). • SPME shows good correlation between bioaccumulation of organic contaminants on a wide range of organisms (You and Landrum, 2006) • SPME and POM-SPE generally are able to predict PAHs concentrations in earthworms (Gomez-Eyles et al., 2011) 	<ul style="list-style-type: none"> • SPME and POM-SPE tended to under predict PAHs root concentrations (Gomez-Eyles et al., 2011) • SPME measures truly dissolved concentrations but it is not able to measure compounds associated with dissolved organic matter (ECETOC, 2014). • Poor correlation between availability of PAHs in soil and bio concentration factors in earthworms (Bergknut et al., 2007)
Fluid-phase extractions	<ul style="list-style-type: none"> • Subcritical Water Extraction (SWE), superheated water technique (SWAT) • supercritical fluid extraction (SFE) , sequential supercritical fluid extraction (SSFE) 	<ul style="list-style-type: none"> • Good correlation between SWE extraction of 14C-activity fraction mineralized by catabolically active Pseudomonas (Latawiec et al., 2008) • Successfully used to predict rates of long-term release of organic compounds (Weber, 2001; Hawthorne et al., 2000; Miller and Hawthorne, 1998) • SWE selectively extracts the PAHs relative to the readily extracted fraction (Smith 2002) • SFE recoveries of the ‘‘mobile’’fraction of PAHs were greater than 90% (Librando and Aresta, 2004) • The amount of PCBs extracted by SFE was very close to the estimated bioavailable fraction in earthworms (Hallgren et al., 2006) 	<ul style="list-style-type: none"> • Potential degradation of analytes subjected to high temperatures. • Lack of significant correlation between the amounts desorbed/amount assimilated by earthworms (Weber et al., 2002). • Contaminants with high molecular weight ((in decreasing polarity and increasing Kaw-coefficients) showed reduced recoveries when applying SSFE (Loibner et al., 2000) • Mild SFE was not able to differentiate pyrene availability in unaged soils (Sun and Li, 2005). • using SFE for predicting bioavailability can be limited due to great variability of soil matrix (Cajthaml and Vaclav, 2005)
non-exhaustive extraction techniques	<ul style="list-style-type: none"> • Mild solvent (butanol, methanol, n-propanol, or ethyl acetate) • combination of solvent and CaCl₂ solution • surfactants (Triton X-100) 	<ul style="list-style-type: none"> • Methanol-water and n-butanol extraction of chemical mixtures were correlated with uptake by earthworms (Elsey and Lexander, 1997) • Good correlation between extractable fraction/ uptake earthworms (Kelsey et al., 1997; Tang et al., 1999) • Good correlations between extractable fractions/ bacterial genotoxicity assay (Alexander and Alexander 2000; Tang et al., 2002) 	<ul style="list-style-type: none"> • Little consistency among different soils (Chung and Alexander, 1998). • High variability in technical operation (type of mixture, shaking time) limit the comparability of data (Cachada et al., 2014). • PAHs extracted by mild solvent extraction show a similar composition to the total soil PAHs (Bergknut et al., 2007) • Mild solvent extractions consistently over predicted

			<p>PAHs biotic concentrations (Gomez-Eyles et al., 2011)</p> <ul style="list-style-type: none"> • PAHs bioavailability (estimated with butanol) and earthworm bioavailability were found to be non-related (Johnson et al., 2002). • Triton X-100 failed to predict PAHs bioavailability in contaminated sediments because extracted both readily and poorly bioavailable PAHs (Cuyppers et al., 2002).
	<ul style="list-style-type: none"> • Tenax 	<ul style="list-style-type: none"> • Tenax was found to be a matrix-independent, cheap and less time-consuming chemical method of estimating bioavailable fraction in PCB-contaminated field sediments (Trimble et al., 2008) • Successfully used to assess the bioavailability of aromatic compounds in sediment (Morrison et al., 2000; Cal et al., 2008; Harwood et al., 2012). • good correlation between “bioavailable” fraction in the sediment and quantity measured by Tenax extraction (You and Pehkonen, 2007) • Rapidly desorbing fractions of PAHs measured by Tenax have been linked to biodegradation rate (Braidia et al., 2004; Shor et al., 2003; Cornelissen et al., 1998) • Tenax extraction of PAHs, PCBs and organochlorine pesticides were correlated to bioavailability to worms (Hulscher et al., 2003) • The Tenax extraction is a good technique to predict bioavailability to earthworms of aged PAHs in soil (Lu et al., 2011) 	<ul style="list-style-type: none"> • The process involves a lot of steps in order to estimate the rapidly desorbing fraction (Xing et al., 2011) • Contaminant desorbing fractions in river sediments extracted by Tenax overestimated the bioavailable fraction of benthic invertebrates (Leppanen et al., 2003) • Few studies where correlations between Tenax extractable amount and biota-sediment accumulation were not related (Sormunen et al., 2008 and 2009; Leppa and Kukkonen, 2006) • The Tenax measure of rapidly desorbing fraction is not the only factor contributing to the bioavailability. A number of ecological factors (species and feeding habits) can affect the bioavailable fraction in model organisms, and remain undetected by Tenax extractions (Akkanen et al., 2007)
	<ul style="list-style-type: none"> • Aqueous hydroxypropyl-B-cyclodextrin (HPCD) 	<ul style="list-style-type: none"> • 1:1 correlation phenantrene extracted/degraded by microorganisms (Reid et al. 2000a) • HPCD was successfully used to predict PAHs bioavailability in contaminated sediments (Cuyppers et al. 2002). • HPCD was successfully used to predict the microbial bioaccessibility and mineralisation rate of aliphatic hydrocarbon (Stroud et al., 2008) • A significant relationship ($p < 0.01$) between HPCD extractability /mineralization (Rhodes et al., 2008) • HPCD β was found to be a good estimation of bioavailable fraction in both single, and multiple contaminants conditions (Stroud et al., 2009) • HPCD ease in sample handling and that no additional device is needed (Cui et al., 2013) 	<ul style="list-style-type: none"> • Predictability of HPCD extraction decreased for higher organisms such as earthworms (Barthe and Pelletier 2007; Hartnik et al., 2008) • Poor indicator of PAH accumulation in benthic invertebrates (Barthe & Pelletier 2007). • Cyclodextrin size and structure can limit the complexation of some PAHs (size dependent) (Villaverde and Pe 2012; Stokes et al., 2005) • Cyclodextrin extractions consistently over predicted PAHs biotic concentrations (Gomez-Eyles et al., 2011)

185 **2.3.1 Passive sampler methods (PSMs)**

186 Passive sampler methods (PSMs) (Table 5), commonly defined as biomimetic extractions
187 (ECETOC 2014), are used to measure the freely dissolved concentration (C_{free}) of contaminants
188 (Parkerton et al., 2012) in equilibrium with the rapidly desorbing fraction. These techniques are
189 based on the molecular diffusion principle, and often use polymer materials such as
190 polydimethylsiloxane, polyethylene, polyoxymethylene, and ethylvinylacetate (Parkerton et al.,
191 2012). Passive sampler methods have been used to predict PAH bioavailability, and have been
192 shown to correlate well with model organism PAH uptake and bioaccumulation (Jonker et al., 2007;
193 Gomez-Eyles et al., 2011; Muijs and Jonker, 2011). Although PSMs are valuable tools for
194 providing weight of evidence and informing regulatory decision-making, there exists a lack of
195 consensus about its implementation and standardization. The following issues were identified and
196 need to be overcome in order to further develop and implement these techniques: (1) PSM has been
197 applied to only a limited number of target compounds; (2) the complexity of the equilibrium
198 sampling methods requires better characterization of potential errors when applied in situ; (3)
199 adoption of robust quality assurance and control strategies are needed (Mayer et al. 2014).

200

201 Correlation with in-vivo measurements and bioaccessibility data are not fully validated, and the
202 complexity of (bio)accumulation mechanisms (Cachada et al., 2014) that govern toxicity responses
203 are not yet fully understood. More information and guidance on the application of passive sampling
204 for the management contaminated sediment sites can be found in the SERDP and ESTCP national
205 guidance documents (Burgess and Driscoll, 2016; Driscoll and Thompson, 2016).

206

207 Supercritical Fluid Extraction (SFE) (Table 5) uses supercritical fluids (e.g. CO₂, combination with
208 solvents) to extract a compound from a soil matrix. This technique used to study
209 sorption/desorption processes and to estimate bioavailability of organic pollutants in sediment. Data
210 obtained from SFE can be fit to a prediction model to obtain information about the
211 bioavailable fraction for PAHs (bioavailable). Different experiments found the amount of organic
212 compounds (PCBs) extracted by SFE to represent the readily extracted fraction (Smith, 2002) and
213 the bioavailable fraction uptake by earthworms (Hallgren et al. 2006). SFE was also found to
214 recover over 90% of the mobile fraction of PAHs from soil and sediments samples (Librando and
215 Aresta, 2004).

216

217 Though SPE can measure the freely dissolved fraction, it is likely to underestimate the
218 concentration/uptake in benthic organisms (with other uptake routes) and its use can be limited due

219 to great variability of soil matrix (Cajthaml and Václav, 2005). In addition, SFE was found to be
220 able to predict degradation for low molecular weight PAHs (three and four ring), with a good
221 correlation between biodegradability and bioavailability (Naidu 2011), but often overestimated
222 bioavailability of high molecular weight PAHs due to their different extractability.

223

224 **2.3.2 Non-exhaustive techniques**

225 Non-exhaustive techniques to assess the bioavailability of organic compounds in soil (Table 5)
226 include mild solvent extraction (Kelsey et al., 1997; Liste and Alexander 2002), solid sorbents (e.g.
227 Tenax) (Cornelissen et al., 2001; Hulscher et al., 2003 ,Lydy et al. 2015) and hydroxypropyl- β -
228 cyclodextrin (HPCD) (Reid et al., 2000; Cuypers et al., 2002; Stokes et al., 2005).

229

230 Mild solvent extraction consists of adding a polar solvent or mixture of solvent and water (e.g.
231 butanol, ethanol, methanol, methanol-water) to a sediment or soil sample, agitating the mixture, and
232 then analysing the extract for contaminants (Cui et al., 2013). This technique has shown good
233 correlation between extractable fraction and uptake in earthworms (Kelsey et al., 1997) and
234 bacterial genotoxicity assay (Alexander and Alexander 2000; Liste and Alexander 2002). Mild
235 solvent extraction could also be used as a proxy to estimate the contaminant bioaccessible fraction,
236 however it shows little consistency between different soil types (Chung & Alexander 1998).

237

238 Tenax is a polymer resin that, when mixed with contaminated sediment, will recover target
239 compounds. Sorbed compounds are eluted from the resin with a solvent and fresh polymeric beads
240 can be added several times (multiple steps) and harvested to measure the recoverable hydrocarbons
241 (Cui et al., 2013). Tenax has been used to assess the bioavailability of aromatic compounds in
242 sediment (Morrison et al. 2000; de la Cal et al. 2008; Harwood et al. 2012).

243

244 Hydroxypropyl-cyclodextrin (HPCD) is a cyclic oligosaccharide formed by α -D-glucopyranoside
245 units linked 1-4 and bound together in a ring (Riding et al. 2013). This structure is highly soluble
246 (hydrophilic outside), and creates a hydrophilic cavity, capable of forming inclusion complexes
247 with hydrophobic compounds such as organic contaminants (Bardi et al. 2000). In these inclusion
248 complex formations, water molecules are released from the HPCD cavity through displacement by a
249 more hydrophobic molecule in solution (Del Valle 2004). HPCD vary in size (α , β , γ) depending
250 on the number of glucose monomers (generally six to eight units) present in the ring. Generally, an
251 aqueous solution of HPCD is mixed with soil or sediment (1:20 ratio) and then the aqueous phase
252 recovered via centrifugation and the supernatant is analysed for the presence of target contaminants

253 (Cui et al., 2013). Alternatively, the supernatant is discarded and the soil pellet resuspended and
254 extracted using exhaustive solvents (total extraction), and cyclodextrin uptake measured by
255 subtraction (comparing the sum totals of PAHs extracted by HPCD against the total amount
256 extracted by exhaustive solvent) (Papadopoulos et al. 2007). Reid et al. (2000b) showed a reliable
257 prediction of the microbial available concentration of PAHs in soil compared to classical methods
258 based on Soxhlet extraction. Positive correlation has been observed between the amount of HPCD
259 extracted and microbial mineralization in activated carbon-amended soils (Rhodes et al. 2008).
260 Potential of HPCD for indicating bioavailable organic contaminant has been recognized, but no
261 clean-up thresholds for CD-extractable PAHs have been implemented yet (CCME, 2006). Main
262 limitations associated with of HPCD extraction are the reduction of potential for indicating
263 bioavailable fraction in higher organisms (earthworms, benthic invertebrates) (Barthe and Pelletier
264 2007; Hartnik et al., 2008) and the size of the HPCD cavity that might prevent PAHs complexation,
265 owing to steric constraints, which would result in poor extraction efficiencies (Stokes et al., 2005).

266

267 In summary, while passive samplers can be used to estimate the readily-available (pore water
268 concentration) for most soil organisms (Brand et al. 2013), bioavailable concentrations measured by
269 Tenax and HPCD correspond to the sum of the fraction dissolved in pore water and the
270 concentration that could become available on a longer term (i.e. rapid and slow desorbing fraction).
271 Methods to estimate potential bioavailable concentrations can be considered more conservative;
272 therefore we believe that both Tenax and HPCD could be more suitable compared to PSMs for the
273 evaluation of receptor's potential exposure and implementation into the risk assessment. Overall the
274 number of laboratory studies and publications on less exhaustive techniques is promising but they
275 require further efforts to obtain an optimised and enhanced procedure that can be applied across
276 different soil samples and a wider range of contaminants. Such methods could assist in evaluating
277 exposure of ecological receptors and facilitate a more proportionate definition of risk. In addition,
278 these measurements may have implication when establishing remediation endpoints.

279

280 **3 Challenges in assessing complex chemical mixtures bioavailability**

281 Methods to assess bioavailability predominantly focus on assessments carried out on individual
282 substances, or a limited number of substances. Humans and ecological receptors, however, are
283 exposed to a wide variety of chemicals and therefore understanding the potential adverse effects of
284 interactions between these chemicals in mixture is fundamental to assessing risk. We can
285 summarise the challenges of assessing the bioavailability of complex chemical mixtures in three
286 key points (1) standards for mixed-pollutants are absent; (2) combination effects studies on complex
287 chemical mixtures are limited; (3) bioavailability is often neglected.

288 In co-contaminated sites, the presence of both inorganic and oil-derived products, in mixture
289 compositions of near infinite character can produce unpredictable effects (Borgert 2004).
290 Uncertainty in the behaviour of these mixtures reflects not only the complexity of the soil matrix,
291 but also the heterogeneous nature of contaminants in soil, as well as temporal variations in chemical
292 structure and concentration. Conventional risk assessments apply risk-based criteria (guideline
293 values) to deterministic models to make decisions about soil remediation, and establish clean-up
294 standards. However, exposure-risk relationships are established on specific quantitative values
295 (maximum acceptable risk), which can largely vary among different countries (Aqeel et al. 2014),
296 depending on the assumption made when modelling exposure. Whereas conventional exposure
297 assessment relied on the measurable effects of individual chemical species, (de Zwart & Posthuma
298 2005), predictive models for exposure assessment, are unlikely to account for (and interpreted) the
299 combinatorial effects of chemical mixtures (Cornelis et al., 2010). Regulatory frameworks address
300 chemical mixtures based on the contribution of each individual compound present in the mixture, if
301 individual compound toxicity does not exceed the threshold, the overall mixture is considered non-
302 toxic (Heys et al. 2016). Over the last decade, the increasing interest in complex chemical mixtures
303 has been reflected by legislative developments and scientific improvement in understanding of the
304 role of bioavailability of single (Elgh-Dalgren, 2009; Bradham et al., 2015; Chen et al., 2015;
305 Fadaei et al., 2015; Henry et al., 2015; Juhasz et al., 2015; Ortega-Calvo et al., 2015; Stegemeier et
306 al., 2015; Tao et al., 2015) and multiple contaminants (Allan et al., 2012; Gouliarmou and Mayer,
307 2012; Cain et al., 2013; Liu et al., 2013; Kuhn & Maurice., 2014; Amato et al., 2014; Jia et al.,
308 2014; Arp et al., 2014) however, to date, potential risks of combined chemicals are rarely examined
309 in risk assessment (Kienzler et al. 2016).

310

311

312

313 Only a few laboratory based studies have attempted at studying complex chemical mixtures toxic
314 effect on in vitro or biological systems (Environment 2002) more data are required on synergistic
315 and antagonistic interactions of these compounds. While the concepts of mixture toxicity have been
316 discussed for decades, their use has been limited by the absence of toxicological data associated
317 with specific substances, the lack of bioavailability data, and generally the uncertainty associated
318 with knowledge on mixtures of compound. The greatest knowledge gap at the present time is the
319 lack of understanding regarding the mode of action of mixture of compounds which limits the
320 definition of a set of criteria, and therefore requires a careful case-by case approach (EU 2011).

321

322 Cumulative risk assessment (CRA) is a relatively new approach that aims to quantify the health or
323 environmental risk by estimating the level of exposure to multiple contaminants (U.S.
324 Environmental Protection Agency (EPA) 2003). CRA represents a conceptual innovation in the
325 decision making process by moving from a single effect approach to a multiple ecological and
326 human approach to the effects caused by multiple exposure of contaminants (Fox, 2002). Although
327 cumulative risk assessment appear to be pragmatic, few ecotoxicological specific guidelines (e.g.
328 pesticides regulations) account for it (European Environment Agency, 2012). At present, CRA may
329 be the best way to add a health dimension to basic contaminant concentration evaluation. It also
330 might support the decision making process creating a more comprehensive understanding of
331 chemicals behaviour in the environment. Further development and additional studies to verify if
332 CRA is a fair representation of the combine risk for compounds that might not be equal in toxicity,
333 ecotoxicity and chemical behaviour are needed. Ultimately, CRA should not be the only measure of
334 risk, but a valuable support to other analytical tools for investigating environmental risk.

335 Bioavailability is often not well understood and therefore neglected in the risk assessment process.
336 Bioavailability is influenced by a wide range of physicochemical (including both soil and nature of
337 contaminant) (Table 1) and biological factors (Table 2) and it can greatly differ among different
338 organisms, therefore designing suitable one-fit-all extraction approach can be challenging. In order
339 to estimating the bioavailable fraction we should question which of these method provide a good
340 representation for the specific species we intend to protect. Several chemical analytical methods
341 have been developed to assess the bioavailability of inorganic and organic compounds, yet few of
342 them were found to correlate with uptake in model organisms. As such, none of these techniques
343 have been applied to complex chemical mixtures (Muijs and Jonker, 2011a).

344

345 **4 Conclusions**

346 Given the multiple variables affecting the availability of chemicals in soil, we should look at
347 bioavailability not as a fixed value (concentration), but as a dynamic process between an organism
348 and the chemical-uptake over time (ageing). Methods to estimate bioavailability are still not
349 sufficiently cost-effective and standardised. While a great deal of studies and results have been
350 achieved in regards to estimating bioavailability of inorganic contaminants, there are still more
351 work to be done for organic contaminants. Bioavailability and bioaccessibility estimations are seen
352 as useful means to inform human health risk assessment and improving cost-effective management
353 of contaminated land. For instance, when the exceedance of the guideline values is minor,
354 bioaccessibility become the main driver on large site investigations where costs associated with soil
355 removal are considerably high. Nevertheless, lack of information regarding other potential routes of
356 exposure (dermal contact and inhalation) contributes in limiting our confidence in integrating these

357 findings into risk assessment. Similarly, bioavailability is still not fully understood and
358 implemented in existing frameworks, because of both multiple definition across different disciplines
359 and lack of standardised test to measure it. Also, a large number of studies and chemical methods
360 have shown that bioavailable fractions can be positively correlated with uptakes in model organisms
361 and microbial mineralization; obtained data are however inconsistent among different type of
362 contaminants and across different receptors tested. An approach based on weigh of evidence should
363 apply chemical-techniques to measure the bioavailable and bioaccessible fraction, along with
364 biological methods (bioassays) to better understand effects of contaminants uptake and related
365 bioavailability in humans. Bioassays provide a direct measure of contaminant's (bioavailable)
366 concentration for a specific organism over time and are able to quantify and detect a wide range of
367 toxins at relatively low cost.

368

369 Understanding and implementing site-specific bioaccessibility and bioavailability data mean being
370 able to represent more realistically the on-site conditions. Implementation of bioavailability can
371 help the revision of exposure estimate, reducing the cost of remediation, and bringing a greater
372 degree of judgment when assessing risk, and allowing greater levels of contamination left safely in
373 soil. However, in order to provide increased confidence in using bioavailability, further
374 investigation is needed on how to incorporate it into risk assessment. Moreover new approaches are
375 required to tackle the complexity of chemical mixtures and the likely effect of exposure. The
376 challenges are understanding the potential risk connected to a complex chemical mixture, and
377 assessing how the physico-chemical interactions, such as co-solvency, sorption, desorption, and
378 saturation, can affect the potential toxicological response. Understanding which chemicals are
379 effectively bioavailable may be the key for future risk assessment.

380

381 **Acknowledgements:** This work was completed as part of the REMEDIATE (Improved decision-
382 making in contaminated land site investigation and risk assessment) Marie-Curie Innovation
383 Training Network. The network has received funding from the European Union's Horizon 2020
384 Programme for research, technological development and demonstration under grant agreement No.
385 643087.

386

387 **5 References**

388 Abdel-Shafy, H.I. & Mansour, M.S.M., 2016. A review on polycyclic aromatic hydrocarbons:
389 Source, environmental impact, effect on human health and remediation. *Egyptian Journal of*

390 *Petroleum*, 25: 107–123.

391 Agbenin, J.O. & Welp, G., 2012. Bioavailability of copper, cadmium, zinc, and lead in tropical
392 savanna soils assessed by diffusive gradient in thin films (DGT) and ion exchange resin
393 membranes. *Environmental Monitoring and Assessment*, 184: 2275–2284.

394 Ahmadipour, F., Bahramifar, N. & Ghasempouri, S.M., 2014. Fractionation and mobility of
395 cadmium and lead in soils of amol area in iran, using the modified BCR sequential extraction
396 method. *Chemical Speciation and Bioavailability*, 26: 31–36.

397 Alegbeleye, O.O., Opeolu, B.O. & Jackson, V.A., 2017. Polycyclic Aromatic Hydrocarbons: A
398 Critical Review of Environmental Occurrence and Bioremediation. *Environmental*
399 *Management*, 60: 758–783.

400 Alexander, R.R. & Alexander, M., 2000. Bioavailability of genotoxic compounds in soils.
401 *Environmental Science & Technology*, 34: 1589–1593.

402 Allan, S.E., Smith, B.W. & Anderson, K.A., 2012. Impact of the deepwater horizon oil spill on
403 bioavailable polycyclic aromatic hydrocarbons in Gulf of Mexico coastal waters.
404 *Environmental science & technology*, 46: 2033–9.

405 Alloway, B.J., 1995. *Heavy Metals in Soils*, Blackie Academic & Professional, London.

406 Almås, Å.R. et al., 2007. Solubility of major cations and Cu, Zn and Cd in soil extracts of some
407 contaminated agricultural soils near a zinc smelter in Norway: modelling with a multisurface
408 extension of WHAM. *European Journal of Soil Science*, 58: 1074–1086.

409 Almeida, M.I.G.S. et al., 2014. Development of a passive sampler for Zinc(II) in urban pond waters
410 using a polymer inclusion membrane. *Environmental Pollution*, 193: 233–239.

411 Akkanen J., Sormunen, A., Lyytikäinen, M., Leppänen, M., Pehkonen, S., & Kukkonen, J.V.K.,
412 2007. Does Tenax extraction based desorption measure (bio)availability of sediment-
413 associated contaminants? available at [www.ices.dk/sites/pub/CM%20Documents/CM-](http://www.ices.dk/sites/pub/CM%20Documents/CM-2007/J/J1407.pdf)
414 [2007/J/J1407.pdf](http://www.ices.dk/sites/pub/CM%20Documents/CM-2007/J/J1407.pdf)

415 Alvarez, M.B., Domini, C.E., Garrido, M., Lista, A.G., Fernández-Band B.S., 2011. Single-step
416 chemical extraction procedures and chemometrics for assessment of heavy metal behaviour in
417 sediment samples from the Bahía Blanca estuary, Argentina. *Journal of Soils and Sediments*,
418 11: 657–666.

419 Amacher, M.C., 1984. Determination of Ionic Activities in Soil Solutions and Suspensions:
420 Principal Limitations1. *Soil Science Society of America Journal*, 48: 519–524.

421 Amato, E.D., Simpson, S.L., Jarolimen, C.V., & Jolley, D.F., 2014. Diffusive gradients in thin films
422 technique provide robust prediction of metal bioavailability and toxicity in estuarine
423 sediments. *Environmental science & technology*, 48: 4485–94.

424 Amiard-Triquet, C., Amiard, J.C. & Mouneyrac, C., 2015. *Aquatic Ecotoxicology: Advancing Tools*

425 for *Dealing with Emerging Risks*, Elsevier Science, Academic Press, 518 pp.

426 An, J., Jho, E.H. & Nam, K., 2015. Effect of dissolved humic acid on the Pb bioavailability in soil
427 solution and its consequence on ecological risk. *Journal of Hazardous Materials*, 286: 236–
428 241.

429 An, X., Xiao, B., Di, X., Dong, H., & Tang, H., 2017. Research progress on aging of organic
430 pollutants in geosorbents: a review. *Acta Geochimica*, 36: 27–43.

431 Apte, S.C. & Batley, G.E., 1995. *Trace metal speciation of labile chemical species in natural*
432 *waters and sediments: non-electrochemical approaches*, John Wiley and Sons.

433 Aqeel, M., Jamil, M. & Yusoff, I., 2014. Soil Contamination, Risk Assessment and Remediation. In
434 *Environmental Risk Assessment of Soil Contamination*. InTech, pp. 3–56.

435 Arp, H.P.H., Lundstedt, S., Josefsson, S., Cornelissen, G., Enell, A., Allard, A.S., Kleja, D.B.,
436 2014. Native oxy-PAHs, N-PACs, and PAHs in historically contaminated soils from Sweden,
437 Belgium, and France: their soil-porewater partitioning behavior, bioaccumulation in
438 *Enchytraeus crypticus*, and bioavailability. *Environmental science & technology*, 48:11187–
439 11195.

440 Ashraf, M.A., Maah, M.J. & Yusoff, I., 2012. Chemical Speciation and Potential Mobility of Heavy
441 Metals in the Soil of Former Tin Mining Catchment. *The Scientific World Journal*, 12: 1–11.

442 Atlas, R.M., 1995. Bioremediation of petroleum pollutants. *International Biodeterioration and*
443 *Biodegradation* 35: 317-327

444 Ayres, R.U., 1992. Toxic heavy metals: materials cycle optimization. *Proceedings of the National*
445 *Academy of Sciences of the United States of America*, 89: 815–820.

446 Baran, A., Tarnawski, M. & Michalec, B., 2015. Assessment of metal leachability and toxicity from
447 sediment potentially stored on land. *Water SA*, 41: 606–613.

448 Bardi, L., Mattei, A., Steffan, S., & Marzona, M., 2000. Hydrocarbon degradation by a soil
449 microbial population with beta-cyclodextrin as surfactant to enhance bioavailability. *Enzyme*
450 *and microbial technology*, 27: 709–713.

451 Barthe, M. & Pelletier, É., 2007. Comparing bulk extraction methods for chemically available
452 polycyclic aromatic hydrocarbons with bioaccumulation in worms. *Environmental Chemistry*,
453 4: 271-283.

454 Bergknut, M., Sehlin, E., Lunstedt, S., Andersson, P.L., Haglund, P., & Tysklind, M., 2007.
455 Comparison of techniques for estimating PAH bioavailability : Uptake in *Eisenia fetida* ,
456 passive samplers and leaching using various solvents and additives. *Environmental Pollution*,
457 145: 154–160.

458 Beriro, D.J., Cave, M.R., Wragg, J., Thomas, R., Willa, G., & Evans, F. 2016. A review of the
459 current state of the art of physiologically-based tests for measuring human dermal in vitro

460 bioavailability of polycyclic aromatic hydrocarbons (PAH) in soil. *Journal of Hazardous*
461 *Materials*, 305: 240–259.

462 Berkowitz, B., Dror, I. & Yaron, B., 2008. Contaminant Geochemistry: Interactions and Transport
463 in the Subsurface Environment, Springer Berlin Heidelberg. 557 pp

464 Berthelot, Y., Valton, E., Auroy, A., & Robidoux, P.Y., 2008. Integration of toxicological and
465 chemical tools to assess the bioavailability of metals and energetic compounds in contaminated
466 soils. *Chemosphere*, 74: 166–177.

467 Biswas, B., Sarkar, B., Mandal, A., & Naidu, R. 2015. Heavy metal-immobilizing organoclay
468 facilitates polycyclic aromatic hydrocarbon biodegradation in mixed-contaminated soil.
469 *Journal of Hazardous Materials*, 298: 129–137.

470 Bolan, N., Kunhikrsihnan, A., Thangarajan, R., Kumpiene, J., Park, J., Makino, T., Kirkham, M.B.,
471 & Scheckel, K. 2014. Remediation of heavy metal(loid)s contaminated soils - To mobilize or
472 to immobilize? *Journal of Hazardous Materials*, 266: 141–166.

473 Borgert, C.J., Quill, T.F., McCarty, L.S., Mason, A.M. 2004. Can mode of action predict mixture
474 toxicity for risk assessment? *Toxicol Appl Pharmacol*. 201: 85-96.

475 Bradham, K.D., Nelson, C., Juhasz, A., Smith, E., Scheckel, K., Obenour, D.R., Bradley, W.M., &
476 Thomas, D.J. 2015. Independent data validation of an in vitro method for the prediction of the
477 relative bioavailability of arsenic in contaminated soils. *Environmental science & technology*,
478 49: 6312–6318.

479 Braida, W.J., White, J.C. & Pignatello, J.J., 2004. Indices for bioavailability and biotransformation
480 potential of contaminants in soils. *Environmental toxicology and chemistry*, 23: 1585–1591.

481 Brand, E. et al., 2012. Advice on implementing bioavailability in the Dutch soil policy framework -
482 User protocols for organic contaminants. RIVM Report 711701102/2012. , p.73. Available at:
483 <http://www.rivm.nl/bibliotheek/rapporten/711701102.pdf>.

484 Brand, E., Lijzen, J., Peijnenburg, W., & Swartjes, F., 2013. Possibilities of implementation of
485 bioavailability methods for organic contaminants in the Dutch Soil Quality Assessment
486 Framework. *Journal of Hazardous Materials*, 261: 833–839.

487 Brand, E., Peijnenburg, W., Goenenberg, B., Vink, J., ten Hulscher, D., Jonker, C., Romkens, P., &
488 Roex, E., 2009. *Towards implementation of bioavailability measurements in the Dutch*
489 *regulatory framework*, BA Bilthoven, the Netherlands.

490 Brassington K., Pollard S.J.T., Coulon F. 2010. Weathered hydrocarbon biotransformation: implications for
491 bioremediation, analysis and risk assessment. Chapter 5. In: *Handbook of Microbiology of*
492 *Hydrocarbons, Oils, Lipids*. Volume 4. Timmis K. ed., Springer Verlag Berlin Heidelberg, Humana
493 Press, [ISBN: 978-3-540-77584-3](https://doi.org/10.1007/978-3-540-77584-3); pp 2487-2501

494 Burgess, R.M. & Driscoll, S.B.K., 2016. P User's Manual (rocedures for Using Passive Sampling in

495 the Evaluation of Contaminated Sediments. U.S. Environmental Protection Agency,
496 Washington, DC, EPA/600/R-16/357, 2016.

497 Cachada, A., Pereira, R., da Silva, E.F., & Duarte, A.C., 2014. The prediction of PAHs
498 bioavailability in soils using chemical methods: state of the art and future challenges. *The*
499 *Science of the total environment*, 472: 463–80.

500 Cain, D.J., Croteau, M.-N. & Fuller, C.C., 2013. Dietary bioavailability of Cu adsorbed to colloidal
501 hydrous ferric oxide. *Environmental science & technology*, 47: 2869–76.

502 Cajthaml, T. & Václav, S., 2005. Application of Supercritical Fluid Extraction (SFE) to Predict
503 Bioremediation Efficacy of Long-Term Composting of PAH-Contaminated Soil. 39: 8448–
504 8452.

505 Canadian Council of Ministers of the Environment (CCME), 2006. *Five-year review of the Canada-*
506 *Wide Standards for Petroleum Hydrocarbons (PHC CWS): Ecological , Direct Soil Contact*
507 *Guidance*,

508 Cantwell, F.F., Nielsen, J.S. & Hrudey, S.E., 1982. Free nickel ion concentration in sewage by an
509 ion exchange column-equilibration method. *Analytical Chemistry*, 54: 1498–1503.

510 Capri, E., Trevisan, M., Jantunen, P.K., 2004. Computer models for characterizing the fate of
511 chemical in soil:Pesticide leaching models and their practical application, Chapter 21. In *Soil-*
512 *Water-Solute Process Characterization: An Integrated Approach*. edited by Alvarez-Benedi J
513 and Munoz-Carpena R., CRC Press 2004, pp. 715-756.

514 Cappuyns, V., 2012. A Critical Evaluation of Single Extractions from the SMT Program to
515 Determine Trace Element Mobility in Sediments. *Applied and Environmental Soil Science*,
516 2012, Article ID 672914, 15 pages

517 Cave, M., Wragg, J., Gowing, C., & Gardner, A. 2015. Measuring the solid-phase fractionation of
518 lead in urban and rural soils using a combination of geochemical survey data and chemical
519 extractions. *Environmental Geochemistry and Health*, 37: 779–790.

520 Cave, M.R., Milodowski, A.E. & Friel, E.N., 2004. Evaluation of a method for identification of host
521 physico-chemical phases for trace metals and measurement of their solid-phase partitioning in
522 soil samples by nitric acid extraction and chemometric mixture resolution. *Geochemistry:*
523 *Exploration, environment, analysis*, 4: 71–86.

524 Cecchin, I., Reginatto, C., & Thomé, A., 2016. Influence of Physicochemical Factors on Biodiesel
525 Retention in Clayey Residual Soil. *Journal of Environmental Engineering*, 142: 4015093.

526 Chen, Z., Zhang, Y., Gao, Y., Boyd, S.A., Zhu, D., & Li, H., 2015. Influence of Dissolved Organic
527 Matter on Tetracycline Bioavailability to an Antibiotic-Resistant Bacterium. *Environmental*
528 *science & technology*, 49: pp.10903–10910.

529 Chigbo, C., Batty, L. & Bartlett, R., 2013. Interactions of copper and pyrene on phytoremediation

530 potential of *Brassica juncea* in copper–pyrene co-contaminated soil. *Chemosphere*, 90(10),
531 pp.2542–2548. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0045653512013549>.

532 Christensen, J.B. & Christensen, T.H., 1999. Complexation of Cd, Ni, and Zn by DOC in polluted
533 groundwater: A comparison of approaches using resin exchange, aquifer material sorption, and
534 computer speciation models (WHAM and MINTEQA2). *Environmental Science and
535 Technology*, 33: 3857–3863.

536 Chung, N. & Alexander, M., 1998. Differences in Sequestration and Bioavailability of Organic
537 Compounds Aged in Dissimilar Soils. *Environmental Science & Technology*, 32: 855–860.

538 Cipullo, S., Brassington, K., Pollard, S.J.T., & Coulon, F., 2017. Weathered hydrocarbons
539 biotransformation: implications for bioremediation, analysis and risk assessment. In K.N.
540 Timmis ed., Springer Verlag Berlin Heidelberg, ed. *Handbook of Microbiology of
541 Hydrocarbons, Oils, Lipids*. Humana Press.

542 Ciszewski, D. & Grygar, T.M., 2016. A Review of Flood-Related Storage and Remobilization of
543 Heavy Metal Pollutants in River Systems. *Water, Air, and Soil Pollution*, 227: 239.

544 Cornelis A. M. van Gestel; Martijs Jonker; Jan E. Kammenga, R.L.C.S., 2010. *Mixture Toxicity:
545 Linking Approaches from Ecological and Human Toxicology*, CRC Press, 320 pages.

546 Cornelissen, G., Rigterink, H., ten Huylscher, D.E., Vrind, B.A., & van Noort, P.C., 2001. A
547 simple Tenax extraction method to determine the availability of sediment-sorbed organic
548 compounds. *Environmental toxicology and chemistry* 20: 706–711.

549 Coulon, F., Al Awadi, M., Cowie, W., Mardlin, D., Pollard, S., Cunningham, C., Risdon, G.,
550 Arthur, P., Semple, K.T., & Paton G.I. 2010. When is a soil remediated? Comparison of
551 biopiled and windrowed soils contaminated with bunker-fuel in a full-scale trial.
552 *Environmental Pollution*, 158: 3032–3040.

553 Cornelissen, G., Rigterink, H., Ferdinandy, M.M.A. & Noort, P.C.M.V.A.N., 1998. Rapidly
554 Desorbing Fractions of PAHs in Contaminated Sediments as a Predictor of the Extent of
555 Bioremediation. *Environmental Science and Technology*, 32: 966–970.

556 Coulthard, T.J. & Macklin, M.G., 2003. Modeling long-term contamination in river systems from
557 historical metal mining. *Geology*, 31: 451–454.

558 Cox, S.F., Chelliah, M.C.M., McKinley, J.M., Palmer, M., Ofterdinger, U., Young, M.E., Cave, M.,
559 & Wragg, J. 2013. The importance of solid-phase distribution on the oral bioaccessibility of Ni
560 and Cr in soils overlying Palaeogene basalt lavas, Northern Ireland. *Environmental
561 Geochemistry and Health*, 35: 553–567.

562 Cui, X., Mayer, P. & Gan, J., 2013. Methods to assess bioavailability of hydrophobic organic
563 contaminants: Principles, operations, and limitations. *Environmental Pollution*, 172: 223–234.

564 Cuypers, C., Pancras, T., Grotenhuis, T., & Rulkens, W., 2002. The estimation of PAH

565 bioavailability in contaminated sediments using hydroxypropyl- β -cyclodextrin and Triton X-
566 100 extraction techniques. *Chemosphere*, 46: 1235–1245.

567 Davidson, I.E. & Smyth, W.F., 1979. Cathodic stripping voltammetric determination of organic
568 halides in drug dissolution studies. *Analytical Chemistry*, 51: 2127–2133.

569 Davlson, W. & Zhang, H., 1994. In situspeciation measurements of trace components in natural
570 waters using thin-film gels. *Nature*, 367: 546–548.

571 Dean, J.R. & Scott, W.C., 2004. Recent developments in assessing the bioavailability of persistent
572 organic pollutants in the environment. *TrAC - Trends in Analytical Chemistry*, 23: 609–618.

573 Del Valle, E.M.M., 2004. Cyclodextrins and their uses: A review. *Process Biochemistry*, 39: 1033–
574 1046.

575 De la Cal, A., Eljarrat, E., Gorenhuis, T., & Barcelo, D. 2008. Tenax® Extraction As a Tool To
576 Evaluate the Availability of Polybrominated Diphenyl Ethers, ddt, and ddt Metabolites in
577 Sediments. *Environmental Toxicology and Chemistry*, 27: 1250–1256.

578 du Bray, E.A., 1995. *Preliminary compilation of descriptive geoenvironmental mineral deposit*
579 *models*, US Geological Survey, available at <https://pubs.usgs.gov/of/1995/0831/report.pdf>

580 Domínguez, J.B., 2008. *Soil Contamination Research Trends*, Nova Science Publishers.

581 Driscoll, S.K. & Thompson, T., 2016. RPM Guide Integrating Passive Sampling Methods into
582 Management of Contaminated Sediment Sites : A Guide for Department of Defense Remedial
583 Project Managers.

584 Duan, L., Naidu, R., Liu, Y., Palanisami, T., Dong, Z., Mallavarapu., M., & Semple, K.T., 2015.
585 Effect of ageing on benzo[a]pyrene extractability in contrasting soils. *Journal of Hazardous*
586 *Materials*, 296: 175–184.

587 Duan, L., Palanisami, T., Liu, Y., Dong, Z., Mallavarapu, M., Kuchel, T., Semple, K.T., & Naidu,
588 R., 2014. Effects of ageing and soil properties on the oral bioavailability of benzo[a]pyrene
589 using a swine model. *Environment International*, 70: 192–202.

590 Duan, L. & Naidu, R., 2013. Effect of ionic strength and index cation on the sorption of
591 phenanthrene. *Water, Air, and Soil Pollution*, 224: 1–17.

592 Dube, A., Zbytniewski, R, Kowalkowski, T., Cukrowska, E., & Buszewski, B., 2001. Adsorption
593 and Migration of Heavy Metals in Soil. *Polish Journal of Environmental Studies*, 10: 1–10.

594 Dyrtrtova, J.J., Šestáková I., Jakl, M., Száková, J., Miholová, D., & Tlustoš, P., 2008. The use of
595 differential pulse anodic stripping voltammetry and diffusive gradient in thin films for heavy metals
596 speciation in soil solution. *Central European Journal of Chemistry*, 6: 71–79.

597 ECETOC, 2014. ECETOC Annual Report 2013. Available at: <http://www.mac.org.my/v3/wp->

598 content/uploads/2014/05/Annual-Report-2013_A4_FA_22052013-web.pdf.

599 Eggleton, J. & Thomas, K. V., 2004. A review of factors affecting the release and bioavailability of
600 contaminants during sediment disturbance events. *Environment International*, 30: 973–980.

601 Elgh-Dalgren, K., 2009. *Remediation of Materials with Mixed Contaminants. Treatability,*
602 *Technology and Final Disposal.* Orebro University. Available at:
603 <http://www.publications.oru.se/>.

604 Elliott, H.A., Dempsey, B.A. & Maille, P.J., 1990. Content and Fractionation of Heavy Metals in
605 Water Treatment Sludges. *Journal of Environment Quality*, 19: 330–334.

606 van Elsas, J.D. et al., 2006. *Modern soil microbiology*, 2nd Edition, CRC Press. 672 pages

607 Environment Agency, 2006. Remedial Target Methodology; Hydrogeological Risk Assessment for
608 Land Contamination. *Environment Agency*, p.25.

609 Environment Agency, 2010. *Verification of remediation of land contamination: SC030114/R1*,
610 Available at: www.environmental-agency.gov.uk. (accessed on 29 September 2017)

611 Environmental Agency, 2008. *Assessment of Metal Mining-Contaminated River Sediments in*
612 *England and Wales, Science Report SC0301/SR4*, Bristol, England

613 European Environment Agency, 2012. Overview of contaminants affecting soil and groundwater in
614 Europe. *Published 12 Nov 2009*. Available at: [http://www.eea.europa.eu/data-and-](http://www.eea.europa.eu/data-and-maps/figures/overview-of-contaminants-affecting-soil-and-groundwater-in-europe)
615 [maps/figures/overview-of-contaminants-affecting-soil-and-groundwater-in-europe](http://www.eea.europa.eu/data-and-maps/figures/overview-of-contaminants-affecting-soil-and-groundwater-in-europe) (Accessed
616 on 8 December 2015].

617 Fadaei, H., Watson, A., Place, A., Connolly, J., & Ghosh, U., 2015. Effect of PCB Bioavailability
618 Changes in Sediments on Bioaccumulation in Fish. *Environmental science & technology*, 49:
619 12405–12413.

620 Fantke, P., Arnot, J.A. & Doucette, W.J., 2016. Improving plant bioaccumulation science through
621 consistent reporting of experimental data. *Journal of Environmental Management*, 181: 374–
622 384.

623 Farmer, W.J., 1997. Role of Microstructural Properties in the Time-Dependent Sorption /
624 Desorption Behavior of 1, 2-Dichloroethane on Humic Substances. *Environmental Science &*
625 *Technology*, 31: 2520–2526.

626 Fernández-Ondoño, E., Bacchetta, G., Lallella A.M., Navarro, F.B., Ortiz, I., & Jimenez, M.N.
627 2017. Use of BCR sequential extraction procedures for soils and plant metal transfer
628 predictions in contaminated mine tailings in Sardinia. *Journal of Geochemical Exploration*,
629 172: 133–141.

630 Fotovat, A. & Naidu, R., 1997. Ion exchange resin and MINTEQA2 speciation of Zn and Cu in

631 alkaline sodic and acidic soil extracts. *Australian Journal of Soil Research*, 35: 711–726.

632 Fox M.A. 2002. Evaluating cumulative risk assessment for environmental justice: a community
633 case study. *Environ Health Perspect.* 110: 203–209.

634 Gál, J., Hursthouse, A.S. & Cuthbert, S.J., 2006. Chemical availability of arsenic and antimony in
635 industrial soils. *Environmental Chemistry Letters*, 3: 149–153.

636 Gauthier, P.T., Norwood, W.P., Prepas, E.E., & Pyle, G.G., 2014. Metal-PAH mixtures in the
637 aquatic environment: A review of co-toxic mechanisms leading to more-than-additive
638 outcomes. *Aquatic Toxicology*, 154: 253–269.

639 Ge, Y., Sauvé, S. & Hendershot, W.H., 2005. Equilibrium Speciation of Cadmium, Copper, and
640 Lead in Soil Solutions. *Communications in Soil Science and Plant Analysis*, 36: 1537–1556.

641 Ghosal, D., Ghosh, S., Dutta, T.K., & Ahn, Y. 2016. Current State of Knowledge in Microbial
642 Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Frontiers in
643 Microbiology*, 7, 1369.

644 Giacomino, A., Abollino, O., Malandrino, M., & Mentasti, E., 2011. The role of chemometrics in
645 single and sequential extraction assays: A Review. Part II. Cluster analysis, multiple linear
646 regression, mixture resolution, experimental design and other techniques. *Analytica Chimica
647 Acta*, 688: 122–139.

648 Gomez-Eyles, J.L., Jonkers, M.T.O., Hodson, M.E., Collins, C.D., 2011. Passive samplers provide
649 a better prediction of PAH bioaccumulation in earthworms and plant roots than exhaustive,
650 mild solvent, and cyclodextrin extractions. *Environmental science & technology*, 46: 962–969.

651 Gouliarmou, V. & Mayer, P., 2012. Sorptive bioaccessibility extraction (SBE) of soils: combining a
652 mobilization medium with an absorption sink. *Environmental science & technology*, 46:
653 10682–10689.

654 Groenenberg, J.E., Römkens, P.F.A.M., Van Zomeren A., Rodrigues, S., & Comans R.B.J., 2017.
655 Evaluation of the Single Dilute (0.43 M) Nitric Acid Extraction to Determine Geochemically
656 Reactive Elements in Soil. *Environmental Science and Technology*, 51: 2246–2253.

657 Gupta, S.K. & Aten, C., 1993. Comparison and Evaluation of Extraction Media and Their
658 Suitability in a Simple Model to Predict the Biological Relevance of Heavy Metal
659 Concentrations in Contaminated Soils. *International Journal of Environmental Analytical
660 Chemistry*, 51: 25–46.

661 Hallgren, P., Westborn, R., Nilsson, T., Sporring S., & Björklund, E., 2006. Measuring
662 bioavailability of polychlorinated biphenyls in soil to earthworms using selective supercritical
663 fluid extraction. *Chemosphere*, 63: 1532–1538.

664 Hartnik, T., Jensen, J. & Hermens, J.L.M., 2008. Nonexhaustive beta-cyclodextrin extraction as a
665 chemical tool to estimate bioavailability of hydrophobic pesticides for earthworms.

666 *Environmental science & technology*, 42: 8419–25.

667 Harmsen J, Naidu R (2013). Bioavailability as a tool in site management. *J. Hazard. Mat* 261: 840-
668 846.

669 Harwood, A.D., Landrum, P.F. & Lydy, M.J., 2012. Can SPME fiber and tenax methods predict the
670 bioavailability of biotransformed insecticides? *Environmental Science and Technology*, 46:
671 2413–2419.

672 Hawthorne, S.B. et al., 2000. Comparisons of Soxhlet extraction , pressurized liquid extraction ,
673 supercritical fluid extraction and subcritical water extraction for environmental solids :
674 recovery , selectivity and effects on sample matrix. , 892: 421–433.

675 Heng, L.-S., Wang, D.Z., Jiang, X., Rao, W., Zhang, W.H., Guo, C.Y., & Li, T. 2010. Relationship
676 between Fe, Al oxides and stable organic carbon, nitrogen in the yellow-brown soils. *Huan*
677 *Jing Ke Xue* 31: 2748–55.

678 Henry, H., Naujokas, M.F., Attanyake, C., Basta, N.T., Cheng, Z., Hettiarachchi, G.M., Maddoloni,
679 M., Schadt, C., & Scheckel, K.G., 2015. Bioavailability-Based In Situ Remediation To Meet
680 Future Lead (Pb) Standards in Urban Soils and Gardens. *Environmental science & technology*,
681 49: 8948–58.

682 Heys, K.A., Shore, R.F., Gloira, M., Jones, K.C, & Martin, F.L. 2016. Risk assessment of
683 environmental mixture effects. *RSC Advances*, 53: 47844–47857.

684 Holm, P.E. et al., 1995. Speciation of Cadmium and Zinc with Application to Soil Solutions.
685 *Journal of Environmental Quality*, 24:183–190.

686 Hooda, P.S., Zhang, H., Davidson, W., & Edwards, A.C., 1999. Measuring bioavailable trace
687 metals by diffusive gradients in thin films (DGT): soil moisture effects on its performance in
688 soils. *European Journal of Soil Science*, 50: 285–294.

689 Houba, V.J.G. et al., 2000. Soil analysis procedures using 0.01 M calcium chloride as extraction
690 reagent. *Communications in Soil Science and Plant Analysis*, 31: 1299–1396.

691 Houben, D., Evrard, L. & Sonnet, P., 2013. Beneficial effects of biochar application to
692 contaminated soils on the bioavailability of Cd, Pb and Zn and the biomass production of
693 rapeseed (*Brassica napus* L.). *Biomass and Bioenergy*, 57: 196–204.

694 Huang, Z.Y., Xie, H., Cao Y-L., Cai, C., & Zhang, Z. 2014. Assessing of distribution, mobility and
695 bioavailability of exogenous Pb in agricultural soils using isotopic labeling method coupled
696 with BCR approach. *Journal of Hazardous Materials*, 266: 182–188.

697 Hughes, J.C. & Noble, A.D., 1991. Extraction of chromium, nickel and iron and the availability of
698 chromium and nickel to plants from some serpentinite-derived soils from the eastern Transvaal
699 as revealed by various single and sequential extraction techniques. *Communications in Soil*
700 *Science and Plant Analysis*, 22: 1753–1766.

701 Hulscher, T.E.M.T., Postma, J., den Besten P.J., Stroomberg, G.J., Belfroid, A., Wegener, J.W.,
702 Faber, J.H., van der Pol, J.J., Hendricks, A.J., van Noort, P.C. 2003. Tenax extraction mimics
703 benthic and terrestrial bioavailability of organic compounds. *Environmental Toxicology and*
704 *Chemistry*, 22: 2258–2265.

705 Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B.B., & Beeregowda, K.N., 2014. Toxicity,
706 mechanism and health effects of some heavy metals. *Interdisciplinary Toxicology*, 7: 60–72.

707 Jia, F., Bao, L.-J., Crago, J., Schlenk, D., & Gan J., 2014. Use of isotope dilution method to predict
708 bioavailability of organic pollutants in historically contaminated sediments. *Environmental*
709 *science & technology*, 48: 7966–73.

710 Jiang, B., Su, D., Wang, X., Liu, J., & Ma, Y., 2017. Field evidence of decreased extractability of
711 copper and nickel added to soils in 6-year field experiments. *Frontiers of Environmental*
712 *Science & Engineering*, 12: 7.

713 Jiang, Y., Brassington, K., Prpich, G., Paton, G.I., Semple, K.T., Pollard, S.J.T., Coulon, F., 2016.
714 Insights into the biodegradation of weathered hydrocarbons in contaminated soils by
715 bioaugmentation and nutrient stimulation. *Chemosphere*, 161: 300–307.

716 John, D.A. & Leventhal, J.S., 1995. Bioavailability of metals. *United States Geological Survey*, 2,
717 pp.10–18. Available at: <http://eprints.utas.edu.au/4774/>.

718 Johnson, D.L., Jones, K.C., Langdon, C.J., Pearce, T.G., & Semple, K.T., 2002. Temporal changes
719 in earthworm availability and extractability of polycyclic aromatic hydrocarbons in soil. *Soil*
720 *Biology and Chemistry*, 34, 1363–1370.

721 Jonker, M.T.O., van der Heijden, S.A., Kreitinger, J.P., & Hawthorne, S.B., 2007. Predicting PAH
722 bioaccumulation and toxicity in earthworms exposed to manufactured gas plant soils with
723 solid-phase microextraction. *Environmental Science and Technology*, 41: 7472–7478.

724 Jørgensen, S., 2000. *Principles of Pollution Abatement*, 1st Edition, Elsevier Science. 532 pages.

725 Juhasz, A.L., Weber, J., Stevenson, G., Slee, D., Gancarz, D., Rofe, A., & Smith, E., 2014. In vivo
726 measurement, in vitro estimation and fugacity prediction of PAH bioavailability in post-
727 remediated creosote-contaminated soil. *Science of The Total Environment*, 473: 147–154.

728 Juhasz, A.L., Herde, P., Herde, C., Boland, J., & Smith, E., 2015. Predicting Arsenic Relative
729 Bioavailability Using Multiple in Vitro Assays: Validation of in Vivo-in Vitro Correlations.
730 *Environmental science & technology*, 49: 11167–11175.

731 Kaakinen, J., Kuokkanen, T., Leskinene, H., Välimäki, I., & Kujala, K., 2015. The use of a four-
732 stage sequential leaching procedure and the corresponding one-phase extractions for risk
733 assessment of potential harmful substances in waste rock utilized in railway ballast. *Chemical*
734 *Speciation and Bioavailability*, 27: 71–80.

735 Kelsey, J.W. & Alexander, M., 1997. Declining bioavailability and inappropriate estimation of risk

736 of persistent compounds. , 16: 582–585.

737 Kelsey, J.W., Kottler, B.D. & Alexander, M., 1997. Selective chemical extractants to predict
738 bioavailability of soil-aged organic chemicals. *Environmental Science and Technology*, 31:
739 214–217.

740 Khan, A., Khan, S., Khan, M.A., Qamar, Z., & Waqas, M., 2015. The uptake and bioaccumulation
741 of heavy metals by food plants, their effects on plants nutrients, and associated health risk: a
742 review. *Environmental Science and Pollution Research*, 22: 13772–13799.

743 Khan, S., Kazi, T.G., Arain, M.B., Kolachi, N.F., Baig, J.A., Afridid, H., I., Shah, A.Q., 2013.
744 Evaluation of bioavailability and partitioning of aluminum in sediment samples of different
745 ecosystems by modified sequential extraction methods. *Clean - Soil, Air, Water*, 41: 808–815.

746 Kienzler, A., Bopp, S.K., van der Linden, S., Berggren, E., & Worth A., 2016. Regulatory
747 assessment of chemical mixtures: Requirements, current approaches and future perspectives.
748 *Regulatory Toxicology and Pharmacology*, 80: 321–334.

749 Kim, R.Y., Yoon, J.K., Kim, T.S., Yang, J.E., Owens, G., Kim, K.R., 2015. Bioavailability of
750 heavy metals in soils: definitions and practical implementation—a critical review.
751 *Environmental Geochemistry and Health*, 37: 1041–1061.

752 Kleber, M., Sollins, P. & Sutton, R., 2007. A conceptual model of organo-mineral interactions in
753 soils: Self-assembly of organic molecular fragments into zonal structures on mineral surfaces.
754 *Biogeochemistry*, 85: 9–24.

755 Koopmans, G.F. et al., 2008. Influence of EDDS on metal speciation in soil extracts: Measurement
756 and mechanistic multicomponent modeling. *Environmental Science and Technology*, 42:
757 1123–1130.

758 Krishnamurti, G.S.R., Subashchandrabose, S.R., Megharaj, M., Naidu, R., 2015. Assessment of
759 bioavailability of heavy metal pollutants using soil isolates of *Chlorella* sp. *Environmental*
760 *Science and Pollution Research*, 22: 8826–8832.

761 Krishnamurti, G.S.R., Huang, P.M., van Ressel, K.J.C., Kozak, L.M., & Rostad, H.P.W., 1995.
762 Speciation of particulate-bound Cadmium of soils and its bioavailability. *The Analyst*, 120:
763 659.

764 Krull, E., Baldock, J. & Skjemstad, J., 2001. Soil Texture Effects on Decomposition and Soil
765 Carbon Storage. *NEE Workshop Proceedings*, pp.103–110.

766 Kuhn, K.M., & Maurice, P.A., 2014. Accessibility of humic-associated Fe to a microbial
767 siderophore: implications for bioavailability. *Environmental science & technology*, 48: 1015–
768 1022.

769 Nikel et al., 2013. Why are chlorinated pollutants so difficult to degrade aerobically? Redox stress
770 limits 1,3-dichloroprop-1-ene metabolism by *Pseudomonas pavonaceae*. *Philos Trans R Soc*

771 Lond B Biol Sci. 368: 20120377.

772 Sormunen, A.J., Leppänen, M.T., Kukkonen, J.V., 2009. Ecotoxicology and Environmental Safety
773 Examining the role of temperature and sediment – chemical contact time on desorption and
774 bioavailability of sediment-associated tetrabromo diphenyl ether and benzo (a) pyrene.
775 *Ecotoxicology and Environmental Safety*, 72, 1234–1241.

776 Kunhikrishnan, A. et al., 2016. Redox Reactions of Heavy Metal (loid) s in Soils and Sediments in
777 Relation to Bioavailability and Remediation. Chapter 3. In: *Trace Elements in Waterlogged*
778 *Soils and Sediments.*, CRC Press, 27-52

779 Kuppusamy, S., Thavamani, P., Venkateswarly, K., Lee, Y.B., Naidu, R., Megharaj, M., 2017.
780 Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils:
781 Technological constraints, emerging trends and future directions. *Chemosphere*, 168: 944–968.

782 Laak, T.L. ter., Agbo, S.O., Barendregt, A., & Hermens, L.M., 2006. Freely Dissolved
783 Concentrations of PAHs in Soil Pore Water : Measurements via Solid-Phase Extraction and
784 Consequences for Soil Tests. *Environmental Science & Technology* 40: 1307–1313.

785 Laak, T.L. ter, Barendregt, A. & Hermens, J.L.M., 2006. Freely dissolved pore water concentrations
786 and sorption coefficients of PAHs in spiked, aged, and field-contaminated soils. *Environmental*
787 *Science & Technology*, 40: 2184–2190.

788 Lal, V., Peng, C. & Ng, J., 2015. A review of non-exhaustive chemical and bioavailability methods
789 for the assessment of polycyclic aromatic hydrocarbons in soil. *Environmental Technology &*
790 *Innovation*, 4: 159–167.

791 Latawiec, A.E., Swindell, A.L. & Reid, B.J., 2008. Environmentally friendly assessment of organic
792 compound bioaccessibility using sub-critical water. *Environmental Pollution*, 156: 467–473..

793 Leggett, G.E. & Argyle, D.P., 1983. The DTPA-extractable iron, manganese, copper, and zinc from
794 neutral and calcareous soils dried under different conditions. *Soil Science Society of America*
795 *Journal*, 47: 518–522.

796 Lepp, N.W., 2012. *Effect of Heavy Metal Pollution on Plants: Metals in the Environment*, Springer
797 Netherlands, Springer, 249 pp

798 Leppä, M.T. & Kukkonen, J.V.K., 2006. Evaluating the role of desorption in bioavailability of
799 sediment-associated contaminants using oligochaetes , semipermeable membrane devices and
800 Tenax extraction. *Environmental Pollution*, 140:150–163.

801 Leppänen, M.T., Landrum, P.F., Kukkonen J.V.K., Greenberg, M.S., Burton Jr, G.A., Robinson,
802 S.D., & Gossiaux, D.C., 2003. Investigating the role of desorption on the bioavailability of
803 sediment-associated 3,4,3',4'-tetrachlorobiphenyl in benthic invertebrates. *Environmental*
804 *toxicology and chemistry*, 22: 2861–2871.

805 Liang, S., Guan, D.X., Ren, J.H., Zhang, M., Luo, J., & Ma, L.Q., 2014. Effect of aging on arsenic

806 and lead fractionation and availability in soils: Coupling sequential extractions with diffusive
807 gradients in thin-films technique. *Journal of Hazardous Materials*, 273: 272–279.

808 Librando, V. & Aresta, M., 2004. Supercritical fluid extraction of polycyclic aromatic hydrocarbons
809 from marine sediments and soil samples Supercritical fluid extraction of polycyclic aromatic
810 hydrocarbons from marine. *Chemosphere*, 54: 1189-1197.

811 Lindsay, W.L. & Norvell, W.A., 1978. Development of a DTPA soil test for zinc, iron, manganese,
812 and copper. *Journal of Soil Science Society of America*. 42: 421-428.

813 Ling, W. et al., 2015. Low-molecular-weight organic acids enhance desorption of polycyclic
814 aromatic hydrocarbons from soil. *European Journal of Soil Science*, 66: 339–347.

815 Liste, H.-H. & Alexander, M., 2002. Butanol extraction to predict bioavailability of PAHs in soil.
816 *Chemosphere*, 46: 1011–1017.

817 Liu, G., Wang, J., Zhang, E., Hou, J., Liu, X., 2016. Heavy metal speciation and risk assessment in
818 dry land and paddy soils near mining areas at Southern China. *Environmental Science and*
819 *Pollution Research*, 23: 8709–8720.

820 Liu, H., Cai, X. & Chen, J., 2013. Mathematical model for cyclodextrin alteration of bioavailability
821 of organic pollutants. *Environmental science & technology*, 47: 5835–5842.

822 Liu, L. & Haderlein, S., 2013. A review on the aging phenomena of organic components and their
823 mass transfer through the NAPL interfacial phase. *Chinese Journal of Geochemistry*, 32: 252–
824 260.

825 Loibner, A., 2006. *Sorption and ageing of soil contamination*, Chapter 2. available at
826 <https://pdfs.semanticscholar.org/8399/52b1b56e930ff38bd21583d45fdffe5eae42.pdf>

827 Loibner, A., Holzer, M., Gartner, M., Szolar, O.H.J., & Braun, R., 2000. The Use of Sequential
828 Supercritical Fluid Extraction for Bioavailability Investigations of PAHs in Soil. *Bodenkultur*,
829 51: 225–233.

830 Lomaglio, T., Hattab-Hambli, N., Miard, F., Lebrun, M., Nadillon, R., Trupiano, D., Scippa., G.S.,
831 Gauthier, A., Motelica-Heino, M., Bourgerie, S., & Morabito, D., 2017. Cd, Pb, and Zn
832 mobility and (bio)availability in contaminated soils from a former smelting site amended with
833 biochar. *Environmental Science and Pollution Research*, 1–13.

834 Lorenz, S.E., Hamon, P.E., Hom, H.C., Domingues, E.M., Sequeira, T.H., Christensen, T.H.,
835 McGrath, S.P., 1997. Cadmium and zinc in plants and soil solutions from contaminated soils.
836 *Plant and Soil*, 189: 21–31.

837 Lu, Z.-Y., Yang, X., Wnag, F., Jiang, X., 2011. The tenax extraction technique to predict the
838 bioavailability of aged PAHs in soil. *China Environmental Science*, 31: 647-656.

839 Lukić, B., Huguenot, D., Panico, A., Fabbicino, M., van Hullebusch, E.D., & Esposito, G., 2016.
840 Importance of organic amendment characteristics on bioremediation of PAH-contaminated

841 soil. *Environmental Science and Pollution Research*, 23: 15041–15052.

842 Lydy, M.J., Harwood, A.D., Nutile, S.A., Landrum, P.F., 2015. Tenax extraction of sediments to
843 estimate desorption and bioavailability of hydrophobic contaminants: A literature review.
844 *Integrated Environmental Assessment and Management*, 11: 208–220.

845 Lynch, S., Batty, L. & Byrne, P., 2014. Environmental Risk of Metal Mining Contaminated River
846 Bank Sediment at Redox-Transitional Zones. *Minerals*, 4: 52–73.

847 Makino, T. et al., 2006. Remediation of cadmium contamination in paddy soils by washing with
848 chemicals: Selection of washing chemicals. *Environmental Pollution*, 144: 2–10.

849 Marquès, M. et al., 2016. Photodegradation of polycyclic aromatic hydrocarbons in soils under a
850 climate change base scenario. *Chemosphere*, 148: 495–503.

851 Matong, J.M., Nyaba, L. & Nomngongo, P.N., 2016. Fractionation of trace elements in agricultural
852 soils using ultrasound assisted sequential extraction prior to inductively coupled plasma mass
853 spectrometric determination. *Chemosphere*, 154: 249–257.

854 Mayer, P. et al., 2014. Passive sampling methods for contaminated sediments: scientific rationale
855 supporting use of freely dissolved concentrations. *Integrated environmental assessment and
856 management*, 10: 197–209.

857 McLachlan, M.S. et al., 2011. Bioaccumulation of organic contaminants in humans: A multimedia
858 perspective and the importance of biotransformation. *Environmental Science and Technology*,
859 45: 197–202.

860 McLean, J.E. & Bledsoe, B.E., 1992. Behavior of Metals in Soils. *Office of Research and
861 Development*, EPA/540/S-, pp.1–25.

862 Meers, E. et al., 2007. Comparison of cadmium extractability from soils by commonly used single
863 extraction protocols. *Geoderma*, 141: 247–259.

864 Menegário, A.A. et al., 2017. Use of diffusive gradient in thin films for in situ measurements: A
865 review on the progress in chemical fractionation, speciation and bioavailability of metals in
866 waters. *Analytica Chimica Acta*, 983: 54–66.

867 Meyer, J.S., Farley, K.J. & Garman, E.R., 2015. Metal Mixtures Modeling Evaluation project: 1.
868 Background. *Environmental Toxicology and Chemistry*, 34: 726–740.

869 Miller, D.J. & Hawthorne, S.B., 1998. Method for Determining the Solubilities of Hydrophobic
870 Organics in Subcritical Water. *Analytical Chemistry*, 70: 1618–1621.

871 Miller, W.P., Martens, D.C. & Zelazny, L.W., 1986. Effect of Sequence in Extraction of Trace
872 Metals from Soils. *Soil Science Society of America Journal*, 50: 598–601.

873 Mills, G.A. et al., 2014. Measurement of environmental pollutants using passive sampling devices –
874 an updated commentary on the current state of the art. *Environ. Sci.: Processes Impacts*, 16:
875 369–373.

876 Moreno, A.M., Pérez, L. & Parra, J.G., 2005. Chemical extractability of copper added to soils at
877 two different concentrations in a batch experiment. *Chemical Speciation and Bioavailability*,
878 17: 11–17.

879 Morrison, D.E., Robertson, B.K. & Alexander, M., 2000. Bioavailability to earthworms of aged
880 DDT, DDE, DDD, and dieldrin in soil. *Environmental Science and Technology*, 34: 709–713.

881 Moyo, F. et al., 2014. Sorption of Hydrophobic Organic Compounds on Natural Sorbents and
882 Organoclays from Aqueous and Non-Aqueous Solutions: A Mini-Review. *International*
883 *Journal of Environmental Research and Public Health*, 11: 5020–5048.

884 Muijs, B. & Jonker, M.T.O., 2011. Does equilibrium passive sampling reflect actual in situ
885 bioaccumulation of PAHs and petroleum hydrocarbon mixtures in aquatic worms?
886 *Environmental science & technology*, 46: 937–944.

887 Naidu, R., 2011. *Chemical Bioavailability in Terrestrial Environments*, Elsevier Science.

888 Naidu, R., Channey R., McConnell S., Johnston N., Semple K.T., McGrath S., Dries V., Nathanail
889 P., Harmsen J., Pruszinski A., MacMillan J., Palanisami T. 2015. Towards bioavailability-
890 based soil criteria: past, present and future perspectives, *Environmental Science Pollution*
891 *Research*, 22: 8779-8785.

892 National Research Council, 2003. *Bioavailability of Contaminants in Soils and Sediments:*
893 *Processes, Tools, and Applications*, Washington, DC: The National Academies Press.
894 Available at: <https://www.nap.edu/>

895 National Research Council, 2014. *Physicochemical Properties and Environmental Fate*, Available
896 at: http://www.nap.edu/catalog.php?record_id=18872.

897 Norvell, W.A. & Lindsay, W.L., 1972. Reactions of DTPA Chelates of Iron, Zinc, Copper, and
898 Manganese with Soils1. *Soil Science Society of America Journal*, 36: 778–783.

899 Norvell, W.A. & Lindsay, W.L., 1969. Reactions of EDTA Complexes of Fe, Zn, Mn, and Cu with
900 Soils1. *Soil Science Society of America Journal*, 33: 86–91.

901 Novozamsky, I., Lexmond, T.M. & Houba, V.J.G., 1993. A Single Extraction Procedure of Soil for
902 Evaluation of Uptake of Some Heavy Metals by Plants. *International Journal of*
903 *Environmental Analytical Chemistry*, 51: 47–58.

904 Olaniran, A.O., Balgobind, A. & Pillay, B., 2013. Bioavailability of heavy metals in soil: Impact on
905 microbial biodegradation of organic compounds and possible improvement strategies.
906 *International Journal of Molecular Sciences*, 14: 10197–10228.

907 Olu-Owolabi, B.I., Diagboya, P.N. & Adebowale, K.O., 2014. Evaluation of pyrene sorption–
908 desorption on tropical soils. *Journal of Environmental Management*, 137: 1–9.

909 Ortega-Calvo, J.-J. et al., 2015. From Bioavailability Science to Regulation of Organic Chemicals.
910 *Environmental Science & Technology*, 49: 10255–10264.

- 911 Palleiro, L. et al., 2016. Metal fractionation in topsoils and bed sediments in the Mero River rural
912 basin: Bioavailability and relationship with soil and sediment properties. *Catena*, 144: 34–44.
- 913 Palumbo-Roe, B. et al., 2013. Effect of weathering product assemblages on Pb bioaccessibility in
914 mine waste: Implications for risk management. *Environmental Science and Pollution*
915 *Research*, 20: 7699–7710.
- 916 Pampanin, D.M., 2017. Introduction to Petrogenic Polycyclic Aromatic Hydrocarbons (PAHs) in
917 the Aquatic Environment. *Petrogenic Polycyclic Aromatic Hydrocarbons in the Aquatic*
918 *Environment: Analysis, Synthesis, Toxicity and Environmental Impact*, 1, 3.
- 919 Pampura, T., Groenenberg, J.E. & Rietra, R.P.J.J., 2006. Comparison of methods for copper free
920 ion activity determination in soil solutions of contaminated and background soils. *Forest Snow*
921 *and Landscape Research*, 80: 305–322.
- 922 Pan, Y. et al., 2016. Solubility of trace metals in two contaminated paddy soils exposed to
923 alternating flooding and drainage. *Geoderma*, 261: 59–69.
- 924 Papadopoulos, A., Paton G.I., Reid, B.J., Semple, K.T., 2007. Prediction of PAH biodegradation in
925 field contaminated soils using a cyclodextrin extraction technique. *Journal of Environmental*
926 *Monitoring*, 516–522.
- 927 Parker, R. et al., 2016. The application of Diffusive Gradients in Thin Films (DGT) for improved
928 understanding of metal behaviour at marine disposal sites. *Sci Total Environ*, 575: 1074–1086.
- 929 Parkerton, T. et al., 2012. Guidance on Passive Sampling Methods to Improve Management of
930 Contaminated Sediments: Summary of a SETAC Technical Workshop Edited by Upal Ghosh. ,
931 (November 2012), pp.1–20.
- 932 Pauget, B. et al., 2011. Use of chemical methods to assess Cd and Pb bioavailability to the snail
933 *Cantareus aspersus*: A first attempt taking into account soil characteristics. *Journal of*
934 *Hazardous Materials*, 192: 1804–1811.
- 935 Peijnenburg, W.J.G.M. et al., 2014. Passive Sampling Methods for Contaminated Sediments : State
936 of the Science for Metals. *Integrated Environmental Assessment and Management*, 10: 179–
937 196.
- 938 Peijnenburg, W.J.G.M., Zablotzkaja, M. & Vijver, M.G., 2007. Monitoring metals in terrestrial
939 environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicology*
940 *and Environmental Safety*, 67:163–179.
- 941 Perez, M. et al., 2015. Development of a new passive sampler based on diffusive milligel beads for
942 copper analysis in water. *Analytica Chimica Acta*, 890: 117–123.
- 943 Perez, M. et al., 2016. Diffusive Milli-Gels (DMG) for in situ assessment of metal bioavailability: A
944 comparison with labile metal measurement using Chelex columns and acute toxicity to
945 *Ceriodaphnia dubia* for copper in freshwaters. *Chemosphere*, 164: 7–13.

946 Peters, R.E. et al., 2016. Is received dose from ingested soil independent of soil PAH
947 concentrations?—Animal model results. *Environmental Toxicology and Chemistry*, 35: 2261–
948 2269.

949 Pham, A.L.-T. et al., 2015. Influence of Sulfide Nanoparticles on Dissolved Mercury and Zinc
950 Quantification by Diffusive Gradient in Thin-Film Passive Samplers. *Environmental Science
951 & Technology*, 49: 12897–12903.

952 Pinto, E., Almeida, A.A. & Ferreira, I.M.P.L.V.O., 2015. Assessment of metal(loid)s
953 phytoavailability in intensive agricultural soils by the application of single extractions to
954 rhizosphere soil. *Ecotoxicology and Environmental Safety*, 113: 418–424.

955 Qasim, B. et al., 2015. Potentially toxic element phytoavailability assessment in Technosols from
956 former smelting and mining areas. *Environmental Science and Pollution Research*, 22: 5961–
957 5974.

958 Qian, P. & Schoenau, J.J., 2002. Practical applications of ion exchange resins in agricultural and
959 environmental soil research. *Canadian Journal of Soil Science*, 82: 9–21.

960 Ramakrishnan, B. et al., 2011. Mixtures of environmental pollutants: effects on microorganisms and
961 their activities in soils. In D. M. Whitacre, ed. *Reviews of Environmental Contamination and
962 Toxicology Volume 211*. Reviews of Environmental Contamination and Toxicology. New
963 York, NY: Springer New York, pp. 63–100.

964 Rauret, G., López-Sánchez, J.F., Sahuquillo, A., Muntau, H., et al., 2000. *EUR 19502 - Addendum
965 to EUR-Report 17554 EN: Indicative values for extractable contents (mass fractions) of Cd,
966 Cr, Cu, Ni, Pb and Zn in sediment (CRM 601) following the modified B C R-sequential
967 extraction (three-step) procedure*, Available at: <http://bookshop.europa.eu>.

968 Reid, B.J. et al., 2000a. Nonexhaustive cyclodextrin-based extraction technique for the evaluation
969 of PAH bioavailability. *Environmental Science and Technology*, 34: 3174–3179.

970 Reid, B.J. et al., 2000b. Nonexhaustive Cyclodextrin-Based Extraction Technique for the
971 Evaluation of PAH Bioavailability. *Environmental Science & Technology*, 34: 3174–3179.

972 Reid, B.J., Jones, K.C. & Semple, K.T., 2000. Bioavailability of persistent organic pollutants in
973 soils and sediments—a perspective on mechanisms, consequences and assessment.
974 *Environmental Pollution*, 108: 103–112.

975 Reis, A.P. et al., 2014. Geochemistry, mineralogy, solid-phase fractionation and oral
976 bioaccessibility of lead in urban soils of Lisbon. *Environmental Geochemistry and Health*, 36:
977 867–881.

978 Ren, J. et al., 2015. Sediment metal bioavailability in Lake Taihu, China: evaluation of sequential
979 extraction, DGT, and PBET techniques. *Environmental Science and Pollution Research*, 22:
980 12919–12928.

- 981 Ren, X. et al., 2018. Sorption, transport and biodegradation – An insight into bioavailability of
982 persistent organic pollutants in soil. *Science of The Total Environment*, 610–611: 1154–1163.
- 983 Renoux, A.Y. et al., 2013. Risk-based management of site soils contaminated with a mixture of
984 hazardous substances: Methodological approach and case study. *Human and Ecological Risk
985 Assessment: An International Journal*, 19: 1127–1146.
- 986 Rhodes, A.H., Carlin, A. & Semple, K.T., 2008. Impact of black carbon in the extraction and
987 mineralization of phenanthrene in soil. *Environmental science & technology*, 42: 740–745.
- 988 Riding, M.J. et al., 2013. Chemical measures of bioavailability/bioaccessibility of PAHs in soil:
989 fundamentals to application. *Journal of hazardous materials*, 261: 687–700.
- 990 Rinklebe, J., Knox, A.S. & Paller, M., 2016. *Trace Elements in Waterlogged Soils and Sediments*,
991 CRC Press, 386 pages.
- 992 Rodgers, K.J., Hursthouse, A. & Cuthbert, S., 2015. The potential of sequential extraction in the
993 characterisation and management of wastes from steel processing: A prospective review.
994 *International Journal of Environmental Research and Public Health*, 12: 11724–11755.
- 995 Romero-Freire, A. et al., 2017. Effects of aging and soil properties on zinc oxide nanoparticle
996 availability and its ecotoxicological effects to the earthworm *Eisenia andrei*. *Environmental
997 Toxicology and Chemistry*, 36: 137–146.
- 998 Rosado, D., Usero, J. & Morillo, J., 2016. Ability of 3 extraction methods (BCR, Tessier and
999 protease K) to estimate bioavailable metals in sediments from Huelva estuary (Southwestern
1000 Spain). *Marine Pollution Bulletin*, 102: 65–71.
- 1001 Rostami, I. & Juhasz, A.L., 2011. Assessment of Persistent Organic Pollutant (POP) Bioavailability
1002 and Bioaccessibility for Human Health Exposure Assessment: A Critical Review. *Critical
1003 Reviews in Environmental Science and Technology*, 41: 623–656.
- 1004 Ruby, M. V. et al., 1996. Estimation of lead and arsenic bioavailability using a physiologically
1005 based extraction test. *Environmental Science and Technology*, 30: 422–430.
- 1006 Rüdél, H. et al., 2015. Consideration of the bioavailability of metal/metalloid species in freshwaters:
1007 experiences regarding the implementation of biotic ligand model-based approaches in risk
1008 assessment frameworks. *Environmental Science and Pollution Research*, 22: 7405–7421.
- 1009 Rurikova, D. & Kudrava, M., 2006. Anodic stripping voltammetric determination of lead and
1010 cadmium in soil extracts. *Chemical Papers-Chemicke Zvesti*, 60: 22–26.
- 1011 Sabljic, A. & Nakagawa, Y., 2014. Sorption and Quantitative Structure-Activity Relationship
1012 (QSAR). In *Non-First Order Degradation and Time-Dependent Sorption of Organic
1013 Chemicals in Soil*. ACS Symposium Series. American Chemical Society, pp. 5–85.
- 1014 Sam K., Coulon F., Prpich G. 2016. Working towards an integrated land contamination management
1015 framework for Nigeria. *Science of the Total Environment*. 571: 916-925

- 1016 Sahito, O.M. et al., 2015. Evaluation of heavy metal bioavailability in soil amended with poultry
1017 manure using single and BCR sequential extractions. *International Journal of Environmental*
1018 *Analytical Chemistry*, 95: 1066–1079.
- 1019 Sakan, S. et al., 2016. Comparison of single extraction procedures and the application of an index
1020 for the assessment of heavy metal bioavailability in river sediments. *Environmental Science*
1021 *and Pollution Research*, 23: 21485–21500.
- 1022 Sauvé, S., McBride, M.B. & Hendershot, W.H., 1997. Speciation of Lead in Contaminated Soils.
1023 *Environmental Pollution*, 98: 149–155.
- 1024 Sayadi, M.H., Rezaei, A. & Sayyed, M.R.G., 2017. Grain size fraction of heavy metals in soil and
1025 their relationship with land use. *Proceedings of the International Academy of Ecology and*
1026 *Environmentl Sciences* 7: 1–11.
- 1027 Schneider, A.R. et al., 2016. Influence of dissolved organic matter and manganese oxides on metal
1028 speciation in soil solution: A modelling approach. *Environmental Pollution*, 213: 618–627.
- 1029 Selim, H.M., 2015. *Phosphate in Soils: Interaction with Micronutrients, Radionuclides and Heavy*
1030 *Metals*, CRC Press, 381 pages.
- 1031 Semple, K.T. et al., 2004. Peer reviewed: defining bioavailability and bioaccessibility of
1032 contaminated soil and sediment is complicated. *Environmental Science and Technology*, 38:
1033 228A-231A.
- 1034 Seshadri, B., Bolan, N.. & Naidu, R., 2015. Rhizosphere-induced heavy metal(loid) transformation
1035 in relation to bioavailability and remediation. *Journal of soil science and plant nutrition*, 15:
1036 524-548
- 1037 Settimo, L. et al., 2014. Fate and lability of silver in soils: Effect of ageing. *Environmental*
1038 *Pollution*, 191: 151–157.
- 1039 Shahid, M. et al., 2017. Chromium speciation, bioavailability, uptake, toxicity and detoxification in
1040 soil-plant system: A review. *Chemosphere*, 178: 513–533.
- 1041 Sharma, P. et al., 2010. Influence of Natural Organic Matter on As Transport and Retention.
1042 *Environmental Science & Technology*, 45: 546–553.
- 1043 Shor, L.M. et al., 2003. Intra-aggregate mass transport-limited bioavailability of polycyclic
1044 aromatic hydrocarbons to Mycobacterium strain PC01. *Environmental science & technology*,
1045 37: 1545–1552.
- 1046 Sihag, S., Pathak, H. & Jaroli, D.P., 2014. Factors Affecting the Rate of Biodegradation of
1047 Polyaromatic Hydrocarbons. *International Journal of Pure and Applied Bioscience*, 2: 185–
1048 202.
- 1049 Smith, M.J. et al., 2011. Study of PAH dissipation and phytoremediation in soils: Comparing
1050 freshly spiked with weathered soil from a former coking works. *Journal of Hazardous*

1051 *Materials*, 192: 1219–1225.

1052 Smith, R.M., 2002. Extractions with superheated water. *Journal of Chromatography A*, 975: 31–46.

1053 Sormunen, A.J. et al., 2008. Desorption of sediment-associated polychlorinated dibenzo- p -dioxins
1054 , dibenzofurans , diphenyl ethers and hydroxydiphenyl ethers from contaminated sediment.
1055 *Chemosphere* , 72: 1–7.

1056 Stegemeier, J.P. et al., 2015. Speciation Matters: Bioavailability of Silver and Silver Sulfide
1057 Nanoparticles to Alfalfa (*Medicago sativa*). *Environmental science & technology*, 49: 8451–
1058 60.

1059 Stokes, J.D. et al., 2005. Prediction of Polycyclic Aromatic Hydrocarbon Biodegradation in
1060 Contaminated Soils Using an Aqueous Hydroxypropyl- β -Cyclodextrin Extraction Technique.
1061 *Environmental Toxicology and Chemistry*, 24: 1325-1330.

1062 Stroud, J.L., Paton, G.I. & Semple, K.T., 2009. Chemosphere Predicting the biodegradation of
1063 target hydrocarbons in the presence of mixed contaminants in soil. *Chemosphere*, 74: 563–567.

1064 Stroud, J.L., Paton, G.I. & Semple, K.T., 2008. Linking chemical extraction to microbial
1065 degradation of 14 C-hexadecane in soil. *Environmental Pollution*, 156: 474–481.

1066 Stroud, J.L., Paton, G.I. & Semple, K.T., 2007. Microbe-aliphatic hydrocarbon interactions in soil:
1067 implications for biodegradation and bioremediation. *Journal of Applied Microbiology*, 102:
1068 1239–1253.

1069 Sun, F. & Zhou, Q., 2010. Interactive effects of 1,4-dichlorobenzene and heavy metals on their
1070 sorption behaviors in two Chinese soils. *Archives of Environmental Contamination and*
1071 *Toxicology*, 58: 33–41.

1072 Sun, H. & Li, J., 2005. Availability of pyrene in unaged and aged soils to earthworm uptake,
1073 butanol extraction and SFE. *Water, Air, and Soil Pollution* 166, 353–365.

1074 Sun, J. et al., 2015. Speciation and determination of bioavailable arsenic species in soil samples by
1075 one-step solvent extraction and high-performance liquid chromatography with inductively
1076 coupled plasma mass spectrometry. *Journal of Separation Science*, 38: 943–950.

1077 Sunda, W.G., 1984. Measurement of manganese, zinc and cadmium complexation in seawater using
1078 Chelex ion exchange equilibria. *Marine Chemistry*, 14: 365–378.

1079 Sungur, A. et al., 2015. Characterization of Heavy Metal Fractions in Agricultural Soils by
1080 Sequential Extraction Procedure: The Relationship Between Soil Properties and Heavy Metal
1081 Fractions. *Soil and Sediment Contamination: An International Journal*, 24: 1–15.

1082 Sungur, A., Soylak, M. & Ozcan, H., 2014. Investigation of heavy metal mobility and availability
1083 by the BCR sequential extraction procedure: Relationship between soil properties and heavy
1084 metals availability. *Chemical Speciation and Bioavailability*, 26: 219–230.

1085 Tahervand, S. & Jalali, M., 2016. Sorption, desorption, and speciation of Cd, Ni, and Fe by four

1086 calcareous soils as affected by pH. *Environmental Monitoring and Assessment*, 188: 322-336.

1087 Tang, J., Liste, H.H. & Alexander, M., 2002. Chemical assays of availability to earthworms of
1088 polycyclic aromatic hydrocarbons in soil. *Chemosphere*, 48: 35–42.

1089 Tang, J., Robertson, B.K. & Alexander, M., 1999. Chemical-Extraction Methods To Estimate
1090 Bioavailability of DDT, DDE, and DDD in Soil. *Environmental Science & Technology*, 33:
1091 4346–4351.

1092 Tangahu, B.V. et al., 2011. A review on heavy metals (As, Pb, and Hg) uptake by plants through
1093 phytoremediation. *International Journal of Chemical Engineering*, Article ID 939161, 31
1094 pages.

1095 Tao, Y., Xue, B. & Yao, S., 2015. Using linoleic acid embedded cellulose acetate membranes to in
1096 situ monitor polycyclic aromatic hydrocarbons in lakes and predict their bioavailability to
1097 submerged macrophytes. *Environmental science & technology*, 49: 6077–6084.

1098 Tchounwou, P.B. et al., 2012. Heavy Metals Toxicity and the Environment. In: Luch A., (eds)
1099 *Molecular, Clinical and Environmental Toxicology, Experientia Supplementum*, 101:133–164.

1100 Tessier, A., Campbell, P.G.C. & Bisson, M., 1979. Sequential Extraction Procedure for the
1101 Speciation of Particulate Trace Metals. *Analytical Chemistry*, 51: 844–851.

1102 Thompson, T., 2016. Bioavailability of Contaminants in Soils and Sediments : Status and
1103 Recommendations. Washington, DC: The National Academies Press.
1104 <https://doi.org/10.17226/10523>.

1105 Tipping, E. et al., 2003. The solid-solution partitioning of heavy metals (Cu, Zn, Cd, Pb) in upland
1106 soils of England and Wales. *Environmental Pollution*, 125: 213–225.

1107 Trelu, C. et al., 2017. Characteristics of PAH tar oil contaminated soils—Black particles, resins and
1108 implications for treatment strategies. *Journal of Hazardous Materials*, 327: 206–215.

1109 Tri-Service Ecological Risk Assessment Workgroup, 2003. *Guide for Incorporating Bioavailability*
1110 *Adjustments into Human Health and Ecological Risk Assessments at US Department of*
1111 *Defense Facilities Part 1 : Overview of Metals Bioavailability June 2003* available at
1112 <http://www.itrcweb.org/>

1113 Trimble, T.A., You, J. & Lydy, M.J., 2008. Bioavailability of PCBs from field-collected sediments :
1114 Application of Tenax extraction and matrix-SPME techniques. *Chemosphere*, 71: 337–344.

1115 Ure, A.M., 1996. Single extraction schemes for soil analysis and related applications. *Science of the*
1116 *Total Environment*, 178: 3–10.

1117 Ure, A.M. et al., 1993. Speciation of Heavy Metals in Soils and Sediments. An Account of the
1118 Improvement and Harmonization of Extraction Techniques Undertaken Under the Auspices of
1119 the BCR of the Commission of the European Communities. *International Journal of*
1120 *Environmental Analytical Chemistry*, 51: 135–151.

- 1121 Van Genderen, E., Adams, W., Dwyer, R., Garman, E., & Gorsuch, J. 2015. Modeling and
1122 interpreting biological effects of mixtures in the environment: Introduction to the metal
1123 mixture modeling evaluation project. *Environmental Toxicology and Chemistry*, 34: 721–725.
- 1124 Vasseur, P. & Bonnard, M., 2014. Ecogenotoxicology in earthworms: A review. *Current Zoology*,
1125 60: 255–272.
- 1126 Vázquez Vázquez, F.A., Pérez Cid, B. & Río Segade, S., 2016. Assessment of metal bioavailability
1127 in the vineyard soil-grapevine system using different extraction methods. *Food Chemistry*,
1128 208: 199–208.
- 1129 Vegter, J., Lowe, J. & Kasamas, H., 2002. *Sustainable management of contaminated land: an*
1130 *overview*, Austria: CLARINET. Available at www.commonforum.eu (accessed on 29/09/2017)
- 1131 Vicent, T. et al., 2013. *Emerging Organic Contaminants in Sludges: Analysis, Fate and Biological*
1132 *Treatment*, Springer Berlin Heidelberg.
- 1133 Vila, J., Tauler, M. & Grifoll, M., 2015. Bacterial PAH degradation in marine and terrestrial
1134 habitats. *Current Opinion in Biotechnology*, 33: 95–102.
- 1135 Villaverde, J. & Pe, I., 2012. Enhanced Solubilisation of Six PAHs by Three Synthetic
1136 Cyclodextrins for Remediation Applications : Molecular Modelling of the Inclusion
1137 Complexes. *PLOS one*, 7: 1–8.
- 1138 Violante, A. et al., 2010. Mobility and Bioavailability of Heavy Metals and Metalloids in Soil
1139 Environments. *Journal of soil science and plant nutrition*, 10: 268–292.
- 1140 Van Der Wal, L. et al., 2004. Solid-Phase Microextraction To Predict Bioavailability and
1141 Accumulation of Organic Micropollutants in Terrestrial Organisms after Exposure to a Field-
1142 Contaminated Soil. *Environmental Science and Technology*, 38: 4842–4848.
- 1143 van der Wal, L., van Gestel, C.A. & Hermens, J.L., 2004. Solid phase microextraction as a tool to
1144 predict internal concentrations of soil contaminants in terrestrial organisms after exposure to a
1145 laboratory standard soil. *Chemosphere*, 54: 561–568.
- 1146 Wang, Y. et al., 2017. Dynamic arsenic aging processes and their mechanisms in nine types of
1147 Chinese soils. *Chemosphere*, 187: 404–412.
- 1148 Wang, Y. et al., 2015. Effect of aging on the bioavailability and fractionation of arsenic in soils
1149 derived from five parent materials in a red soil region of Southern China. *Environmental*
1150 *Pollution*, 207: 79–87.
- 1151 Wang, Y. et al., 2016. Impact of Humic on Soil Adsorption and Remediation of Cd(II), Pb(II), and
1152 Cu(II). *Soil and Sediment Contamination*, 25: 700–715.
- 1153 Weber, W.J. et al., 2002. Annual Report for Michigan Great Lakes Protection Fund Grant. , pp.1–
1154 22.
- 1155 Weber, W.J., 2001. Rapid Prediction of Long-Term Rates of Contaminant Desorption from Soils

1156 and Sediments. *Environmental Science and Technology*, 35: 427–433.

1157 Weber, W.J., McGinley, P.M. & Katz, L.E., 1992. A distributed reactivity model for sorption by
1158 soils and sediments. 1. Conceptual basis and equilibrium assessments. *Environmental Science*
1159 *& Technology*, 26: 1955–1962.

1160 Weng, L., Van Riemsdijk, W.H. & Temminghoff, E.J.M., 2005. Kinetic aspects of Donnan
1161 membrane technique for measuring free trace cation concentration. *Analytical Chemistry*, 77:
1162 2852–2861.

1163 Weng, N. et al., 2015. Insight into unresolved complex mixtures of aromatic hydrocarbons in heavy
1164 oil via two-dimensional gas chromatography coupled with time-of-flight mass spectrometry
1165 analysis. *Journal of Chromatography A*, 1398: 94–107.

1166 Wijayawardena, M. a. A. et al., 2015. Influence of ageing on lead bioavailability in soils: a swine
1167 study. *Environmental Science and Pollution Research*, 22: 8979–8988.

1168 Workman, S.M. & Lindsay, W.L., 1990. Estimating Divalent Cadmium Activities Measured in
1169 Arid-Zone Soils using Competitive Chelation. *Soil Science Society of America Journal*, 54:
1170 987–993.

1171 Wu, G. et al., 2013. Influence of mature compost amendment on total and bioavailable polycyclic
1172 aromatic hydrocarbons in contaminated soils. *Chemosphere*, 90: 2240–2246.

1173 Wuana, R. a, Okieimen, F.E. & Vesuwe, R.N., 2014. Mixed contaminant interactions in soil :
1174 Implications for bioavailability , risk assessment and remediation. *African Journal of*
1175 *Environmental Science and Technology*, 8: 691–706.

1176 Wyszowska, J. et al., 2012. Effect of cadmium, copper and zinc on plants, soil microorganisms
1177 and soil enzymes. *Journal of Elementology*, 18: 769-793.

1178 Xiao, R. et al., 2014. Polycyclic aromatic hydrocarbons (PAHs) in wetland soils under different
1179 land uses in a coastal estuary: Toxic levels, sources and relationships with soil organic matter
1180 and water-stable aggregates. *Chemosphere*, 110: 8–16.

1181 Xing, B., Senesi, N. & Huang, P.M., 2011. *Biophysico-Chemical Processes of Anthropogenic*
1182 *Organic Compounds in Environmental Systems*, Wiley, 592 pages.

1183 Xue, H., Kistler, D. & Sigg, L., 1995. Competition of copper and zinc for strong ligands in a
1184 eutrophic lake. *Limnology and Oceanography*, 40: 1142–1152.

1185 Yang, X. et al., 2016. Bioavailability of Polycyclic Aromatic Hydrocarbons and their Potential
1186 Application in Eco-risk Assessment and Source Apportionment in Urban River Sediment.
1187 *Scientific Reports*, 6: 23134. Available at: <http://www.nature.com/articles/srep23134> (accessed
1188 on 29 September 2017).

1189 Yin, Y. et al., 1997. Adsorption/Desorption Isotherms of Hg(II) by Soil. *Soil Science*, 162: 35-45.

1190 You, J. & Landrum, P.F., 2006. Comparison of Chemical Approaches for Assessing Bioavailability

1191 of Sediment-Associated Contaminants. *Environmental Science and Technology*, 40: 6348–
1192 6353.

1193 You, J. & Pehkonen, S., 2007. Desorption of Hydrophobic Compounds from Laboratory-Spiked
1194 Sediments Measured by Tenax Absorbent and Matrix Solid-Phase Microextraction.
1195 *Environmental Science and Technology*, 41(16), pp.5672–5678.

1196 Young, S.D., 2013. Chemistry of Heavy Metals and Metalloids in Soils BT - Heavy Metals in
1197 Soils: Trace Metals and Metalloids in Soils and their Bioavailability. In B. J. Alloway, ed.
1198 Dordrecht: Springer Netherlands, pp. 51–95.

1199 Young, S.D. et al., 2000. Methods for determining labile cadmium and zinc in soil. *European*
1200 *Journal of Soil Science*, 51: 129–136.

1201 Yu, H.-Y. et al., 2016. Iron Redox Cycling Coupled to Transformation and Immobilization of
1202 Heavy Metals: Implications for Paddy Rice Safety in the Red Soil of South China. Chapter 5
1203 In D. L. B. T.-A. in A. Sparks, ed. *Advances in Agronomy*. Academic Press, pp. 279–317.

1204 Yu, H.Y. et al., 2016. Cadmium availability in rice paddy fields from a mining area: The effects of
1205 soil properties highlighting iron fractions and pH value. *Environmental Pollution*, 209: 38–45.

1206 Yu, L. et al., 2018. Abiotic factors controlling bioavailability and bioaccessibility of polycyclic
1207 aromatic hydrocarbons in soil : Putting together a bigger picture. *Science of the Total*
1208 *Environment*, 614: 1140–1153.

1209 Zhang, D. et al., 2014. Synergistic role of different soil components in slow sorption kinetics of
1210 polar organic contaminants. *Environmental Pollution*, 184: 123–130.

1211 Zhang, H. et al., 1998. In situ measurements of solution concentrations and fluxes of trace metals in
1212 sells using DGT. *Environmental Science and Technology*, 32: 704–710.

1213 Zhang, J. & Fan, S., 2016. Influence of PAH speciation in soils on vegetative uptake of PAHs using
1214 successive extraction. *Journal of Hazardous Materials*, 320: 114–122.

1215 Zhang, M. et al., 2014. Sorption of polycyclic aromatic hydrocarbons (PAHs) to lignin: Effects of
1216 hydrophobicity and temperature. *Bulletin of Environmental Contamination and Toxicology*,
1217 93: 84–88.

1218 Zhang, W. et al., 2015. The roles of humic substances in the interactions of phenanthrene and heavy
1219 metals on the bentonite surface. *Journal of Soils and Sediments*, 15: 1463–1472.

1220 Zhao, C.M., Campbell, P.G.C. & Wilkinson, K.J., 2016. When are metal complexes bioavailable?
1221 *Environmental Chemistry*, 13: 425–433.

1222 Zima, J. & van den Berg, C.M.G., 1994. Determination of arsenic in sea water by cathodic stripping
1223 voltammetry in the presence of pyrrolidine dithiocarbamate. *Analytica Chimica Acta*, 289:
1224 291–298.

1225 Zimmerman, A.J. & Weindorf, D.C., 2010. Heavy Metal and Trace Metal Analysis in Soil by

- 1226 Sequential Extraction: A Review of Procedures. *International Journal of Analytical Chemistry*,
1227 Article ID 387803, 7 pages.
- 1228 de Zwart, D. & Posthuma, L., 2005. Complex mixture toxicity for single and multiple species.
1229 *Environmental Toxicology and Chemistry*, 24: 2665-2676.