

Factors governing the solid phase distribution of Cr, Cu and As in contaminated soil after 40 years of ageing

Stacie Tardif^{1*}, Sabrina Cipullo², Helle U. Sørensen³, Joanna Wragg⁴, Peter E. Holm¹, Frederic Coulon²,
Kristian K. Brandt¹, Mark Cave⁴

¹University of Copenhagen, Department of Plant and Environmental Sciences, Frederiksberg, DK-1871, Denmark

²Cranfield University, School of Water, Energy and Environment, Cranfield, MK43 0AL, UK

³Geological Survey of Denmark and Greenland, Copenhagen K, DK-1350, Denmark

⁴British Geological Survey, Keyworth, Nottingham, NG12 5GG, UK

*Corresponding author: stta@plen.ku.dk

Abstract:

The physico-chemical factors affecting the distribution, behavior and speciation of chromium (Cr), copper (Cu) and arsenic (As) was investigated at a former wood impregnation site (Fredensborg, Denmark). Forty soil samples were collected and extracted using a sequential extraction technique known as the Chemometric Identification of Substrates and Element Distributions (CISED) and multivariate statistical tools (redundancy analysis) were applied. CISED data was linked to water-extractable Cr, Cu and As and bioavailable Cu as determined by a whole-cell bacterial bioreporter assay. Results showed that soil pH significantly affected the solid phase distribution of all three elements on site. Additionally, elements competing for binding sites, Ca, Mg and Mn in the case of Cu, and P, in the case of As, played a major role in the distribution of these elements in soil. Element-specific distributions were observed amongst the six identified soil phases including residual pore salts, exchangeable, carbonates (tentative designation), Mn-Al oxide, amorphous Fe oxide, and crystalline Fe oxide. While Cr was strongly bound to non-extractable crystalline Fe oxide in the oxic top soil, Cu and notably, As were associated with readily extractable phases. While water-extractable Cu and As significantly correlated to CISED identified soil phases, water-extractable Cr and bioavailable Cu did not, the latter suggesting that sequential extraction schemes may not be ideally suited for inferring bioavailability and toxicity of elements to ecological endpoints such as microbes in soil. Findings from this study suggest that after 40 years of ageing, Cu and As at CCA-contaminated field sites constitute the highest risk to environmental and human health through ecotoxicological impact and leaching to groundwater reservoirs.

1. Introduction

Multi-element contaminated sites resulting from anthropogenic activity are of serious concern as: (1) these types of contaminants accumulate and persist in the environment, (2) complex chemical mixtures of elements behave differently from each other in soil and (3) the interactions between elements and potential synergistic effects are not well understood. Chromated copper arsenate (CCA) contaminated sites are a result of the wood impregnation industry which, for several decades, used a solution of salts (CrO_3 and CuO) and arsenic acid (H_3AsO_4) to impregnate wood to preserve it from decay by biotic and abiotic factors (Humphrey, 2002) [\a00143.science.domain\vxr881\Documents\DATA \(D\)\Abstracts-papers\CISED Stacie\Revision STOTEN Oct 2018\](#) - Humphrey, 2002. As environmental legislations in Europe (Directive, 2003) and North America became more stringent over the years, the use of these impregnation mixtures was severely curtailed and numerous sites were closed down, remaining contaminated to this today (Bhattacharya et al., 2002).

During the fixation of CCA-treated wood a number of different chemical reactions occur and their resulting speciation can have significant effects on leachate toxicity and mobility (Hingston et al., 2001). Namely, reactive and mobile hexavalent Cr is reduced to insoluble Cr(III); however a fraction of Cr(VI) may remain oxidized (Gezer and Cooper, 2016). Arsenic is fixed as chromium arsenates, but As(V) can be reduced to the toxic and mobile As(III) in soil (Nielsen et al., 2010). Chromium, Cu and As, in the form of Cr(VI), Cu(II), and As(III), can cause significant environmental impact, primarily as a result of leaching to groundwater reservoirs and transport to downstream lakes and streams (Nielsen et al., 2013) and due to their high toxicity to soil and water-dwelling organism alike (Gezer and Cooper, 2016; Nunes et al., 2016). Past studies of CCA mixtures have reported that Cu was generally immobilized in the oxic top soil while Cr and As exhibited high spatial mobility and risk of leaching (Allinson et al., 2000; Andersen et al., 1996; Carey et al., 2002; Hopp et al., 2006). However, these studies have been focused on short-term monitoring and/or under controlled laboratory environments and therefore do not provide a useful

basis for determining element speciation and mobility in the field after substantial ageing processes (Allinson et al., 2000; Andersen et al., 1996; Carey et al., 2002; Hopp et al., 2006).

Several sequential extraction techniques are available to quantify the distribution of elements in soil phases, providing valuable tools to make assumptions on element speciation and mobility in the field (Bacon and Davidson, 2008). In general, the extractable phases are categorized as exchangeable, specifically adsorbed, carbonates, Fe and Mn-oxides, organic matter and sulphides, and mineral lattice or residual (Cave and Harmon, 1997; Cave and Wragg, 1997). Among sequential extraction methodologies, the Chemometric Identification of Substrates and Element Distributions (CISED) provides an estimate of how much of the potentially harmful elements can be found in each soil solid phase. This approach was developed as a result of limitations/weaknesses associated with 'classical' sequential extractions (Cave et al., 2004): lack of specificity for the target phase, methodological definition of the distribution of elements and re-adsorption of liberated metals, the varied (and sometimes long) extraction times required to undertake the methodology, and the complexity of the extraction matrices for analysis. The CISED methodology considers that a soil and its associated phases are multi-elemental and has taken the approach of using a non-specific acid extractant at increasing strengths combined with mixture resolution analysis to combat the limitations outlined above (Cave et al., 2004). It has been successfully validated by comparison to other sequential extraction procedures such as the Tessier method (Cave and Wragg, 1997), to backscattered scanning electron microscopy (BSEM) analysis and digital energy-dispersive X-ray microanalysis (Cave et al., 2004) and applied to diverse soil samples (Cipullo et al., 2018; Wragg and Cave, 2012; Wragg et al., 2014). Furthermore, although the link between human oral bioaccessibility and CISED has been studied (Cox et al., 2013; Wragg and Cave, 2012; Wragg et al., 2014), the relationship between elements associated with CISED identified solid phases and bioavailable elements, in the context of biological receptors such as bacteria, remains unexplored. Physico-chemical factors, such as pH (Lu et al., 2005), organic C (Andersen et al., 1996), reactive Fe oxides (Yang et al., 2002) and P content (Bolan et al., 2013; Cao and Ma, 2004), all play an

important role in influencing the adsorption and occlusion of these elements between the soil solid and liquid phases. However, studying the complex relationships between physico-chemical factors and element distribution amongst phases in the field can be very challenging and requires the use of multivariate statistical tools. Redundancy analysis (RDA), a form of constrained ordination founded on the principles of multiple linear regression, can test the relationships between multivariate explanatory and response datasets (Legendre and Anderson, 1999) and allows us to explore such complex relationships in the field.

To our knowledge, no field studies investigating the physico-chemical factors governing the solid-phase distribution of CCA in aged soils have occurred. To this end, we collected soil samples along a contamination gradient at a CCA legacy site and assessed the factors affecting the solid-phase distribution of Cr, Cu and As in the field. The study objectives were: (i) to identify the common soil solid phases on site and the distribution of Cr, Cu and As amongst these phases, (ii) to study the influence of different physico-chemical factors on this solid phase distribution, and finally, (iii) to test the relationship between water extractable Cr, Cu and As and bioavailable Cu, as determined by a whole-cell bacterial bioreporter assay, with elements distributed amongst the CISED easily extractable phase.

2. Materials and Methods

2.1. Field site history and geology

The multi-element field contaminated site, located in Fredensborg, Denmark (55°57N, 12 °21E), was used as a wood impregnation site, coating wood telephone poles with a mixture of CCA for over 20 years (**Fig. 1a**). The most recent impregnation liquid being used on site was a water soluble formulation of salts CrO_3 (Cr(VI)) and CuO (Cu(II)) and arsenic acid As_2O_5 (As(V)) as 34%, 27% and 25% respectively (Nielsen et al., 2014). The main contamination on site was derived from drippings of surplus preservatives into the soil. In 1976, the entire site was abandoned and left as a

brownfield in the surrounding forest. It was replanted in clearly visible rows of birch (*Betula*) and oak (*Quercus*), and today, trees are found in various states of growth because of the heterogeneous soil contamination.

In 1989, a full-scale site investigation was launched and reports on the bedrock geology and soil chemistry by the company Samfundsteknik were published (Samfundsteknik, 1989a; Samfundsteknik, 1989b) and an updated site investigation for the period between 1977 and 2009 was reported by Nielsen et al. (2010). Briefly, the 1989 investigation reported that soil contamination was widespread across the 65,000 m² site as a result of direct spillage and dripping from highly contaminated bark during the drying process. The main contamination on sites derive from the metals Cr (Cr³⁺ and Cr⁶⁺), Cu and metalloid As (As³⁺, As⁵⁺), in the top soil and shallow groundwater. Top soil (0-0.5m) was found to be the most heavily contaminated with average concentrations of 4-700 mg kg⁻¹, 6-1500 mg kg⁻¹ and 2-1000 mg kg⁻¹ for Cr, Cu and As, respectively (**Fig. 1b**: modified from Samfundsteknik report).

The bedrock geology at this site is predominantly a sandy glaciofluvial formation with lenses of sand and clay. The site is located on the edge of the regional formation of the Alnarp Valley, which extends across Northern Zealand and South of Sweden. The contaminated site is found upstream of the Esum Lake and connected via shallow, intermediate and deep aquifers, with local sediments constituting one of the most important groundwater reservoirs in North Zealand, Denmark. Nielsen et al. postulated that the most likely route for CCA leaching to the lake being the intermediary and deep aquifers (Nielsen et al., 2010).

2.2. Experimental design and soil sampling

Ten sampling stations (1-10) were established at this site in October 2016. The sample selection was based on the 1989 site investigation concentrations and a preliminary sampling (data not shown) validating the establishment of a contamination gradient (**Fig. 1b**). At each location a plot of 1 m² was laid out and each plot separated into 4 quadrants with the intention of sampling

biological replicates. It became apparent after soil chemical analysis, however, that heterogeneity on site was extensive and as a result, samples were kept separate for downstream analysis. Samples were named from 1-40 based on total cumulative Cr, Cu and As concentrations in the samples.

Approximately 500 g of top soil (~10 cm in depth) was collected from each quadrant in a randomized sampling design within the quadrant and mixed thoroughly on site in plastic bags in order to obtain a representative sample (10 sampling locations \times 4 quadrants = 40 samples).

Individual fresh soil samples were divided and processed for analysis in the following way

(Supplementary Fig.1): (1) 2 g into 15mL falcon tube for water-extraction/bacterial bioreporter; (2) 5 g into 50mL falcon tube for pH measurement; (3) 10 g in petri-dish for water content measurement; (4) 5 g in airtight anoxic vials and stored at -20 °C for reactive Fe oxide measurement and (5) 10 g was pooled from each quadrant into one composite sample for each sampling location and thoroughly homogenized for texture analysis. Remaining soil was air-dried, sieved (2 mm mesh size) and ground to a fine powder with mechanical pestle and mortar (henceforth referred to as pulverized) for downstream physical-chemical analyses (Supplementary F1).

2.3. Soil physico-chemical analyses

Particle size distribution was determined on composite fresh soil by the hydrometer method (Gee and Bauder, 1986) and classified according to international standard particle size classes. Water content was measured after drying samples for 12h at 105°C. Soil pH was measured on fresh soil in MilliQ water at a 1:2.5 ratio (soil liquid, w/v) using a combined glass electrode (Metrohm, 6.0228.000). Poorly crystalline Fe oxides, henceforth referred to as reactive fraction of Fe oxides, were measured based on the experimental procedures described in Nguyen et al. (2014). Two chemical extractions were performed in parallel for 24 hours in 1:100 (soil liquid, w/v) using: 1) 0.5M formic acid at pH 3, targeting the carbonates and phosphates and 10mM ascorbic acid and 2) 0.5M formic acid at pH 3, to target the ferrihydrite and very poorly crystalline goethite. Poorly crystalline Fe oxides was determined by subtracting Fe^{2+} found in the first extraction from Fe^{2+} in

the second extraction. Total carbon and total nitrogen concentrations were analyzed by dry combustion at 1200° using a Vario Macro cube C/N elemental analyzer on dry, 2mm sieved and pulverized soil samples (Elementar Analysen systeme GmbH, Hanau, Germany). Data quality of C/N analysis was evaluated by inclusion of standard reference materials (1515 Apple leaves and 141d acetanilide, National Institute of Standards and Technology (NIST), Gaithers-burg, MD, USA). Plant-available P was determined by the Olsen P method, extracting the dry, 2mm sieved and pulverized soil with a 0.5M hydrogen carbonate solution at pH 8.5 and then spectrophotometrically measuring the P content in solution, following ISO 11263:1994 protocol.

2.4. Water-extractable, total and CISED extractions

Water-extractable elements (Al, As, Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mo, Mn, Na, Ni, P, Pb, S, Sb, Si, Se, Sr, V and Zn) were analyzed in soil-water extracts at a 1:5 ratio (soil liquid, w/v), extracted on a horizontal shaker at 200rpm for 2 hours.

Total element concentrations (Al, As, Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mo, Mn, Na, Ni, P, Pb, S, Sb, Si, Se, Sr, V and Zn) were determined following ISO 11047 protocol, which extracts the elements with aqua regia (HCl/HNO₃) using a microwave digestion system.

The solid phase distribution of elements (Al, As, Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mo, Mn, Na, Ni, P, Pb, S, Sb, Si, Se, Sr, V and Zn) was investigated using a modified non-specific sequential extraction method, based on the CISED methodology (Cave et al., 2004), as described in (Cipullo et al., 2018). Briefly, 2 g of pulverized soil was sequentially extracted (twice per reagent) with deionized water (DI) and HNO₃ in increasing dissolution strength: DI (E1-E2), 0.01 M (E3-E4), 0.05 M (E5-E6), 0.1 M (E7-E8), 0.5 M (E9-E10), 1.0 M (E11-E12) and 5.0 M (E13-E14), with H₂O₂ (100 Volumes) addition in the last 4 extracts.

2.5. Bacterial bioreporter assay

Henceforth, Bioavailable Cu (Cu_{bio}) is operationally defined as Cu species in soil-water extracts that are able to induce gene expression in a Cu-specific *Pseudomonas fluorescens* bioreporter during a 90 minute incubation period. A dual strain whole-cell bacterial bioreporter assay was performed using the same soil-water extracts as above, as described previously (Brandt et al., 2008). No sample matrix correction factors were required.

2.6. Element measurement by Inductively Couple Plasma-Mass Spectrometry

Water-extractable, aqua regia and CISED samples were diluted (1:10) with HNO_3 , spiked with $5 \mu g mL^{-1}$ of internal standard mix (Se, Ge, Rh, and Bi), and subsequently measured for elements listed above using an Inductively Couple Plasma-Mass Spectrometry (NexION® 350D ICP-MS, Perkin Elmer). The instrument was calibrated with 9 major (Ca, Fe, K, Mg, Mn, Na, S, Si, P) and 17 trace (Al, As, Ba, Cd, Co, Cr, Cu, Hg, Li, Mo, Ni, Pb, Sb, Se, Sr, V, Zn) elements. The concentration ranged from 1 to $40 mg L^{-1}$ for major elements and 0.01 to $2 mg L^{-1}$ for trace elements. For data quality control, acid blanks (1% nitric acid), digestion blank, and guidance material (BGS102) were analyzed systematically, to account for blank contamination, sensitivity, operating conditions and extraction accuracy. No recovery corrections were made as the mean repeatability (expressed as relative standard deviation %) of the guidance material (BGS102) was below 15% for individual elements. Sample 8 was removed from subsequent data analyses, as it did not meet the data quality control criteria.

2.7. CISED self-modelling mixture resolution

The self-modelling mixture resolution (SMMR) algorithm (MatLab Version R2015a) was used to model the solid-phase distribution of elements in soil as studied by the CISED sequential extraction technique (Cave et al., 2004). The algorithm assumes that the chemical analysis data represents different proportions of chemical components (which have fixed chemical composition) that have been dissolved out of the soil by the extraction media. The algorithm then uses chemometric methods to identify the number of components being extracted, the chemical composition (Al, As,

Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mo, Mn, Na, Ni, P, Pb, S, Sb, Si, Se, Sr, V and Zn) of each component (CMP = composition) and the amount of each component in each extraction solution. The extraction profile of each component is derived from the amount of each component found in the 14 extracts (PRF = profile). In addition, the relative proportions of each element in each component can also be calculated (DST = distribution). The SMMR algorithm was run separately for each sampling location, including the 4 quadrants, resulting in 10 distinct sets of components (6-13/sampling location), resulting in a total 89 individual components in the entire dataset. From here, hierarchical clustering, geochemical profile interpretations and previous study with a certified reference material (Cave et al., 2004; Wragg and Cave, 2012) was used to classify the 89 components into the common, distinct soil phases (i.e. residual pore salts and exchangeable phases). Briefly, a matrix with mean-centered and scaled CMP and PRF, as described above, of each component was performed with Ward's method using the 'agnes' function in the cluster package (Maechler et al., 2012) in R (v.3.4.1, R Foundation for Statistical Computing; available at <https://www.r-project.org/>). The result from the clustering was visualized using a heatmap (**Fig. 2**) created using ggplot2, reshape2, grid, and ggdendro packages (Wickham, 2007; Wickham, 2010).

2.8. Statistical analyses

All statistical analyses were performed in R (v.3.4.1, R Foundation for Statistical Computing). The significant influence of specific physico-chemical soil parameters on the solid phase distribution of Cr, Cu and As was tested using redundancy analyses (RDA) (Legendre and Anderson, 1999). In the context of this research, it tested the relationship between physico-chemical properties of soil on the solid phase distribution of either Cr, Cu or As (separate analysis for each element). In order to choose which explanatory variables (physico-chemical factors) to include in the model, we pre-selected physico-chemical parameters based on geological interpretation of the system, keeping in mind that one of the central assumptions of this analysis is independence between explanatory variables. Variables selected for hypothesis testing were based on evidence from the literature and

included pH, total C, reactive Fe oxide, Olsen P, total P as well as total and water-extractable Ca, Mg and Mn. A forward selection of variables using the 'ordistep' function in the 'vegan' package was performed in order to select the variables to include in the final model. The final model was built using the 'rda' and 'anova.cca' functions of the 'vegan' package (Oksanen et al., 2013). Spatial autocorrelation between samples due to sampling design tested positive using the 'mso' function in the 'vegan' package. Thus, it was accounted for using partial redundancy analysis, conditioning for sampling location. Reported adjusted redundancy statistics R^2 were obtained using the 'RsquareAdj' function of the 'vegan' package.

The relationship between elements in the water-extracts, bioavailable Cu and the common CISED identified soil phases was not reported as it is generally assumed that soil particle associated-elements are not directly available to bacteria (Brandt et al., 2006; Noll, 2003). Therefore, we tested the relationships between bioavailable Cu ($\mu\text{g g}^{-1}$), water-extractable Cr, Cu and As ($\mu\text{g g}^{-1}$) and elements ($\mu\text{g g}^{-1}$) in the most easily extractable soil component identified by CISED for each sampling location by using Pearson correlation coefficient with the 'cor' function in R.

3. Results

3.1 Soil physico-chemical properties

Selected physico-chemical soil properties are summarized in **Table 1** and detailed characterization is provided in **Supplementary Table 1** and **Table 2**. Total concentrations of Cr, Cu and As ranged between 26.1-1819 mg kg^{-1} , 17.2-2205 mg kg^{-1} and 32.4-2839 mg kg^{-1} , respectively, while water-extractable concentrations ranged from 0.02-0.78 mg Cr kg^{-1} , 0.11-5.99 mg Cu kg^{-1} and 0.17-18.3 mg As kg^{-1} and bioavailable Cu ranged from 0.04-3.52 mg kg^{-1} .

Soil samples collected from location 1, 2, 3, 5, 6 and 8 were classified as sandy soils (sand > 85.0%) while 4, 7, 9 and 10 were loamy sand (sand = 81.0- 85.0 %). pH ranged from 3.53-6.83

across all samples. Total C, ranged from 1.47-17.4% and total N from 0.09-0.83%. Reactive Fe oxides in samples ranged from 1.48-5.97 mg kg⁻¹, Olsen P 14.7-73.4 mg kg⁻¹ and total P 319-713 µg g⁻¹. Total and water-extractable Ca ranged from 1112-7414 µg g⁻¹ and 3.42-50.8 µg g⁻¹, respectively. Total and water-extractable Mg ranged from 692-1951 µg g⁻¹ and 0.66-6.72 µg g⁻¹, respectively. Total and water-extractable Mn ranged from 225.4-1174 µg g⁻¹ and 0.08-2.45 µg g⁻¹, respectively.

3.2 Soil solid phases identified by CISED

Extraction recoveries with the CISED extraction methodology compared with total element concentration averaged 61 ± 13 %, 90 ± 11 % and 81 ± 11 % for Cr, Cu and As, respectively. The cumulative CISED extraction recoveries were less than 100 % because the CISED extraction protocol mainly targets the easily soluble surface coatings and not the silicate matrix of soil.

The SMMR algorithm identified 6-13 soil components per sampling location giving 89 individual components in the entire dataset. Using a combination of the chemical composition of the clusters, the relative ease of their extraction as indicated by the amount extracted at each step and a general knowledge of soil geochemistry, 6 common physico-chemical phases were identified (**Fig. 2**).

Table 2 describes the 6 common soil solid phases amongst the sampling locations, listed here in decreasing ease of extractability: Residual pore salts, exchangeable, carbonates (tentative designation), Mn-Al oxide, Fe oxide (amorphous) and Fe oxide (crystalline) (**Table 2**).

Residual pore salts-this cluster was extracted primarily during the first two extractions of CISED using deionized water. It is dominated by Ca, Na, K, and Si, which are highly mobile elements and can be derived from soluble pore water salts in soil.

Exchangeable-this cluster does not result in defined peaks like the other phases, likely as a result of data processing of soil components of higher variability, and is comprised of several components from each location. It is mainly comprised of exchangeable cations such as Na, Mg, S, Sb, Se and P, which are typically bound to soil particle surfaces by weak electrostatic forces and can be displaced by competing cations and released into solution, as exhibited in the saw toothed pattern of the

extraction profile (**Table 2**). Some of these extractable components also come out at higher acid strengths, conceivably being encased in more recalcitrant components which are only being released later on in the extraction profile.

Carbonates (tentative designation)- typically we would designate this component as carbonates however, due to the inherent physico-chemical properties (pH below 7) on site, which would suggest that carbonates are unlikely to occur, we only tentatively assign this soil phase. This is further discussed in Supplementary Material. This cluster is extracted on the first addition of acids and is comprised of two sub-clusters which are carbonate (Ca-Mg) and carbonate (Ca). Carbonate (Ca-Mg) sub-cluster is dominated with Ca and Mg, and comes out over a narrow range in the extractions, primarily during the addition of 0.01M nitric acid. Other elements present in this partition include Al, Ba, Cd, Cu, Sr and Zn. Carbonate (Ca) sub-cluster comes out a little later in the extraction profile, indicating that this carbonate phase is less available than the previous. It is composed of the same elements as the previous cluster, with the notable absence of Mg. Location 3, 4, 5, 8 and 10 have soil components in both sub-clusters while Location 1, 2, 6, 7 and 9 only have soil components associated with more readily available carbonate phase (Ca-Mg).

Mn-Al oxide- this cluster is dominated by Mn and Al elements and tends to be extracted upon addition of peroxide, a reagent commonly known to target Al oxides.

Two distinct clusters of Fe oxides were identified and suggest the presence of both amorphous and crystalline Fe oxides:

Fe oxide (amorphous)- this cluster comprises soil components that are extracted from 0.1M-1M nitric acid and are primarily made up of Fe and to a lesser extent, Al, Pb, Se and V.

Fe oxide (crystalline)- this cluster is comprised primarily of Fe and to a lesser degree Al and Si, and requires high acid concentrations (0.5-5M) to extract. Within this cluster, there is a subcluster with 3 soil components belonging to the locations with the highest contamination (Location 3, 4 and 5) and appear to be a CCA component, primarily made up of Fe, Cr and As.

3.3 Solid Phase Distribution of Cr, Cu and As in soil

Chromium, Cu and As at this site were associated with specific soil phases along the contamination gradient. **Fig. 3** presents the data based on percentage of the individual element in the soil phases as compared with the total Cr, Cu, As extracted in that sample with CISED and as such, the changes observed are relative. The distribution of Cr, Cu and As ($\mu\text{g g}^{-1}$) in each soil phase can be found in **Supplementary Table 3**. Samples were separated into low, moderate and high contamination the subsequent text in order to discuss their distribution along the contamination gradient, as follows: (i) Low concentrations, samples 1-14 (below $175 \mu\text{g g}^{-1}$ for Cr, Cu and As), (ii) moderate concentrations, samples 15-27 (between 175-200 for Cr, 175-600 for Cu and 175-400 $\mu\text{g g}^{-1}$ for As), and (iii) high concentrations, samples 28-40 (above 200 for Cr, 600 for Cu and 400 $\mu\text{g g}^{-1}$ for As).

Cr is almost exclusively found in the least reactive phase, in the crystalline Fe oxide. In the low contamination, particularly in samples coming from location 9, Cr is present in the exchangeable (avg. 78 ± 11 %) and Mn-Al oxide (avg. 22 ± 11 %) phases. In the high contamination, at location 3, some Cr is found in the exchangeable phase (avg. 42 ± 10 %). In the low and moderate contamination, some Cr is found in the amorphous Fe oxide phase (avg. 18 ± 8 %) at location 8 and 2.

Cu is primarily found in Mn-Al oxides and to a lesser extent, the Fe oxide (amorphous) phases in the lower end of the contamination gradient. In the moderate and high contamination, Cu is almost exclusively found in the carbonate (tentative designation) phase (avg. 57 ± 29 %). It is primarily found in the less extractable carbonate sub-cluster, carbonate (Ca). At location 9 (avg. 33 ± 25 %) as well as locations 3 (avg. 17 ± 8 %) and 4 (avg. 18 ± 6 %), a portion of the Cu is found in the exchangeable phase.

The As solid phase distribution pattern along the contamination gradient was more variable. At low contamination, it was primarily in the crystalline Fe oxides (avg. 46 ± 21 %), in the Mn-Al oxides

(avg. 10 ± 8 %), and amorphous Fe oxides (avg. 16 ± 6 %). For samples from location 9, it is also in the exchangeable phase (avg. 24 ± 23 %). In the moderate contamination, the As contamination shifts into the exchangeable (avg. 18 ± 13 %), Mn-Al oxides (avg. 18 ± 13 %) and amorphous Fe oxides (avg. 34 ± 22 %). In the high contamination, the As was no longer found in the Mn-Al oxides but in the carbonates (tentative designation) (avg. 22 ± 7 %), amorphous Fe oxides (avg. 18 ± 10 %), crystalline Fe oxides (avg. 33 ± 11 %) and also in the exchangeable (avg. 24 ± 3 %) for locations 3 and 4. In the moderate (avg. $4 \pm 3 \mu\text{g g}^{-1}$) and high contamination (avg. 2 ± 1 %), a small proportion of As could be detected in the residual pore salts, corresponding to alarmingly high concentrations in specific samples (up to $37.5 \mu\text{g g}^{-1}$).

3. 4 Factors affecting the Solid-Phase Partitioning of Cr, Cu and As in Soil

Specific physico-chemical parameters significantly affected the solid-phase partitioning of Cr, Cu and As in soil as shown with partial redundancy analyses (**Fig. 4**). Cr solid phase distribution was significantly affected by pH ($F=10.3$, $p=0.002$), with the model adjusted $R^2=0.194$ (**Fig. 4a**). The amount of variance explained by this model was 20.9% and an additional 6.1% of the variation could be attributed to spatial autocorrelation correction. Cu solid phase distribution was significantly affected by total Mg ($F=29.1$, $p=0.001$), Olsen P ($F=24.9$, $p=0.001$), pH ($F=22.4$, $p=0.001$), total Mn ($F=14.0$, $p=0.003$), water-extractable Mg ($F=13.3$, $p=0.001$) and water-extractable Ca ($F=6.2$, $p=0.018$), with the model adjusted $R^2=0.759$ (**Fig. 4b**). The amount of variance explained by this model was 70.0% and an additional 8.1% of the variation could be attributed to spatial autocorrelation correction. As solid phase distribution was significantly affected by pH ($F=24.3$, $p=0.001$), reactive Fe oxides ($F=5.2$, $p=0.022$) and Olsen P ($F=3.4$, $p=0.05$), with the model adjusted $R^2=0.437$ (**Fig. 4c**). The amount of variance explained by this model was 46.5% and an additional 9.0% of the variation could be attributed to spatial autocorrelation correction.

3. 5 Linking Cr, Cu and As solid phases to bioavailability

While water-extractable Cu ($r=0.374$, $p=0.019$) and As ($r=0.612$, $p<0.001$) were significantly correlated to the most readily available CISED component at each site, water-extractable Cr and bioavailable Cu were not.

4. Discussion

To our knowledge, this is the first study to investigate and elucidate specific effects of physico-chemical parameters on the solid phase distribution of elements at a complex field site after substantial ageing processes. Element-specific distributions and behaviors in soil were observed along a CCA-contamination gradient at former wood impregnation site.

4.1 Factors affecting the Solid-Phase Partitioning of Cr, Cu and As in Soil

Although past studies have already demonstrated that pH can alter the distribution of these individual elements in soil (Kabata-Pendias, 2010; Lu et al., 2005), here we report the importance of pH in affecting the distribution of all three elements simultaneously along a field contamination gradient. These findings suggest that acidification of soil, as a result of events such as rainwater leaching, decomposition of organic matter and release of carbon dioxide and/or plant root exudation of reducing and chelating compounds for example, could lead to redistribution of Cr, Cu and As amongst soil solid phases.

Additionally, element-specific influences were observed. Cu distribution was influenced by divalent cations such as Ca^{2+} , Mg^{2+} and Mn^{2+} , which are known to be structurally similar and may compete with Cu for binding sites in soil. For example, competition for binding sites and translocation into plant roots between divalent cations such as Ca and Mn has been reported with Cd (Eller and Brix, 2016). Plant available P (Olsen P) was also shown to be important in influencing the distribution of Cu, and could compete with Cu for binding to different oxides, such as Mn-Al oxides. In the case of As, reactive Fe oxides were significantly affecting its distribution, a phenomenon generally known (Yang, Barnett et al. 2002). Available P (Olsen P) also played a role in the distribution of As on soil

solid phases. Interestingly, As(V) is a well-known phosphate analog which competes for sorption sites in soil (Bolan et al., 2013). Thus, an increase in P could result in an increase in available As for plant uptake, as demonstrated at a CCA site by Cao et al. (2004). Indeed, this suggests that sudden changes in these specific physico-chemical factors could lead to shifts in the distribution of these elements between soil liquid and solid phases, as discussed in the subsequent section 4.2.

4.2 Solid Phase Distribution of Cr, Cu and As in soil

In the wood impregnation mixture, Cr was found as Cr(VI), a highly reactive, mobile and toxic form of Cr. During the fixation process, it is reduced to Cr(III) but residues of Cr(VI) may still occur and leach into the soil. After a substantial ageing period, Cr on site was found almost exclusively bound to the least extractable phase, the crystalline Fe oxide. Although the crystalline Fe oxide minerals typically have lower adsorption capacity than amorphous phases such as ferrihydrite minerals, the dominance of Cr in the aforementioned phase could be for several reasons. Either there are more of these oxides present at this site and therefore a larger fraction is absorbed onto this fraction, and/or perhaps more importantly, ferrihydrites age to crystalline Fe oxide over time, a process which can be further aided by microbes in soil (Zachara et al., 2002) and as such, Cr may have started off associated with ferrihydrites but this fraction became crystalline over time. As a result, and in line with a previous report at this site (Nielsen et al., 2016), we suggest that the dominant form of Cr in the top soil is Cr (III) because in this form, Cr is typically bound to positively charged surfaces such as Fe and Al oxides and is very stable in soil (Namiesnik and Rabajczyk, 2012). Although past laboratory and field studies have suggested that Cr from the CCA mixtures is highly mobile, these studies have focused on short-term monitoring (Allinson et al., 2000; Hopp et al., 2006). After 100 days of ageing, however, Carey et al. demonstrated in a controlled leaching study, that less than 1% of Cr(VI) could be extracted in the top layer (Carey et al., 2002). Groundwater investigations of the terrain magazines at this site have suggested that the mobile pools of Cr have been washed out, as concentrations in the groundwater have fallen

significantly between 1977 and 1995, from 21-1500 $\mu\text{g L}^{-1}$ to $<3 \mu\text{g L}^{-1}$ (Nielsen et al., 2010).

Remaining Cr(VI) in the top soil was likely reduced to chromic cation Cr(III) by soil organic matter, as previously shown (Carey et al., 2002; Kabata-Pendias, 2010). Although total concentrations of Cr at the higher end of the contamination gradient exceed guideline values, it is unlikely to leach and cause adverse ecological impact because of its association with crystalline Fe oxide structures which are stable under oxic top soil conditions.

Cu, added as CuO salts in the impregnation mixture, is typically found as a divalent cation in soil. After 40 years of ageing, Cu(II) occurred in readily available phases such as exchangeable, and less available phases such as carbonates (tentative designation) and Mn-Al oxides. Previously, Cu(II) in the oxic top soil has been shown to be highly reactive with soil organic matter (SOM), typically exhibiting low mobility in soil (Andersen et al., 1996; Kabata-Pendias, 2010). At this site, however, soil organic matter content is generally low (avg. $5.6\pm 4.2\%$), particularly in the highly contaminated end of the gradient where substantial amounts of Cu are bound (avg. $2.0\pm 0.5\%$), suggesting that Cu might not efficiently bind to SOM. In recent work at this contaminated site, Frick et al. (*submitted*) also found that dissolved organic matter content in soil from the highly contaminated samples was very low. Indeed, in their artificially CCA contaminated soils with low organic content, Balasoïu et al., found that Cu was predominantly bound to soluble or exchangeable soil phases (Balasoïu et al., 2001). In samples from the low contamination gradient at this site, Cu binds readily to Mn-Al oxides and amorphous Fe oxide, which has been demonstrated in several soils with low organic matter soils (Agbenin and Olojo, 2004; Yu et al., 2004). When moving towards the moderate to highly contaminated samples, Cu shifted into the more extractable phase, carbonates (tentative designation) (primarily the less extractable carbonate sub-cluster). Indeed, the differential vegetation cover and, changes in plant species and physiology between the sampling location on site will affect the geochemistry through direct (e.g. input and quality of organic matter, oxygen content and water holding capacity) and indirect effects on the soil fauna by processes such as reduced organic material breakdown by microorganisms and/or increased compaction of soil due to reduced earthworm activity

(Arthur et al., 2012). The absence of trees, as seen in the higher end of the contamination gradient which can only support the growth of mosses, can lead to increased mobility of elements via absence of interception of incident precipitation, root accumulation and precipitation of elements in the rhizosphere and could in part explain the shift of Cu into a more extractable phase. Determining the isolated impact of individual effects discussed above on the solid phase distribution of elements, however, is nearly impossible as these processes add-up and merge to form what is called the legacy effect. Although carbonate associated elements are generally considered non-available to plants, a drop in pH could release Cu to more readily-extractable phases (Martinez and Motto, 2000). In the highly contaminated samples, Cu was also associated to the exchangeable phase at specific locations, a phase which has been described as readily available to plants and microorganisms (Maderova et al., 2011). Furthermore, the cation characteristics of this phase (e.g. Na, Mg and S), are bound to soil particle surfaces by weak electrostatic forces and can be displaced by competing cations (Rowell, 1994). Indeed, this study has reported that competing divalent cations such as Ca^{2+} , Mg^{2+} and Mn^{2+} play a significant role in the distribution of Cu and as such, the divalent cations in this phase could compete for binding with Cu, leading to the release of Cu in the soil solution. In this study we report that Cu, particularly in highly contaminated samples, is present in soil phases that are readily extractable and/or in phases that are could replenish exchangeable and soil pore water element concentrations, under altering physico-chemical conditions (Degryse et al., 2009).

Arsenic was added as arsenate (As(V)) in the impregnation mixture, which is the less mobile and toxic inorganic form of As as compared with arsenite (As(III)) (Masscheleyn et al., 1991). At this CCA site (Frick et al., *submitted*) and in other studies (BalasoIU et al., 2001; Gräfe et al., 2008; Hopp et al., 2008), As(V) is reported as the dominant form of As, which is consistent with our knowledge that As (V) dominates in oxic top soil environments (Kabata-Pendias, 2010). Along our contamination gradient, we observed that As was associated with all soil solid phases identified with the CISED methodology. A large portion of the As on site was associated with Mn-Al and Fe oxides, in line with our knowledge that the inorganic oxyanion As(V) strongly binds to positively

charged Fe and Al oxides, both in natural (Lin and Puls, 2000; Wragg et al., 2014) and contaminated soils (Bhattacharya et al., 2002). Arsenic bound in these phases is least likely to be dissolved in the soil solution and leach to groundwater aquifers. But, As is also associated with readily extractable phases such as residual pore salts and exchangeable. Arsenic concentrations in those phases, reported at up to 37 and 437 $\mu\text{g g}^{-1}$, respectively, are particularly alarming, given that the soil guidelines for ecological risk are set at 20 $\mu\text{g g}^{-1}$. Like Cu, As is also found in the carbonates (tentative designation) phase and changes in specific physico-chemical factors such as pH, reactive Fe oxides or available P, as described in section 4.1, could lead to redistribution of As into different soil phases and thus, increase the risk of leaching and contamination of groundwater reservoirs.

4.3 Linking Cu and As in solid phases to bioavailability

There is consensus in the literature that insignificant fractions of soil particle associated-Cu are directly available to bacteria (Brandt et al., 2006; Noll, 2003; Ore et al., 2010). As a result, soil pore water elements are often used to represent the biologically relevant element fraction (Giller et al., 2009; Peijnenburg et al., 2007). This study found that although water-extractable Cu correlated to Cu in the most extractable CISED component, bioavailable Cu measured with a bacterial bioreporter, did not. This is in line with knowledge that bioavailability of Cu is influenced by complexation with dissolved organic matter (Brandt et al., 2008; Nybroe et al., 2008). Results from sequential extraction schemes have been used in the past to make inferences about bioavailability and toxicity to soil dwelling organisms such as plants, invertebrates and microorganisms. Our study has shown, however, that bioavailability of Cu to microorganisms cannot be fully inferred using sequential extraction methodologies such as CISED. Indeed, Maderova et al. (2011) also concluded that no single chemical method, including the BCR sequential extraction methodology they employed in their study, could quantify bioavailability of Cu to microbes. This suggests that it is

imperative to have receptor-targeted tests of bioavailability and highlights the importance of including these tests alongside chemical methods.

Water-extractable As was also significantly correlated with As in the most extractable CISED phase, and interestingly, past research at this site reported that water-extractable As was almost exclusively bioavailable, as measured with an arsenic bioreporter assay (Frick, 2016). Collectively these findings suggest that while no inferences can be made about Cu bioavailability from Cu in the CISED residual pore salt phase, As bioavailability to bacteria could tentatively be inferred at this contaminated site.

5. Conclusion/Summary

In this study we demonstrated that physico-chemical factors governing the solid phase distribution of Cr, Cu and As at a CCA-contaminated site are element-specific. Hence, pH and elements competing for binding sites, P in the case of As and divalent cations Ca, Mg and Mn for Cu, affected this distribution. Changes in specific physico-chemical factors through natural processes such as soil acidification from rainwater leaching, organic matter degradation and/or plant exudation or targeted remediation approaches which increase available P or reactive Fe oxides in the field, could alter the solid phase distribution and therefore, mobility of Cr, Cu and As. After substantial ageing processes, Cu and particularly, As remained in readily extractable phases such as residual pore salts and exchangeable while Cr was largely bound to the least extractable solid phase, crystalline Fe oxides. Findings from this study suggest that Cu and As at aged CCA-contaminated sites constitute the largest risks for environmental and human health and should be closely monitored. In the case of Cu, risk is mainly related to ecotoxicological impact while As, due to its higher mobility and severe human health effects, may represent a risk to adjacent aquatic ecosystems and to human health via contamination of drinking water reservoirs. In addition, Cu bioavailability to microbes could not be predicted solely using a sequential extraction procedure.

This suggests that receptor-specific tests of bioavailability such as whole cell bacterial bioreporters for microbes are needed to infer bioavailability in the soil environment and should be integrated into risk assessments to complement analytical methods.

Acknowledgements/Funding

This work was supported financially by REMEDIATE (Improved decision-making in contaminated land site investigation and risk assessment) Marie-Curie Innovation Training Network from the European Union's Horizon 2020 Program (grant agreement n. 643087). In addition, we acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC), [PGS D 396155934]. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), [PGS D 396155934]. The author's would like to thank the Danish Soil Partnership for facilitating access to the contaminated site.

Conflict of Interests

The authors declare no conflicts of interests

References

- Agbenin J, Olojo L. Competitive adsorption of copper and zinc by a Bt horizon of a savanna Alfisol as affected by pH and selective removal of hydrous oxides and organic matter. *Geoderma* 2004; 119: 85-95.
- Allinson G, Turoczy N, Kelsall Y, Allinson M, Stagnitti F, Lloyd-Smith J, et al. Mobility of the constituents of chromated copper arsenate in a shallow sandy soil. *New Zealand Journal of Agricultural Research* 2000; 43: 149-156.
- Andersen S, Rasmussen G, Snilsberg P, Amundsen C, Westby T. Assessing toxicity and mobilisation of impregnation salts at a contaminated site. *Fresenius' journal of analytical chemistry* 1996; 354: 676-680.
- Arthur E, Moldrup P, Holmstrup M, Schjønning P, Winding A, Mayer P, et al. Soil microbial and physical properties and their relations along a steep copper gradient. *Agriculture, ecosystems & environment* 2012; 159: 9-18.
- Bacon JR, Davidson CM. Is there a future for sequential chemical extraction? *Analyst* 2008; 133: 25-46.

- Balasoiu CF, Zagury GJ, Deschenes L. Partitioning and speciation of chromium, copper, and arsenic in CCA-contaminated soils: influence of soil composition. *Science of the Total Environment* 2001; 280: 239-255.
- Bhattacharya P, Frisbie S, Smith E, Naidu R, Jacks G, Sarkar B. Arsenic in the environment: a global perspective. *Handbook of heavy metals in the environment*. Marcell Dekker Inc., New York 2002: 147-215.
- Bolan N, Mahimairaja S, Kunhikrishnan A, Choppala G. Phosphorus–arsenic interactions in variable-charge soils in relation to arsenic mobility and bioavailability. *Science of the Total Environment* 2013; 463: 1154-1162.
- Brandt KK, Holm PE, Nybroe O. Bioavailability and toxicity of soil particle- associated copper as determined by two bioluminescent *Pseudomonas fluorescens* biosensor strains. *Environmental Toxicology and Chemistry: An International Journal* 2006; 25: 1738-1741.
- Brandt KK, Holm PE, Nybroe O. Evidence for Bioavailable Copper– Dissolved Organic Matter Complexes and Transiently Increased Copper Bioavailability in Manure-Amended Soils as Determined by Bioluminescent Bacterial Biosensors. *Environmental science & technology* 2008; 42: 3102-3108.
- Cao X, Ma LQ. Effects of compost and phosphate on plant arsenic accumulation from soils near pressure-treated wood. *Environmental Pollution* 2004; 132: 435-442.
- Carey P, Bidwell V, McLaren R. Chromium (VI) leaching from large undisturbed soil lysimeters following application of a simulated copper-chromium-arsenic (CCA) timber preservative. *Soil Research* 2002; 40: 351-365.
- Cave MR, Harmon K. Determination of trace metal distributions in the iron oxide phases of red bed sandstones by chemometric analysis of whole rock and selective leachate data. *Analyst* 1997; 122: 501-512.
- Cave MR, Milodowski AE, Friel EN. Evaluation of a method for identification of host physico-chemical phases for trace metals and measurement of their solid-phase partitioning in soil samples by nitric acid extraction and chemometric mixture resolution. *Geochemistry: Exploration, environment, analysis* 2004; 4: 71-86.
- Cave MR, Wragg J. Measurement of trace element distributions in soils and sediments using sequential leach data and a non-specific extraction system with chemometric data processing. *Analyst* 1997; 122: 1211-1221.
- Cipullo S, Snapir B, Tardif S, Campo P, Prpich G, Coulon F. Insights into mixed contaminants interactions and its implication for heavy metals and metalloids mobility, bioavailability and risk assessment. *Science of The Total Environment* 2018; 645: 662-673.
- Cornell R, Schindler P. Photochemical dissolution of goethite in acid/oxalate solution. *Clays Clay Miner* 1987; 35: 347-352.
- Cornell RM, Schwertmann U. *The iron oxides: structure, properties, reactions, occurrences and uses*: John Wiley & Sons, 2003.
- Cox SF, Chelliah MC, McKinley JM, Palmer S, Ofterdinger U, Young ME, et al. The importance of solid-phase distribution on the oral bioaccessibility of Ni and Cr in soils overlying Palaeogene basalt lavas, Northern Ireland. *Environmental geochemistry and health* 2013; 35: 553-567.
- Degryse F, Smolders E, Parker D. Partitioning of metals (Cd, Co, Cu, Ni, Pb, Zn) in soils: concepts, methodologies, prediction and applications—a review. *European Journal of Soil Science* 2009; 60: 590-612.
- Directive E. Commission Directive 2003/100/EC of 31 October 2003 amending Annex I of Directive 2002/32/EC of the European Parliament and of the Council on undesirable substances in animal feed. *Off. J. Eur. Union* 2003; 50.
- Eller F, Brix H. Influence of low calcium availability on cadmium uptake and translocation in a fast-growing shrub and a metal-accumulating herb. *AoB Plants* 2016; 8.
- Frick HM, Tardif S, Kandeler E, Holm PH, Brandt KK. Assessment of biochar and zero-valent iron for in-situ remediation of chromated copper arsenate contaminated soil. *STOTEN* (submitted)

- Frick HM. Remediation of chromated copper arsenate (CCA) contaminated sites with biochar and zero-valent iron. Faculty of Science. Master's thesis. University of Copenhagen, 2016.
- Gee GW, Bauder JW. Particle-size analysis. *Methods of soil analysis: Part 1—Physical and mineralogical methods* 1986; 383-411.
- Gezer ED, Cooper PA. Effects of wood species and retention levels on removal of copper, chromium, and arsenic from CCA-treated wood using sodium hypochlorite. *Journal of forestry research* 2016; 27: 433-442.
- Giller KE, Witter E, McGrath SP. Heavy metals and soil microbes. *Soil Biology and Biochemistry* 2009; 41: 2031-2037.
- Gräfe M, Tappero RV, Marcus MA, Sparks DL. Arsenic speciation in multiple metal environments: II. Micro-spectroscopic investigation of a CCA contaminated soil. *Journal of Colloid and Interface Science* 2008; 321: 1-20.
- Hingston J, Collins C, Murphy R, Lester J. Leaching of chromated copper arsenate wood preservatives: a review. *Environmental Pollution* 2001; 111: 53-66.
- Hopp L, Nico PS, Marcus MA, Peiffer S. Arsenic and chromium partitioning in a podzolic soil contaminated by chromated copper arsenate. *Environmental science & technology* 2008; 42: 6481-6486.
- Hopp L, Peiffer S, Durner W. Spatial variability of arsenic and chromium in the soil water at a former wood preserving site. *Journal of contaminant hydrology* 2006; 85: 159-178.
- Humphrey DG. The chemistry of chromated copper arsenate wood preservatives. *Reviews in inorganic chemistry* 2002; 22: 1-40.
- Kabata-Pendias A. Trace elements in soils and plants: CRC press, 2010.
- Legendre P, Anderson MJ. Distance- based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological monographs* 1999; 69: 1-24.
- Lin Z, Puls R. Adsorption, desorption and oxidation of arsenic affected by clay minerals and aging process. *Environmental Geology* 2000; 39: 753-759.
- Loeppert RH, Suarez DL. Carbonate and gypsum. 1996.
- Lu A, Zhang S, Shan X-q. Time effect on the fractionation of heavy metals in soils. *Geoderma* 2005; 125: 225-234.
- Maderova L, Watson M, Paton G. . *Soil Biology and Biochemistry* 2011; 43: 1162-1168.
- Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. Cluster: cluster analysis basics and extensions. R package version 2012; 1: 56.
- Manning BA, Goldberg S. Modeling competitive adsorption of arsenate with phosphate and molybdate on oxide minerals. *Soil Science Society of America Journal* 1996; 60: 121-131.
- Martinez C, Motto H. Solubility of lead, zinc and copper added to mineral soils. *Environmental Pollution* 2000; 107: 153-158.
- Masscheleyn PH, Delaune RD, Patrick Jr WH. Effect of redox potential and pH on arsenic speciation and solubility in a contaminated soil. *Environmental science & technology* 1991; 25: 1414-1419.
- Namiesnik J, Rabajczyk A. Speciation analysis of chromium in environmental samples. *Critical reviews in environmental science and technology* 2012; 42: 327-377.
- Nguyen MTH, Postma D, Trang PTK, Jessen S, Viet PH, Larsen F. Adsorption and desorption of arsenic to aquifer sediment on the Red River floodplain at Nam Du, Vietnam. *Geochimica et cosmochimica acta* 2014; 142: 587-600.
- Nielsen SS, Jakobsen R, Kjeldsen P. Lokalitet nr. 219-3 Collstrupgrunden: Udredning vedr. forureningssituationen på og omkring grunden 1977–2009 (In Danish). 2010.
- Nielsen SS, Jakobsen R, Kjeldsen P. Stabilization of arsenic and chromium polluted soils using water treatment residues. 2013.
- Nielsen SS, Kjeldsen P, Hansen HCB, Jakobsen R. Transformation of natural ferrihydrite aged in situ in As, Cr and Cu contaminated soil studied by reduction kinetics. *Applied geochemistry* 2014; 51: 293-302.

- Nielsen SS, Kjeldsen P, Jakobsen R. Full scale amendment of a contaminated wood impregnation site with iron water treatment residues. *Frontiers of Environmental Science & Engineering* 2016; 10: 3.
- Noll MR. Trace elements in terrestrial environments: biogeochemistry, bioavailability, and risks of metals. *Journal of Environmental Quality* 2003; 32: 374.
- Nunes I, Jacquioud S, Brejnrod A, Holm PE, Johansen A, Brandt KK, et al. Coping with copper: legacy effect of copper on potential activity of soil bacteria following a century of exposure. *FEMS microbiology ecology* 2016; 92: fiw175.
- Nybroe O, Brandt KK, Ibrahim YM, Tom- Petersen A, Holm PE. Differential bioavailability of copper complexes to bioluminescent *Pseudomonas fluorescens* reporter strains. *Environmental toxicology and chemistry* 2008; 27: 2246-2252.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R, et al. Package 'vegan'. *Community ecology package, version 2013*; 2.
- Ore S, Mertens J, Brandt KK, Smolders E. Copper toxicity to bioluminescent *Nitrosomonas europaea* in soil is explained by the free metal ion activity in pore water. *Environmental science & technology* 2010; 44: 9201-9206.
- Peijnenburg WJ, Zablotkaja M, Vijver MG. Monitoring metals in terrestrial environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicology and environmental safety* 2007; 67: 163-179.
- Rowell D. *SoilScience: methods & applications.*, (Longman Scientific & Technical: Harlow, UK). *Soil science: Methods and applications.* Longman Scientific and Technical, Harlow, UK. 1994: -.
- Samfundsteknik. *Pollution Investigation of the Former Impregnation Facility, Stenholtsvang – Fase 1* (in Danish). County of Frederiksborg, Denmark (1989) 1989a.
- Samfundsteknik. *Stenholtsvang, Pollution Investigation – Fase 2* (In Danish). County of Frederiksborg, Denmark (1989) 1989b.
- Wickham H. Reshaping data with the reshape package. *Journal of Statistical Software* 2007; 21: 1-20.
- Wickham H. ggplot2: elegant graphics for data analysis. *J Stat Softw* 2010; 35: 65-88.
- Wragg J, Cave M. Assessment of a geochemical extraction procedure to determine the solid phase fractionation and bioaccessibility of potentially harmful elements in soils: A case study using the NIST 2710 reference soil. *Analytica Chimica Acta* 2012; 722: 43-54.
- Wragg J, Cave M, Gregory S. The solid phase distribution and bioaccessibility of arsenic, chromium, and nickel in natural ironstone soils in the UK. *Applied and Environmental Soil Science* 2014; 2014.
- Yang J-K, Barnett MO, Jardine PM, Basta NT, Casteel SW. Adsorption, sequestration, and bioaccessibility of As (V) in soils. *Environmental Science & Technology* 2002; 36: 4562-4569.
- Young SD. *Chemistry of heavy metals and metalloids in soils.* Heavy metals in soils. Springer, 2013, pp. 51-95.
- Yu S, He Z, Huang C, Chen G, Calvert D. Copper fractionation and extractability in two contaminated variable charge soils. *Geoderma* 2004; 123: 163-175.
- Zachara JM, Kukkadapu RK, Fredrickson JK, Gorby YA, Smith SC. Biomineralization of poorly crystalline Fe (III) oxides by dissimilatory metal reducing bacteria (DMRB). *Geomicrobiology Journal* 2002; 19: 179-207.

Figures and Tables




Figure 1: (a) Map of Denmark and location of contaminated field site and (b) field site with cumulative total Cr, Cu and As concentrations measured in 1990 by SAMFUNDSTEKNIK during phase 2 risk assessment (Samfundsteknik, 1989b) with designated 10 sampling locations (1-10).




Figure 2: Heatmap and hierarchal clustering of components identified by self-modelling mixture resolution (SMMR) algorithm for 10 sampling locations. The heatmap color gradient represents the mean-centered concentrations of elements and the extraction profiles (E1-E14) reported as total extracted solids ($\mu\text{g g}^{-1}$) of the individual components, where black is low and white represents high concentrations. Each row represents a component identified by the SMMR algorithm with the name indicating the origin of the sampling location and the major elemental composition ($\geq 10\%$). The stipple line on the ward cluster tree is where the cut-off was made to inform the geochemical assignment of common soil solid phases, listed in decreasing extractability; 1) Residual pore salts, 2) Exchangeable, 3) Carbonates (tentative designation), 4) Mn-Al oxide, 5) Fe oxide (amorphous) and 6) Fe oxide (crystalline).




Figure 3: Distribution (%) of Cr, Cu and As in different geochemical soil phases as compared to total extracted using CISED, in decreasing ease of extractability from the soil samples, along contamination gradient in Fredensborg, Denmark.




Figure 4: The effect of significant ($p \geq 0.05$) physico-chemical factors on the solid phase distribution measured using CISED of (a) Cr (one-dimensional), (b) Cu and (c) As shown using constrained partial redundancy analysis, while controlling for the effect of autospatial correlation of sample.

1 **Table 1:** Selected physico-chemical properties of the top 10 cm of soil sampled ($n=40$) along the contamination gradient in Fredensborg, Denmark

Sample ^a	Sampling Location ^b	Clay ^c	Silt ^c	Sand ^c	Texture ^c	pH	C	N	Olsen P	Reactive Fe oxides	Cu _{Bio} ^d	As _{tot} ^d	Ca _{we} ^d	Cr _{tot} ^d	Cu _{tot} ^d	Mg _{tot} ^d	Mn _{tot} ^d	P _{tot} ^d	As _{we} ^d	Ca _{we} ^d	Cr _{we} ^d	Cu _{we} ^d	Mg _{we} ^d	Mn _{we} ^d
		%			USDA		%			mg g ⁻¹	µg g ⁻¹													
1	9	6	13	81	Loamy sand	4.56	2.93	0.18	20.2	5.97	0.06	32.5	2441	28.9	17.2	1565	521	460	0.17	8.04	0.04	0.11	1.05	0.39
2	7	4.4	13.6	82	Loamy sand	5.27	2.69	0.14	32.1	1.48	0.40	39.8	2538	30.4	32.3	1672	457	403	0.40	18.8	0.13	0.56	4.78	2.45
3	9	6	13	81	Loamy sand	5.01	2.92	0.17	19.4	4.98	0.05	46.1	2524	33.4	24.0	1648	492	443	0.22	7.12	0.02	0.11	0.77	0.20
4	7	4.4	13.6	82	Loamy sand	5.26	2.5	0.13	27.3	1.98	0.25	36.7	2480	26.1	42.4	1622	445	399	0.24	11.2	0.06	0.38	2.27	0.70
5	9	6	13	81	Loamy sand	4.29	3.21	0.19	24.0	2.19	0.10	61.5	1997	30.4	22.9	1334	441	403	0.37	7.35	0.03	0.18	0.97	0.40
6	7	4.4	13.6	82	Loamy sand	5.64	3.83	0.21	27.9	2.2	0.32	48.0	2759	47.1	52.2	1455	447	394	0.67	26.0	0.17	0.70	4.70	1.38
7	7	4.4	13.6	82	Loamy sand	5.29	5.91	0.27	41.2	1.87	0.34	54.0	3137	59.0	52.8	1747	496	492	0.61	15.7	0.13	0.48	2.91	1.02
8	10	4	13	83	Loamy sand	6.42	11.6	0.6	29.6	2.95	0.14	109	3307	67.2	75.3	1550	463	538	2.19	36.1	0.04	0.38	2.02	0.09
9	1	6.4	8	85.6	Sand	4.53	9.55	0.56	14.7	1.73	0.19	109	2869	94.4	70.0	1535	276	455	1.22	13.7	0.15	0.60	1.73	0.47
10	1	6.4	8	85.6	Sand	3.53	12.6	0.69	18.2	1.93	0.21	113	3408	110	69.6	1601	304	486	1.79	17.8	0.21	0.48	2.55	1.03
11	10	4	13	83	Loamy sand	6.73	7.89	0.41	22.1	1.61	0.04	169	4596	94.0	93.9	1627	543	539	2.02	30.6	0.03	0.23	2.72	0.13
12	1	6.4	8	85.6	Sand	3.64	9.15	0.51	19.4	2.02	0.38	143	2847	128	94.6	1520	282	459	1.62	25.8	0.27	0.96	3.08	1.90
13	9	6	13	81	Loamy sand	5.62	4.24	0.25	31.0	1.89	0.09	178	4584	107	137	1584	533	633	0.54	6.31	0.03	0.14	0.98	0.24
14	10	4	13	83	Loamy sand	6.78	12.9	0.62	37.3	1.78	0.05	175	7414	142	122	1591	616	671	1.97	45.6	0.04	0.24	4.96	0.10
15	8	4	9	87	Sand	5.55	10.3	0.56	59.4	1.99	0.18	213	4775	154	158	1707	449	557	5.44	27.4	0.29	0.79	6.72	0.66
16	8	4	9	87	Sand	5.78	11.3	0.58	42.2	2.16	0.62	288	5021	128	149	1787	508	562	4.98	22.7	0.15	0.65	4.95	0.25
17	2	5.6	5.4	89	Sand	5.08	2.63	0.17	73.4	4.52	1.93	251	1399	74.2	289	872	225	516	4.80	5.11	0.13	1.69	1.34	0.70
18	10	4	13	83	Loamy sand	6.75	10.3	0.51	35.0	2.57	0.09	294	6797	161	203	1839	678	712	2.49	41.0	0.04	0.35	2.70	0.09
19	1	6.4	8	85.6	Sand	4.73	11	0.58	20.4	2.15	0.59	284	3637	244	134	1594	339	463	2.94	26.8	0.44	1.15	3.17	1.25
20	8	4	9	87	Sand	5.67	10	0.55	39.0	1.95	0.28	313	5294	169	188	1924	687	699	5.24	24.3	0.15	0.68	5.01	0.31
21	8	4	9	87	Sand	5.96	7.1	0.39	45.3	2.15	0.71	228	6096	213	251	1951	747	645	6.21	23.9	0.22	0.86	5.13	0.32
22	2	5.6	5.4	89	Sand	4.45	2.85	0.17	68.6	2.25	0.87	334	1144	216	281	755	230	384	7.63	3.42	0.23	1.51	1.00	0.69
23	2	5.6	5.4	89	Sand	4.51	2.56	0.16	59.5	1.91	1.53	221	1112	147	493	725	258	428	3.18	3.54	0.14	2.09	0.66	0.46
24	6	5.6	9.4	85	Sand	5.07	5.43	0.29	47.3	2.2	1.00	308	1564	267	371	898	623	576	5.33	4.71	0.29	1.47	1.21	0.51
25	6	5.6	9.4	85	Sand	4.38	7.93	0.41	40.8	2.1	0.72	386	1519	440	468	919	536	567	8.43	5.02	0.72	1.57	1.64	0.93
26	2	5.6	5.4	89	Sand	4.67	2.96	0.18	65.9	1.96	1.22	397	1169	372	561	721	434	438	3.87	4.02	0.21	1.59	0.94	0.60
27	6	5.6	9.4	85	Sand	4.2	17.4	0.83	22.2	2.17	3.52	203	1749	187	1129	692	484	445	7.51	11.2	0.35	5.99	2.36	0.90

28	6	5.6	9.4	85	Sand	4.59	7.03	0.38	34.5	2.44	1.92	404	1925	568	674	864	683	582	5.84	10.9	0.78	2.98	2.71	1.06
29	5	2.4	6.6	91	Sand	6	2.57	0.17	26.3	2.4	1.03	459	1899	283	965	1095	1142	474	4.79	6.53	0.09	1.64	1.03	0.15
30	5	2.4	6.6	91	Sand	6.21	1.88	0.14	26.1	2.67	1.28	470	1818	239	1298	978	834	428	6.52	12.1	0.13	2.69	2.16	0.17
31	3	4.4	4.6	91	Sand	6.83	1.6	0.1	29.3	1.94	0.23	857	3285	336	879	1106	536	356	4.15	36.9	0.11	0.67	2.36	0.08
32	3	4.4	4.6	91	Sand	6.65	1.66	0.09	37.5	2.35	1.31	1088	3692	399	1097	1079	461	361	4.59	50.8	0.09	0.84	2.56	0.10
33	4	4.4	10.6	85	Loamy sand	6.35	2.4	0.12	27.4	3.87	1.01	1139	2400	434	1158	1124	385	319	18.4	15.7	0.29	3.66	4.05	0.86
34	3	4.4	4.6	91	Sand	6.21	1.47	0.09	26.6	2.43	1.34	1097	2509	555	1168	1144	343	363	7.65	37.5	0.24	2.02	3.02	0.16
35	5	2.4	6.6	91	Sand	6.23	1.76	0.11	24.8	1.48	0.76	973	2394	526	1462	1083	869	344	9.09	21.7	0.11	1.62	4.12	0.82
36	5	2.4	6.6	91	Sand	6.08	1.79	0.11	30.4	2.37	0.51	859	2442	456	1935	1172	1174	372	6.45	12.8	0.09	1.60	2.33	0.29
37	3	4.4	4.6	91	Sand	6.54	1.54	0.1	25.1	3.27	1.52	1357	2633	641	1785	977	397	350	7.63	47.3	0.24	2.05	2.70	0.13
38	4	4.4	10.6	85	Loamy sand	6.42	1.7	0.09	29.5	3.6	1.15	1795	4030	263	2205	1239	459	341	11.8	20.3	0.52	4.80	4.83	0.38
39	4	4.4	10.6	85	Loamy sand	6.27	2.16	0.11	23.9	3.49	0.60	2028	3867	1171	1844	1221	445	378	9.51	19.7	0.15	1.69	4.19	0.13
40	4	4.4	10.6	85	Loamy sand	6.26	3.08	0.11	25.3	3.99	0.58	2839	3038	1819	1630	1201	469	341	10.8	12.9	0.31	2.31	3.80	0.37

2

3 ^a Samples were named 1-40 by total cumulative concentrations of Cr, Cu and As in ascending order

4 ^b Four samples were obtained from each sampling location (see Figure 1 for details)

5 ^c Clay, silt, sand and texture classification were performed on composite sample at each sampling location

6 ^d Cu_{bio}=bioavailable Cu as determined by whole-cell bacterial bioreporter assay, tot=total extracted by aqua regia digestion and we= water-extractable
7 concentrations

8

9

10

11






12

13

14

15 **Table 2:** Common soil solid phases identified by CISED sequential extraction along the contamination gradient in Fredensborg, Denmark

16

Soil Phase	Major element composition	Minor element composition	Extraction profile example	References ^a
Residual pore-salts	Al, Ca, Na, K, and S	Si		(Cave et al., 2004; Rowell, 1994)
Exchangeable	K, Na, Mg, S, Si and K	Sb, Se and P		(Rowell, 1994)
Carbonates (tentative designation)	Ca and Mg	Al, Ba, Cd, Cu, Sr and Zn		(Loeppert and Suarez, 1996)
	Ca	Al, Ba, Cd, Cu, Sr and Zn		(Loeppert and Suarez, 1996)
Mn-Al oxides	Mn and Al			(Manning and Goldberg, 1996; Young, 2013)

Fe oxide
(amorphous)

Fe and Al

Pb, Se and V


(Cave et al., 2004; Cave and Wragg, 1997;
Cornell and Schindler, 1987)

Fe oxide
(crystalline)

Fe

Al and Si

(Cave et al., 2004; Cave and Wragg, 1997;
Cornell and Schwertmann, 2003)



17

18 ^aReferences which informed the grouping into common soil phases