

**The potential for constructed wetlands to treat alkaline bauxite residue leachate:
Phragmites australis growth**

Higgins, D.^{1,3}, Curtin, T.^{2,3}, Pawlett, M⁴ & Courtney, R.^{1,3*}

¹ Department of Life Sciences, University of Limerick, Limerick, Ireland.

² Department of Chemistry and Environmental Science, University of Limerick, Limerick, Ireland.

³ Bernal Institute, University of Limerick, Limerick, Ireland.

⁴ School of Water Energy and Environment, Cranfield University, Bedfordshire, U.K

* ronan.courtney@ul.ie Tel: +353 61202427

Abstract

High alkalinity (pH>12) of bauxite residue leachates presents challenges for the long-term storage and managements of the residue. Recent evidence has highlighted the potential for constructed wetlands to effectively buffer the alkalinity, but there is limited evidence on the potential for wetland plants to establish and grow in soils inundated with residue leachate. A pot based trial was conducted to investigate the potential for *Phragmites australis* to establish and grow in substrate treated with residue leachate over a pH range of 8.6-11.1. The trial ran for 3 months, after which plant growth and biomass were determined. Concentrations of soluble and exchangeable trace elements in the soil substrate, and also in the aboveground and belowground biomass were determined. Residue leachate pH did not affect plant biomass or microbial biomass. With the exception of Na there was no effect on exchangeable trace elements in the substrate, however increases in soluble metals (As, Cd, Na) were observed with increasing leachate concentration. Furthermore, increases in Al, As and V were observed in belowground biomass and for Cd and Cr in aboveground biomass. Concentrations within the vegetation biomass were less than critical phytotoxic levels. Results demonstrate the ability for *P. australis* to grow in bauxite residue leachate inundated growth media without adverse effects.

Keywords: Hyperalkaline wastes, macrophytes, metal removal, mine water, passive treatment, red mud

Introduction

Bauxite residue is the waste by-product generated by the extraction of alumina from bauxite ore via the Bayer Process. The annual global production of bauxite residue is about 120 million tonnes, with an estimated global stockpile of 3 billion tonnes (Klauber et al. 2011). The high alkalinity of bauxite residue presents challenging conditions for the long-term environmental management of bauxite residue disposal areas (BRDAs). Where BRDAs are not managed correctly there is potential for leakage to the surrounding environment representing a high environmental risk (Wang et al. 2015). Requirements for the management and treatment of such drainage waters may persist for many decades following closure (Hua et al. 2015).

Treatment of drainage waters by conventional methods is likely to be expensive, especially if it is to be continued for many decades after closure. Recent research has focused attention on the potential for low-cost, passive technology (constructed wetlands) to treat the alkaline leachates from BRDAs (Hua et al. 2015; Buckley et al. 2016) but the focus of these experiments has been on batch trials at laboratory scale.

Wetlands have been effective in buffering alkaline steel slag leachates (Banks et al. 2006; Mayes et al. 2008) and there is emerging evidence of their ability to treat bauxite residue leachate (Hua et al. 2015; Buckley et al. 2016). The ability of vegetation to establish within the stressful environment of bauxite residue leachate wetlands is unknown. Vegetation of wetlands treating alkaline leachates serve several functions such as flow baffling, provision of a large surface area for inorganic precipitates to be lost from solution, and provision of a continued carbon source for the microbial decomposers responsible for elevating CO₂ and thereby the partial pressure of CO₂ (pCO₂) in wetlands (Mayes et al. 2008). Establishment and continued growth of macrophytes is seen as an essential component for an effective wetland system to treat bauxite residue leachates. Potential impacts of alkaline loading on wetland ecosystems include: reduced solubility of micronutrients, reduced microbial activity, reduced availability of potassium and, direct toxicity of the OH⁻ (Mayes et al. 2009).

Alkaline steel slag leachate reduces shoot production, shoot growth and causes higher root:shoot weight ratios in both *P. australis* and *T. latifolia* (Lawson 2004). Similarly, Mayes et al. (2009) observed reduced *T. latifolia* biomass in calcareous substrates receiving

lime spoil drainage (pH up to 12.7). Reduced growth in both of these studies has been attributed to reduced nutrient availability in high pH substrates (Mayes et al. 2009). Other characteristics that can inhibit vegetation growth in the steel slag wetland environments include low organic matter content, calcareous crusts, poor physical structure and elevated boron (Mayes et al. 2009). Microbial community structure and activity within wetlands play important roles in pollutant removal (Vymazal 2005; Truu et al. 2009). Microbial activity also consumes alkalinity but microbe activity within wetlands is often overlooked in short term laboratory trials (Mayes et al. 2009). The effects of residue leachate loading on soil microbial populations are unknown.

In addition to high alkalinity (ca. pH 13 Buckley et al. 2016) bauxite residue leachates often present high salinity (up to 160 mS cm⁻¹) (Mayes et al. 2011). Several oxyanionic forming elements are very soluble at alkaline conditions and high concentrations of Al (500–1000 mg L⁻¹), As (3–5 mg L⁻¹) and V (5–10 mg L⁻¹) can occur in the leachate (Burke et al. 2012; Hua et al. 2015). Contamination of soils with “red mud” bauxite residue has resulted in increased levels of trace elements (Rutyers et al. 2011; Lehoux et al. 2013) but there is limited knowledge on the impact of residue leachate to growth media and soils.

For constructed wetlands to be considered as a potential mechanism for buffering bauxite residue derived leachates, the impact of leachate on plant establishment and growth needs to be assessed. We determined whether residue leachate (pH range 8.6-11.1) affects *Phragmites australis* biomass and nutrient uptake in a 10 week pots based experiment. Potential phytoavailability of residue associated trace elements was also assessed by measuring soluble and exchangeable fractions. Substrate microbial biomass was also determined as an indicator of leachate effects on soil microbial properties.

Materials and methods

Bauxite residue leachate was collected from an operating residue disposal area, filtered (0.45 µm) and trace element content determined using a Perkin-Elmer Elan DRCII inductively coupled plasma-mass spectrometer (ICP-MS; As and Cr) and an Optima 5300 DV inductively coupled plasma optical emission spectrometer (ICP-OES; all other elements) (Table 1). Dilutions were prepared in the laboratory representing the anticipated range of closed BRDA leachate (Buckley et al. 2016) (Table 2).

Juvenile *Phragmites australis* plants were provided by a local nursery and were planted in 1 L pots. Growth medium used was a proprietary all-purpose compost (pH 6, organic content 86%), as a previous study by Buckley et al. (2016) demonstrated greater potential for buffering pH of NaOH solutions in high organic matter substrates. All plants were pre-grown for one month to acclimatise them to their new environment, which was in a nutrient solution containing: KNO_3 (1.5 mmol L⁻¹), $\text{Ca}(\text{NO}_3)_2$ (1 mmol L⁻¹), $\text{NH}_4\text{H}_2\text{PO}_4$ (0.5 mmol L⁻¹), $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$ (0.25 mmol L⁻¹), KCl (1 mmol L⁻¹), H_3BO_3 (25 mmol L⁻¹), $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ (0.1 mmol L⁻¹), $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot 4\text{H}_2\text{O}$ (mmol L⁻¹) and $\text{Fe}(\text{Na})\text{EDTA}$ (0.1 mmol L⁻¹) (Matthews et al. 2005). Following the acclimatisation period replicate 1L pots (n=5) were placed in basins and subjected to inundation with residue leachate treatments (Table 2). Following the method of Matthews et al. (2005) leachate treatments were fed to the plants at the same volume and on a continuous basis over a 3-month period ensuring the plant pot was submerged in solution at all times.

Measurements of leaf length were carried out weekly and root length was determined at end of experiment. The length of the longest root was measured, by measuring with a ruler from the initial point of growth to the endpoint of the root at the tip. Dry weight measurements of aboveground and belowground biomass were determined at harvest.

Post treatment the wetland plants were carefully removed from the plastic pots to ensure all roots stayed intact. The majority of the compost substrate mass was removed by hand before the roots were washed. Plant material was placed in an oven at 60°C for 48 hours. Once dry, the plant aboveground and belowground mass was recorded. Plant samples were acid digested in HNO_3 and elemental content determined by ICP (see above).

Compost substrate samples were split and one half were air-dried prior to physico-chemical analysis. Substrate pH and EC determined in 1:5 solutions and elemental content was determined in water soluble and exchangeable (1 M NH₄OAc) extractions. Microbial biomass-C was determined on field fresh (<2mm sieve) substrate using the fumigation-extraction procedure (Jenkinson and Powlson 1976) using the correction factor K_{EC} of 0.45 (Vance et al. 1987; Joergensen 1996).

Table 1. Selected parameters of bauxite residue leachate

	This study	Literature values*
pH	13.2	12.6 -13.1
EC ($\mu\text{S cm}^{-1}$)	67400	-
Al ($\mu\text{g l}^{-1}$)	1077000	352000 - 833500
As ($\mu\text{g l}^{-1}$)	3125	6325- 8140
Cd ($\mu\text{g l}^{-1}$)	3.2	-
Cr ($\mu\text{g l}^{-1}$)	148	65 -188
Ni ($\mu\text{g l}^{-1}$)	35	-
V ($\mu\text{g l}^{-1}$)	13500	8977-15600

*Burke et al. 2012; 2013

Table 2. Dilution rates and pH of residue leachate treatments

Treatment	Dilution	pH
Control	-	6.9
1	1:400	8.5
2	1:300	8.8
3	1:200	9.05
4	1:100	9.9
5	1:75	10.4
6	1:40	11.1

Data were analysed statistically using one-way analysis of variance, descriptive measures and Pearson's bivariate correlations on SPSS, version 19.0. Following the implementation of Kolmogorov– Smirnov one-sample normality tests on SPSS, differences between the residue leachate treatments on substrate and plant properties were individually determined by one-way ANOVA and differences were calculated at the 5% level using Tukey's test. Figures were constructed using GraphPad Prism, Version 6.

Results

After the three month growth trial alkaline residue leachate caused a significant increase in substrate pH that increased with leachate concentration (Fig. 1). At the highest concentration rate of residue leachate (1:40 dilution), pH increase was 0.6 pH units. Substrate EC increased significantly in all treatments with an EC high of 0.72 mS cm^{-1} recorded for the highest leachate application rate (3.5 * that of the control). Soluble levels of Mg and Ca were decreased with increasing application rate of leachate and soluble Na was increased. Increased soluble As and Cd was observed with increasing amounts of leachate additions but this increase was only significant for Cd with the highest leachate pH loading. With the exception of Na, leachate pH did not significantly affect exchangeable elements (Ca, Mg, Cr, Al) (Table 3). Soluble and exchangeable V was below detection limits.

164 **Table 3.** Ammonium acetate extractable elements (mg kg⁻¹) in bauxite residue leachate
 165 treated substrate following *Phragmites australis* growth trial (means ± SE, n = 5)

Treatments	Ca	Mg	Na	Al	As	Ni	Cr
Control	4480±228a	1280±156a	9.5±3a	2.5±0.1a	0.19±0.02a	4.8±0.7a	0.25±0.01a
pH 8.5	3810±229a	1280±25a	93±7.5a	2.4±0.1a	0.35±0.04a	5.0±0.6a	0.26±0.01a
pH 8.8	5300±122a	1830±76a	194±18ab	3.8±0.7a	0.24±0.02a	4.4±0.1a	0.26±0.01a
pH 9.0	4400±201a	1550±119a	243±33b	2.6±0.1a	0.33±0.03a	4.7±0.081	0.26±0.01a
pH 9.9	4110±206a	1420±71a	461±55bc	2.6±0.1a	0.46±0.02a	4.9±0.2a	0.26±0.01a
pH 10.4	4850±221a	1870±127a	848±105c	3.1±0.3a	0.20±0.04a	5.4±0.2a	0.29±0.02a
pH 11.3	4330±131a	1620±111a	2180±174d	2.5±0.1a	0.27±0.03a	5.1±0.1a	0.27±0.01a

166 Means followed by the same letter in a column are not significantly different at P≤0.05

167 **Table 4.** Plant growth parameters for *Phragmites australis* and microbial biomass in
 168 substrate samples (means ± SE, n = 5)

Treatments	Root Biomass (g/pot)	Root length (cm)	Shoot Biomass (g/pot)	Shoot length (cm)	Microbial biomass (µg C/g)
Control	7.6±1.6a	53±10a	1.3±0.3a	56.4±4.9a	780±152a
pH 8.5	5.8±1.1a	43±7.5a	1.2±0.2a	56.5±9.9a	583±278a
pH 8.8	6.5±1.5a	42±5.7a	1.3±0.2a	56.4±4.8a	604±199a
pH 9.0	8.6±2.6a	36±3.7a	2.2±0.5a	74.2±8.6a	460±376a
pH 9.9	8.5±1.6a	38±5.2a	1.9±0.4a	63.5±6.1a	798±178a
pH 10.4	5.7±1.4a	29±2.3a	1.0±0.3a	48.4±8.1a	900±285a
pH 11.3	7.1±1.0a	32±4.2a	1.9±0.2a	68.7±8.3a	516±145a

169 Means followed by the same letter in a column are not significantly different at P≤0.05

170

171

172 Although decreased root length was observed with increasing concentrations of leachate pH
 173 the difference were not significant (Table 4). There was also no significant differences
 174 between treatments for shoot length. Similarly there were no significant differences between
 175 treatments for belowground and aboveground biomass. No significant differences were
 176 observed for microbial biomass.

177

178 There was no significant difference between the treatments for *Phragmites australis* nutrient
 179 content in the belowground samples (Figure 2). Na content significantly increased between
 180 each treatment with the highest leachate application displaying results higher than all other
 181 treatments but no significant differences were observed for aboveground samples.

182 For aboveground biomass samples increased leachate pH did not result in any significant
183 differences for N, K, Na and Mn content (Figure 3). Significantly lower values for Ca, Mg, S
184 and Mn were observed with increased residue leachate pH.

185 No significant differences were observed for Ni, Cd or Cr content in belowground biomass
186 (Figure 4). There was a pattern of increased concentrations with increased rate of leachate
187 for Al, As and V with significant differences for the higher pH treatments. Significant
188 differences were also observed for As and Al content at the highest application rates. For
189 aboveground samples only Cd and Cr displayed significant differences and these were at the
190 higher application rates (Figure 5).

191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209

210 **Discussion**

211 Residue leachate was both alkaline and saline with pH marginally higher than values
212 previously reported (Burke et al. 2012; 2013). Elements determined were also in the range of
213 the previously reported values with the exception of Al which was higher (Burke et al. 2013),
214 possibly due to the slightly higher pH. Whilst there have been no published studies on
215 application of residue leachates to soils and substrates, Lehoux et al. (2013) and Mišík et al.
216 (2014) reported similar findings (increase in pH, increase in salinity and increase in aqueous
217 metal(loid)s concentrations) upon addition of residue to soils.

218 Al and V (except for pH 11.1) concentrations in the current study were below the limit of
219 detection, and soluble levels for other elements determined were below values previously
220 reported (Czop et al. 2011; Lehoux et al. 2013) over the same pH range. This may be due to
221 the buffering capacity of the organic substrate used in our study. The most effective soil for
222 buffering residue additions was attributed to those with high organic carbon content (Lehoux
223 et al. 2013; Buckley et al. 2016). Reduction in solution pH is a critical factor for removing
224 trace elements from alkaline leachate water (Hua et al. 2015).

225 Adsorption of arsenate and vanadate to mineral surfaces occurs at circumneutral pH (Peacock
226 and Sherman 2004) and aluminate becomes highly insoluble below about pH 10.5 and
227 precipitates as an amorphous oxyhydroxide phase (Burke et al. 2012). Significant decreases
228 in aqueous Al, As and V concentrations with red mud additions were observed below
229 approximately pH 8.5 and resulted from an enhancement in both sorption (As and V) and
230 precipitation (Al) that effectively inhibited metal(loid) release to solution (Lehoux et al.
231 2013). Soluble V in the pH 11.1 treatment was a magnitude of 10 times lower than values
232 reported likely to cause genotoxic effects in plants (Mišík et al. 2014). Although leachate
233 additions in the current study raised substrate pH levels, all values were considerably lower
234 than pH 9. Mišík et al. (2014) suggested that pH control of soil to pH 9 in residue affected
235 soils would mitigate high aqueous concentrations phytotoxic effects.

236

237 Increased alkalinity, salinity and sodicity has led to inhibition of root growth in a range of
238 plants growing in bauxite residue aqueous extracts (Courtney and Mullen 2009) and residue
239 additions to soil resulted in decreased plant growth and increased trace element uptake
240 (Ruyters et al. 2011). Decreases in plant growth in alkaline wetland substrates was
241 previously attributed to reduced N and P availability in the high pH substrates (Lawson,

242 2004; Mayes et al, 2009). Increased levels of soluble and exchangeable Na in the compost
243 substrate highlight concern for long-term soil quality and performance in constructed
244 wetlands. Soils with high Na content (sodicity) can have reduced permeability and this may
245 impede wetland performance.

246

247 Deficiency levels for macronutrients in *P. australis* of 1.45 % for N, 0.06 % for P and 0.7 –
248 0.75 % for K were reported by Allen and Pearsall (1963). Although leachate additions
249 resulted in a trend of decreased N content, this was not significantly different and indeed, N
250 values of >1.5 % were recorded for the second highest leachate application rate. No
251 treatment displayed nutrient deficiencies in the aerial portions for P and K. Although
252 significant differences were recorded for Ca, Mg and S content, the Ca and Mg content for
253 above ground samples were within the normal range reported by Vymazal and Šveha (2012)
254 for natural wetlands.

255 Concentrations of Ca for adequate growth in plants are normally around 0.5% shoot dry
256 matter (Batty and Younger 2004). For leachate applications of 1:100 this value was not
257 reached. The lower concentrations in the higher pH loading treatment occurred despite
258 exchangeable reserves for all treatments being adequate although soluble Ca was
259 significantly decreased. The presence of elevated concentrations of Na in soluble and
260 exchangeable form may inhibit uptake of Ca into root tissues. Mean ranges for Na in
261 belowground tissues in natural wetlands are 0.14 – 0.27 % and 0.27 – 0.75 % for constructed
262 wetlands (Vymazal and Šveha 2012). Even with the higher application rates the Na content
263 of belowground samples is at the lower end of the constructed wetlands values. Transfer of
264 Na into aerial portions as a result of higher leachate loadings was not evident.

265 Out of 6 non-nutrient elements detected in plant samples, concentrations in belowground
266 biomass were several magnitudes higher than in aboveground for Al, As, and V. A trend of
267 higher content in belowground compared to aboveground was also observed for Cd, Cr and
268 Ni. Restriction of translocation to aboveground biomass is believed to be a strategy of metal
269 tolerance and plants can avoid the potential effects of high metal concentrations on the
270 photosynthetic tissue (Bragato et al. 2006).

271 Bonanno (2011) has proposed *P. australis* as a potential bioindicator of Al trace content in
272 wetlands. In the current study results showed higher concentrations in belowground samples

273 with low mobility through to the aerial portions. This phenomenon is well reported as Al is
274 immobilized in the roots of metallophytes (e.g. Baker et al. 2000; Vymazal et al. 2009;
275 Bonanno 2011). Vegetation content of Al has also been shown to be affected by
276 concentrations in water and sediment with lower concentrations for both sediment and plant
277 content reported with distance from influent (Lesage et al. 2007). While additions of residue
278 leachate increased levels of Al in belowground biomass, all concentrations were below those
279 reported for other constructed wetlands treating domestic waste water (Lesage et al. 2007;
280 Vymazal et al. 2009) and much lower than the 1000-3000 mg kg⁻¹ phytotoxic levels cited by
281 Kabata-Pendias and Pendias (2001).

282 Cd content in aboveground biomass recorded was significantly increased with leachate
283 applications but values detected were much lower than values reported for constructed
284 wetlands (Lesage et al. 2007; Vymazal et al. 2007), river areas (Bonanno and Giudice 2010)
285 and are well below the phytotoxic range (5–700 mg kg⁻¹) reported by Chaney (1989).

286 Leachate additions had no effect on belowground Cr levels and values are considerably lower
287 than the 15-22 mg kg⁻¹ reported by Lesage et al (2007) for *P. australis* in a constructed
288 wetland. Leachate additions resulted in increased levels of Cr in aboveground biomass and
289 fall within the same range reported by Lesage et al. (2007) and are at the lower end of the
290 range reported by Vymazal et al. (2007) for a range of constructed wetlands and natural
291 stands. The toxicity range of Cr levels is 5–30 mg kg⁻¹ (Kabata-Pendias and Pendias 2001)
292 and all values are well below the threshold.

293 Bonanno (2011) reported that roots in *P. australis* may have inherent protective mechanisms
294 to prevent V from penetrating into other organs. Such exclusion mechanisms have been
295 demonstrated in several plant species (e.g. Baker 2000) and Qian et al. (2014) have suggested
296 the limited translocation of V to plant aerial parts provides a selective advantage during
297 colonization and establishment. Soluble forms of V in sediment appear to be readily taken up
298 by roots and soluble vanadium was below limits of detection in all treatment except treatment
299 6. Although this treatment yielded the highest V content in belowground samples (ca 5.4 mg
300 kg⁻¹) this is less than the values reported by Bonanno (2011) for the riverside *P. australis* in
301 an urbanised area and is considerably lower than values recorded for brownfield root content
302 in *P. australis* (200 µg g⁻¹). Similarly, the low V content in aboveground biomass (< 0.26 µg
303 g⁻¹) are several magnitudes lower than those recorded by Qian et al. (2014).

304 Whilst Ni levels above 5 mg kg⁻¹ are considered toxic (Allen 1989) by the end of the growing
305 period, leaves can accumulate up to 60 mg kg⁻¹ (Bragato et al. 2006). Concentrations in the
306 current study are at the lower end of the range reported for *P. australis* growing in wetlands
307 range from 0.5 – 9 mg kg⁻¹ (Bonanno and Giudice 2010), 0.6 – 11 mg kg⁻¹ (Vymazal et al.
308 2007).

309 As content in the current study are similar to those reported by Vymazal et al. (2009) for
310 constructed wetlands but are considerably lower than reported by Allende et al. (2014) which
311 examined acidic wastewater. Results are in agreement with Vymazal et al. (2009) and
312 Allende et al. (2014) that roots of *P. australis* have greater As concentration than shoots.
313 Allende et al. (2014) attributed elevated As in wetland plants to the immature state of the
314 wetland system.

315 Buckley et al. (2015) demonstrated the ability of organic soils to buffer high pH of NaOH
316 solutions. Similarly, buffering of high pH residue leachate allows for such soils to support
317 plant growth and avoid excessive uptake of trace elements. Trace element levels in *P.*
318 *australis* detected in the current study are lower than levels reported by Ruyters et al. (2011)
319 for plants growing in red mud affected soils. Ruyters et al. (2011) reported that although
320 these trace elements were detected they did not exceed toxic limit and hypothesized Na as the
321 prime cause which affected plant growth. Although not investigated in the current study,
322 chlorophyll content (indicator of stress) has been found to have correlations with total N
323 content (Lippert et al. 2001). N content did not significantly vary between treatments or
324 display any negative correlations with trace element uptake.

325

326 Factors influencing plant uptake in wetland systems include age of wetland system and
327 period of acclimatization, duration of exposure, the type of vegetation and the type of
328 substrate. Accumulation of Na and metal(loids) in wetland substrates may be problematic in
329 their long-term application for treating hyperalkaline residue leachates. Analysis of stream
330 sediment samples following the red mud spill at Ajka, Hungary demonstrated that the bulk of
331 trace elements were in forms that were not readily bioavailable (Mayes et al 2011). Further
332 work is recommended to investigate trace element accumulation and potential bioavailability
333 in constructed wetlands treating bauxite residue leachate.

334

335

336

337 **Conclusions**

338 Growth of *Phragmites australis* (below and above ground growth and biomass) was not
339 adversely affected in bauxite residue leachate treatments (pH 8.5-11.1). Whilst some
340 substrate pH, EC and Na content were increased these were not to levels of concern. A
341 pattern of increased trace element content in vegetation was found with increased leachate
342 rates but no treatments were at levels of concern.

343

344 Previously, batch trials have shown the potential for wetland mechanisms to decrease pH and
345 trace element concentration in bauxite residue leachate and NaOH solutions. Plant growth
346 and biological activity are integral components of constructed wetlands treating mine waters.
347 The ability of *Phragmites australis* to grow in the residue leachate treatments provides strong
348 encouragement for constructed wetland approaches to effectively treat alkaline bauxite
349 residue leachate.

350

351 **Acknowledgements**

352 This research was supported by funding from the International Aluminium Institute.

353

355 **References**

- 356 Allen SE, Pearsall W (1963) Leaf analysis and shoot production in *Phragmites*. *Oikos*
357 14:176-189.
- 358 Allen SE (1989) *Chemical Analysis of Ecological Material*, 2nd edition. Blackwell Scientific
359 Publications, Oxford, 368 pp.
- 360 Allende KL, McCarthy DT, Fletcher TD (2014) The influence of media type on removal of
361 arsenic, iron and boron from acidic wastewater in horizontal flow wetland microcosms
362 planted with *Phragmites australis*. *Chem Eng J* 246:217-228
- 363 Baker AJ (2000) Metal hyperaccumulator plants: a review of the biological resource for
364 possible exploitation in the phytoremediation of metal-polluted soils. In: Terry N, Baneulos
365 GS (eds) *Phytoremediation of contaminated soil and water*. CRC Press, Boca Raton, FL, pp
366 85–107
- 367 Banks MK, Schwab AP, Alleman JE, Hunter JG, Hickey JC (2006) Constructed wetlands for
368 the remediation of blast furnace slag leachates. Indiana: Joint Transportation Research
369 Program.
- 370 Batty LC, Younger PL (2004) Growth of *Phragmites australis* (Cav.) Trin. ex. Steudel in mine
371 water treatment wetlands: effects of metal and nutrient uptake. *Environ Pollut* 132:85-93
- 372 Bonanno G (2011) Trace element accumulation and distribution in the organs of *Phragmites*
373 *australis* (common reed) and biomonitoring applications. *Ecotox Environ Safe* 74:1057-1064
- 374 Bonanno G, Giudice RL (2010) Heavy metal bioaccumulation by the organs of *Phragmites*
375 *australis* (common reed) and their potential use as contamination indicators. *Ecol Indic*
376 10:639-645.
- 377 Bragato C, Brix H, Malagoli M (2006) Accumulation of nutrients and heavy metals in
378 *Phragmites australis* (Cav.) Trin. ex Steudel and *Bolboschoenus maritimus* (L.) Palla in a
379 constructed wetland of the Venice lagoon watershed. *Environ Pollut* 144:967-975.
- 380 Buckley R, Curtin T, Courtney R (2016) The potential for constructed wetlands to treat
381 alkaline bauxite residue leachate: laboratory investigations. *Environ Sci Pollut Res* 23:14115–
382 14122
- 383 Burke IT, Mayes WM, Peacock CL, Brown AP, Jarvis AP, Gruiz K (2012) Speciation of
384 arsenic, chromium, and vanadium in red mud samples from the Ajka spill site,
385 Hungary. *Environ Sci Technol* 46:3085-3092.
- 386 Burke IT, Peacock CL, Lockwood CL, Stewart DI, Mortimer RJ, Ward MB, Renforth P,
387 Gruiz K, Mayes WM (2013) Behavior of aluminum, arsenic, and vanadium during the

388 neutralization of red mud leachate by HCl, gypsum, or seawater. *Environ Sci Technol*
389 47:6527-6535.

390 Chaney RL (1989) Toxic element accumulation in soils and crops: protecting soil fertility and
391 agricultural food chains. In: Bar-Yosef, B., Barrow, N.J., Goldshmid, J. (Eds.), *Inorganic*
392 *Contaminants in the Vadose Zone*. Springer-Verlag, Berlin, pp. 140–158.

393 Courtney R, Mullen G (2009) Use of germination and seedling performance bioassays for
394 assessing revegetation strategies on bauxite residue. *Water Air Soil Pollut* 197:15-22

395 Czop M, Motyka J, Sracek O, Szuwarzyński M (2011) Geochemistry of the hyperalkaline
396 Gorka Pit Lake (pH> 13) in the Chrzanow region, southern Poland. *Water Air Soil Pollut*
397 214:423-434

398 Hua T, Haynes RJ, Zhou YF, Boullemant A, Chandrawana I (2015) Potential for use of
399 industrial waste materials as filter media for removal of Al, Mo, As, V and Ga from alkaline
400 drainage in constructed wetlands—adsorption studies. *Water Res* 71:32-41

401 Jenkinson DS, Powelson DS (1976) The effects of biocidal treatments on metabolism in
402 soil—I. Fumigation with chloroform. *Soil Biol Biochem* 8:167–177

403 Joergensen R, Wichern F (2008) Quantitative assessment of the fungal contribution to
404 microbial tissue in soil. *Soil Biol Biochem* 40:2977 – 2991

405 Kabata-Pendias A, Pendias H (2001) *Trace Element in Soils and Plants*, 3rd edn. CRC Press,
406 Boca Raton

407 Klauber C, Gräfe M, Power G (2011) Bauxite residue issues: II. options for residue
408 utilization. *Hydrometallurgy*, 108:11-32

409 Lawson CJ (2004) A Preliminary Analysis into the use of Passive Remediation in Calcareous
410 Alkaline Waters, Unpublished MSc. thesis. University of Newcastle upon Tyne

411 Lehoux AP, Lockwood CL, Mayes WM, Stewart DI, Mortimer RJ, Gruiz K, Burke IT (2013)
412 Gypsum addition to soils contaminated by red mud: implications for aluminium, arsenic,
413 molybdenum and vanadium solubility. *Environ Geochem Health* 35:643-656

414 Lesage E, Rousseau DPL, Meers E, Tack FMG, De Pauw N (2007) Accumulation of metals
415 in a horizontal subsurface flow constructed wetland treating domestic wastewater in Flanders,
416 Belgium. *Sci Total Environ* 380:102-115

417 Lippert I, Rolletschek H, Kohl JG (2001) Photosynthetic pigments and efficiencies of two
418 *Phragmites australis* stands in different nitrogen availabilities. *Aquat Bot* 69:359-365

419 Matthews DJ, Moran BM, Otte ML (2005) Screening the wetland plant species *Alisma*
420 *plantago-aquatica*, *Carex rostrata* and *Phalaris arundinacea* for innate tolerance to zinc and
421 comparison with *Eriophorum angustifolium* and *Festuca rubra* Merlin. *Environ Pollut* 134:43-
422 351

423 Mayes WM, Younger PL, Aumônier J (2008) Hydrogeochemistry of alkaline steel slag
424 leachates in the UK. *Water Air Soil Pollut* 195:35-50

425 Mayes WM, Batty LC, Younger PL, Jarvis AP, Kõiv M, Vohla C, Mander U (2009) Wetland
426 treatment at extremes of pH: A review. *Sci Total Environ* 407:3944-3957

427 Mayes WM, Jarvis AP, Burke IT, Walton M, Feigl V, Klebercz O, Gruiz K (2011) Dispersal
428 and attenuation of trace contaminants downstream of the Ajka bauxite residue (red mud)
429 depository failure, Hungary. *Environ Sci Technol* 45:5147-5155

430 Mišík M, Burke IT, Reismüller M, Pichler C, Rainer B, Mišíková K, Mayes WM,
431 Knasmueller S (2014) Red mud a byproduct of aluminum production contains soluble
432 vanadium that causes genotoxic and cytotoxic effects in higher plants. *Sci Total*
433 *Environ* 493:883-890

434 Peacock CL, Sherman DM (2004) Vanadium (V) adsorption onto goethite (α -FeOOH) at pH
435 1.5 to 12: a surface complexation model based on ab initio molecular geometries and EXAFS
436 spectroscopy. *Geochim Cosmochim Acta* 68:1723-1733

437 Qian Y, Gallagher FJ, Feng H, Wu M, Zhu Q (2014) Vanadium uptake and translocation in
438 dominant plant species on an urban coastal brownfield site. *Sci Total Environ* 476:696-704

439 Ruyters S, Mertens J, Vassilieva E, Dehandschutter B, Poffijn A, Smolders E (2011) The red
440 mud accident in Ajka (Hungary): plant toxicity and trace metal bioavailability in red mud
441 contaminated soil. *Environ Sci Technol* 45:1616-1622

442 Truu M, Juhanson J, Truu J (2009) Microbial biomass, activity and community composition
443 in constructed wetlands. *Sci Total Environ* 407:3958-3971

444 Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil
445 microbial biomass C. *Soil Biol Biochem* 19:703-707

446 Vymazal J (2005) Horizontal sub-surface flow and hybrid constructed wetlands systems for
447 wastewater treatment. *Ecol Eng* 25:478-490

448 Vymazal J, Švehla J, Kröpfelová L, Chrastný V (2007) Trace metals in *Phragmites australis*
449 and *Phalaris arundinacea* growing in constructed and natural wetlands. *Sci Total Environ*
450 380:154-162

451 Vymazal J, Kröpfelová L, Švehla J, Chrastný V, Štíhová J (2009) Trace elements in
452 *Phragmites australis* growing in constructed wetlands for treatment of municipal wastewater.
453 *Ecol Eng* 35:303-309

454 Vymazal J, Šveha J (2012) Removal of alkali metals and their sequestration in plants in
455 constructed wetlands treating municipal sewage. *Hydrobiologia* 692:131-143

456

457 Wang X, Zhang Y, Lv F, An Q, Lu R, Hu P, Jiang S (2015) Removal of alkali in the red mud
458 by SO₂ and simulated flue gas under mild conditions. *Environ Prog Sustain Energy* 34:81-87

459

460

461

462

463

464

465 **Figure captions**

466

467 **Figure 1.** Substrate pH, EC and soluble elements in residue leachate treatments. Means
468 followed by the same letter are not significantly different at $P \leq 0.05$

469

470 **Figure 2.** Nutrient and Na content in *Phragmites australis* belowground biomass. Means
471 followed by the same letter are not significantly different at $P \leq 0.05$

472

473 **Figure 3.** Nutrient content with Na in *Phragmites australis* aboveground biomass. Means
474 followed by the same letter are not significantly different at $P \leq 0.05$

475

476 **Figure 4.** Trace element content in *Phragmites australis* belowground biomass. Means
477 followed by the same letter are not significantly different at $P \leq 0.05$

478

479 **Figure 5.** Trace element content in *Phragmites australis* aboveground biomass. Means
480 followed by the same letter are not significantly different at $P \leq 0.05$

481

482

483

484

485

486

487


488

489

490

491

492



494


495

496

497

498


499



501

502


503




505

506

507





511
512

513

514

515

516

517

518

519

The potential for constructed wetlands to treat alkaline bauxite-residue leachate: phragmites australis growth

Higgins, D.

2016-09-21

Attribution-NonCommercial 4.0 International

Higgins D, Curtin T, Pawlett M, Courtney R. (2016) The potential for constructed wetlands to treat alkaline bauxite-residue leachate: phragmites australis growth. *Environmental Science and Pollution Research*, Volume 23, Volume 23, December 2016, pp. 24305-24315

<https://doi.org/10.1007/s11356-016-7702-1>

Downloaded from CERES Research Repository, Cranfield University