

Journal Pre-proof

Aspergillus ullungdoensis sp. nov., *Penicillium jeongsukae* sp. nov., and other fungi from Korea

Hyang Burm Lee, Thuong T.T. Nguyen, So Jeong Noh, Dong Hee Kim, Ki Hyun Kang, Su Jin Kim, Paul M. Kirk, Simon V. Avery, Angel Medina, John E. Hallsworth



PII: S1878-6146(24)00079-5

DOI: <https://doi.org/10.1016/j.funbio.2024.05.014>

Reference: FUNBIO 1475

To appear in: *Fungal Biology*

Received Date: 14 August 2023

Revised Date: 1 April 2024

Accepted Date: 14 May 2024

Please cite this article as: Lee, H.B., Nguyen, T.T.T., Noh, S.J., Kim, D.H., Kang, K.H., Kim, S.J., Kirk, P.M., Avery, S.V., Medina, A., Hallsworth, J.E., *Aspergillus ullungdoensis* sp. nov., *Penicillium jeongsukae* sp. nov., and other fungi from Korea, *Fungal Biology*, <https://doi.org/10.1016/j.funbio.2024.05.014>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier Ltd on behalf of British Mycological Society.

1 ***Aspergillus ullungdoensis* sp. nov., *Penicillium jeongsukae* sp. nov., and other fungi from**
2 **Korea**

3

4 Hyang Burm Lee ^{a,*}, Thuong T.T. Nguyen ^a, So Jeong Noh ^a, Dong Hee Kim ^a, Ki Hyun Kang ^a,
5 Su Jin Kim ^a, Paul M. Kirk ^b, Simon V. Avery ^c, Angel Medina ^d, John E. Hallsworth ^e

6

7 ^a *Environmental Microbiology Lab, Department of Agricultural Biological Chemistry, College of*
8 *Agriculture and Life Sciences, Chonnam National University, Gwangju 61186, Korea*

9 ^b *Biodiversity Informatics and Spatial Analysis, Jodrell Laboratory, Royal Botanic Gardens Kew,*
10 *Surrey TW9 3DS, UK*

11 ^c *School of Life Sciences, University of Nottingham, University Park, Nottingham, NG7 2RD, UK*

12 ^d *Applied Mycology, Cranfield University, Cranfield, MK43 0AL, UK*

13 ^e *Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast,*
14 *19 Chlorine Gardens, Belfast, BT9 5DL, UK*

15

16 * Corresponding author. Environmental Microbiology Lab, Department of Agricultural
17 Biological Chemistry, College of Agriculture and Life Sciences, Chonnam National University,
18 Gwangju 61186, Korea

19 *E-mail address:* hblee@jnu.ac.kr (H.B. Lee).

20

21 **ABSTRACT**

22 *Eurotiales* fungi are thought to be distributed worldwide but there is a paucity of information
23 about their occurrence on diverse substrates or hosts and at specific localities. Some of the
24 *Eurotiales*, including *Aspergillus* and *Penicillium* species, produce an array of secondary
25 metabolites of use for agricultural, medicinal, and pharmaceutical applications. Here, we carried
26 out a survey of the *Eurotiales* in Korea, focusing on soil, freshwater, and plants (dried persimmon
27 fruits and seeds of *Perilla frutescens*, known commonly as shiso). We obtained 11 species that—
28 based on morphology, physiology, and multi-locus (ITS, *BenA*, *CaM*, and *RPB2*) phylogenetic
29 analyses—include two new species, *Aspergillus ullungdoensis* sp. nov. and *Penicillium*
30 *jeongsukae* sp. nov., and nine species that were known, but previously not described in Korea,
31 *Aspergillus aculeatinus*, *Aspergillus aurantiacoflavus*, *Aspergillus croceiaffinis*, *Aspergillus*
32 *pseudoviridinutans*, *Aspergillus uvarum*, *Penicillium ferraniaense*, *Penicillium glaucoroseum*,
33 *Penicillium sajarovii*, and one, *Penicillium charlesii*, that was isolated from previously unknown
34 host, woodlouse (*Porcellio scaber*). We believe that biodiversity surveys and identifying new
35 species can contribute to set a baseline for future changes in the context of humanitarian crises
36 such as climate change.

37

38 **Keywords:** *Aspergillus ullungdoensis* sp. nov.; ascomycete phylogeny, biodiversity surveys,
39 *Eurotiales*, fungal ecology, *Penicillium jeongsukae* sp. nov.

40

41 **1. Introduction**

42

43 *Aspergillus* and *Penicillium* are fungal genera belonging to the ascomycete order *Eurotiales*. This
44 order includes five families:—*Aspergillaceae*, *Elaphomycetaceae*, *Penicillaginaceae*,
45 *Thermoascaceae*, and *Trichocomaceae*—that represent a total of 28 genera—*Acidotalaromyces*,

1 *Ascospirella*, *Aspergillago*, *Aspergillus*, *Dendrosphaera*, *Elaphomyces*, *Evansstolkia*, *Hamigera*,
2 *Leiothecium*, *Monascus*, *Paecilomyces*, *Penicillago*, *Penicilliopsis*, *Penicillium*, *Phialomyces*,
3 *Pseudohamigera*, *Pseudopenicillium*, *Pseudotulostoma*, *Rasamsonia*, *Sagenomella*, *Sclerocleista*,
4 *Talaromyces*, *Thermoascus*, *Thermomyces*, *Trichocoma*, *Warcupiella*, *Xerochrysum*, and
5 *Xeromyces* (Houbraken et al., 2020). These fungi are commonly found in air/bioaerosols, soil,
6 plants and animals, freshwater and seawater, the subsurface, and the built environment and many
7 can grow at low water activities and low temperatures (Nguyen et al., 2021a; Nguyen et al., 2020;
8 Houbraken et al., 2016; Hubka et al., 2014; Samson et al., 2014, 2010; Frisvad and Samson, 2004;
9 Lee and Magan, 1999; Micheluz et al., 2021). Some of these fungi are economically important in
10 the fields of biotechnology, food, and medicine (Hyde et al., 2019; Houbraken et al., 2014; Frisvad
11 and Samson, 2004) and can make important contributions towards Sustainability Development
12 Goals (Purchase, 2016; Noel et al., 2023).

13 The genus *Aspergillus* (family *Aspergillaceae*, order *Eurotiales*) comprises a large group
14 of filamentous fungi subdivided into six subgenera, 27 sect. and 75 ser. by a phylogenetic
15 approach, often supported by morphological, physiological, and/or extrolite profile data
16 (Houbraken et al., 2020). Members of this genus are mainly saprophytes that can decompose
17 organic materials that can be found in plants, fruits, foods, indoor environments, water, soil, air,
18 and rock as well as on various other surfaces (Pangging et al., 2022; Sklenář et al., 2017; Chen et
19 al., 2017; Hyde et al., 2016; Hubka et al., 2014; Samson et al., 2014; Perrone et al. 2008;
20 Hallsworth, 2019). Species of *Aspergillus* can have positive or negative impacts on human
21 activities. Indeed, some species of *Aspergillus* have economic importance in industry, agriculture
22 and medicine (Meyer et al., 2011). For example, *A. terreus* is known for producing itaconic acid
23 used in polymer manufacture, the cholesterol-lowering drug lovastatin (Wierckx et al., 2020;
24 Boruta and Bizukojc, 2017), and the phytohormone gibberellin (produced by *A. fumigatus*) which
25 improves plant growth (Khan et al., 2011). Furthermore, penicillixanthone A produced by *A.*
26 *fumigatus* is reported to have effective anti-HIV-1 activity (Tan et al., 2019). *Aspergillus niger* is
27 known to produce citric acid which is widely used in beverages, foods, detergents, cosmetics and
28 pharmaceutical industries (Behera, 2020).

29 On the contrary, some species can produce mycotoxins, cause food spoilage, and
30 sometimes cause infections in humans and animals (Kim et al., 2022; Ráduly et al., 2020; Frisvad
31 et al., 2019; Mousavi et al., 2016; Magan, 2006; Paulussen et al., 2017). *Aspergillus ochraceus*
32 belonging to sect. *Circumdati* produces the well-known the mycotoxin ochratoxin A (Visagie et
33 al., 2014; Magan and Aldred, 2005; van der Merwe et al., 1965). Aflatoxins produced by *A. flavus*
34 belonging to sect. *Flavi* are major toxic, carcinogenic secondary metabolites (Frisvad et al., 2019).

35 Currently, the *Aspergillus* genus consists of 480 species (Index Fungorum 2023; Sung et
36 al., 2022; Wang and Zhuang, 2022; Houbraken et al., 2020), only two of which are registered in
37 Korea according to the Index Fungorum database (<http://www.indexfungorum.org>). These are
38 *Aspergillus cibarius* S.B. Hong & Samson and *Aspergillus koreanus* Hyang B. Lee, T.T. Duong
39 & T.T. Nguyen.

40 The genus *Penicillium* (family *Aspergillaceae*, order *Eurotiales*), first described by Link
41 in 1809, is one of the most common fungal genera. Species of *Penicillium* play an important role
42 in diverse natural environments by degrading and cycling the organic matter in microbial
43 necromass. They also have medicinal and industrial uses. For example, *Penicillium* species have
44 been known to produce hydrolytic enzymes (Hamlyn et al., 1987). Important medical compounds,
45 such as penicillin, are produced by *P. chrysogenum* (now reclassified as *P. rubens*) (Houbraken

1 et al., 2011). *Penicillium compactum* and *P. coprobium* produce pyripyropens that act to inhibit
2 acyl-CoA:cholesterol acyltransferase (ACAT) (Frisvad et al., 2004). However, some species
3 cause food spoilage, produce various mycotoxins, and also cause diseases in animals (including
4 humans).

5 The genus *Penicillium* is made up of two subgenera, 32 sect., 89 ser. (Houbraken et al.
6 2020), with a total of more than 500 species (Index Fungorum 2023; Visagie and Yilmaz, 2022;
7 Nguyen et al., 2021a; Houbraken et al., 2020). So far only these 10 species are registered in Korea
8 according to the Index Fungorum database. These are *Penicillium acidum* Hyang B. Lee, T.T.
9 Duong & T.T.T. Nguyen, *Penicillium aquadulcis* Hyang B. Lee & T.T.T. Nguyen, *Penicillium*
10 *aquaticum* Hyang B. Lee, T.T. Duong & T.T.T. Nguyen, *Penicillium circulare* Hyang B. Lee,
11 P.M. Kirk & T.T.T. Nguyen, *Penicillium daejeonium* S.H. Yu & H.K. Sang, *Penicillium*
12 *geumsanense* Hyang B. Lee, P.M. Kirk & T.T.T. Nguyen, *Penicillium koreense* S.B. Hong, D.H.
13 Kim & Y.H. You, *Penicillium mali-pumilae* Hyang B. Lee, T.T.T. Nguyen & Houbraken,
14 *Penicillium psychrotrophicum* Hyang B. Lee, H.Y. Mun, J.C. Frisvad & Houbraken, and
15 *Penicillium ulleungdoense* D.H. Choi & J.G. Kim.

16 Whereas some known *Aspergillus* and *Penicillium* species originate in Korea, there has
17 been little systematic work to understand the diversity of these fungal taxa across environmental
18 niches of this geographical region. Yet Korea has a peculiar climate with dry cold winters and
19 humid hot summers, is coastal and subject to the North Pacific anticyclone, and has extensive
20 natural environments but also much land disturbed by anthropogenic activity, all of which
21 collectively creates a unique combination of niches for fungi. Here, we carried out a survey of
22 *Aspergillus* and *Penicillium* species in different environmental settings in Korea. We discovered
23 11 species, including two new species, proposed as *Aspergillus ullungdoensis* sp. nov. and
24 *Penicillium jeongsukae* sp. nov.

26 **2. Materials and methods**

28 *2.1. Sampling and isolation*

30 Soil samples, freshwater, dried persimmon fruits, and seeds of the green shiso plant (*Perilla*
31 *frutescens*) were collected from different areas in Korea (Fig. 1). The soil was collected from
32 Cheongyang, located in Chungnam Province, and from Ulleung Island, North Gyeongsang
33 Province, South Korea. The freshwater sample was taken from a depth of 0–10 cm in lake located
34 at Chonnam National University, Gwangju and from waterfall freshwater located in Jeju Island,
35 Jeju Province, South Korea. Dried persimmon fruits purchased from a grocery store located at
36 Cheongyang, Chungnam Province, and then stored at the room temperature. Finally, the seeds of
37 *Per. frutescens* were obtained from plants growing on agricultural land in Jeongeup, Jeollabuk-
38 do, South Korea.

39 Samples were individually stored in paper bags or sterile 50-mL conical tubes and brought back
40 to the laboratory for fungal isolations. For the soil samples, 1 g of soil was added to 9 mL of
41 sterile distilled water. Then, serial dilutions of the mixture (from 10 parts water to 1 part of sample
42 mixture to 10 parts of water to 5 parts of sample mixture, by volume) were prepared. A 100- μ L
43 aliquot of the soil solution was dispensed onto potato dextrose agar (PDA; Becton, Dickinson and
44 Co., Sparks, MD, USA) containing the antibiotic neomycin (50 mg/L). For freshwater samples, a
45 100- μ L aliquot of water was dispensed onto PDA containing neomycin (50 mg/L). For dried

1 persimmon samples, spores were carefully examined under a stereomicroscope and transferred to
2 PDA plates. The Petri plates were incubated at 25 °C for 3–7 days. To isolate fungi from seed
3 samples, 7–10 seeds were directly plated on a moist blotter and incubated at 25 °C in the dark for
4 7–14 days. Fungal spores on the seeds were transferred to PDA (Difco™, Sparks, MD, USA)
5 using a capillary tube under a stereomicroscope.

6 Pure isolates were obtained and stored on PDA slants at 4 °C and in 20% (v/v) glycerol at
7 –80 °C at the Environmental Microbiology Laboratory Fungarium (Chonnam National University,
8 Gwangju, Korea). The isolates were also deposited at the National Institute of Biological
9 Resources (NIBR, Incheon, Korea) under the accession numbers NIBRFG0000510049,
10 NIBRFGC000508992, NIBRFGC000508615, NIBRFG0000509597, and NIBRFGC000510099
11 at the Nakdonggang National Institute of Biological Resources (NNIBR, Sangju, Korea) under
12 the accession number NNIBRFG31720 and NNIBRFG46695.

14 2.2. Fungal morphology studies

16 Fungal isolates were three-point inoculated onto Czapek yeast autolysate agar (CYA), malt
17 extract agar (MEA), yeast extract sucrose agar (YES), oatmeal agar (OA), dichloran 18% (w/v)
18 glycerol agar (DG18), CYA agar supplemented with 5% (w/v) NaCl (CYAS), and creatine
19 sucrose agar (CREA). For each medium type and each isolate, Petri plates were inoculated at 20,
20 25, 30, and 37 °C for 7 days. Medium preparation, inoculation, and incubation were performed
21 according to Samson et al. (2014). Colony characteristics such as colour, margin, diameter and
22 texture were recorded after 7 days of incubation. Slides were prepared using 60% (v/v) lactic acid,
23 and 96% (v/v) ethanol was used to remove excess conidia. An Olympus BX53 light microscope
24 with differential interference contrast optics (Olympus, Tokyo, Japan) was used to obtain digital
25 images of conidiophores, vesicles, metulae, phialides, and conidia.

27 2.3. Extraction, PCR amplification, and sequencing of DNA

29 Total genomic DNA was extracted from fungal mycelia using the Solg™ Genomic DNA
30 Preparation Kit (Solgent Co. Ltd., Daejeon, Korea) following the manufacturer's instructions.
31 Marker genes were selected as follows: the internal transcribed spacer (ITS) region of the rRNA,
32 β -tubulin (*benA*), calmodulin (*CaM*), and RNA polymerase II second largest subunit (*RPB2*).
33 Amplification of the ITS region, *BenA*, *CaM* and *RPB2* was performed using the following
34 conditions: initial denaturation 95 °C for 5 min, 30 cycles of denaturation (95 °C for 30 s),
35 annealing [temperature according to the primer for 45 s, (Table S1)], and extension (72 °C for 45
36 s), final extension 72 °C for 7 min. The PCR products were purified using an Accuprep PCR
37 Purification Kit (Bioneer Corp.). Sequencing was performed using the same primers on an ABI
38 PRISM 3730XL Genetic Analyzer (Applied Biosystems, California, USA).

40 2.4. Molecular analyses

42 The raw forward and reverse reads were edited for ambiguous bases at both ends and assembled
43 into contigs using SeqMan Version 7.1.0. The edited sequences were BLAST-searched against
44 the NCBI GenBank nucleotide database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to identify their
45 closest relatives. Sequences of previously published species of *Aspergillus* and *Penicillium* were

1 obtained from the GenBank sequence database (<http://www.ncbi.nlm.nih.gov/>). The sequences
2 were aligned using MAFFT (<http://mafft.cbrc.jp/alignment/server>) (Katoh et al. 2019), then
3 confirmed manually in MEGA v. 7 (Kumar et al. 2016). Data were converted from a FASTA
4 format to NEXUS and PHYLIP formats using the online tool Alignment Transformation
5 Environment (<https://sing.ei.uvigo.es/ALTER/>) (Glez-Peña et al. 2010). Phylogenetic
6 reconstructions based on the maximum likelihood (ML) method were performed using RAxML-
7 HPC2 on XSEDE via the CIPRES web portal (<https://www.phylo.org/portal2>), with a default
8 GTR substitution matrix and 1,000 rapid bootstraps. Bayesian inference (BI) analyses were
9 performed using MrBayes v. 3.2.6 (Ronquist et al. 2012). Four Markov chain Monte Carlo
10 (MCMC) chains were run from a random starting tree for 5 million generations, and trees were
11 sampled every 100th generation. The first 25% of the trees were removed as burn-in, and the
12 remaining trees were used to calculate posterior probabilities. Final consensus trees were viewed
13 using FigTree v. 1.3.1 (Rambaut 2009). Support values were provided at the branches (ML
14 bootstrap support (BS) and BI posterior probability (PP)). The newly obtained sequences were
15 deposited in the GenBank sequence database under the accession numbers provided in Table S2.
16

17 3. Results

18 3.1. Fungal phylogeny

19
20
21 The phylogenetic relationship of the new *Aspergillus* species with accepted species in ser.
22 *Spelaei* (sect. *Flavipedes*) was determined by analysing single and concatenated sequence datasets
23 of four loci (ITS, *BenA*, *CaM*, and *RPB2*). The concatenated alignment consisted of 3084
24 characters (including alignment gaps): 751 (ITS), 537 (*BenA*), 783 (*CaM*), and 1013 (*RPB2*).
25 Strains CNUFC UD06 and CNUFC UD07 clustered close to but separated from *A. lanuginosus*,
26 *A. albovidis* in the single (ITS, *BenA*, *CaM*, and *RPB2*) and combined phylogenies (Fig. 2 and
27 Figs. S1–S4). The combined ITS, *BenA*, *CaM*, and *RPB2* phylogeny data indicated that strains
28 CNUFC YB6 and CNUFC UDS2111, placed in ser. *Japonici* (sect. *Nigri*), were closely related
29 to the ex-type strains of *A. uvarum* CBS 121591 and *A. aculeatinus* CBS 121060, respectively
30 (Fig. S5). Strain CNUFC HSW10 grouped with ex-type strain of *A. aurantiacoflavus* CBS 141930
31 (Fig. S6) in ser. *Aspergillus* (sect. *Aspergillus*). Strain CNUFC 92316 clustered with the ex-type
32 strain of *A. croceiaffinis* CCF 6035 in ser. *Unguium* (sect. *Nidulantes*) (Fig. S7). CNUFC JPS72
33 clustered with the ex-type strain of *A. pseudoviridinutans* NRRL 62904 in ser. *Viridinutantes*
34 (sect. *Fumigati*) (Fig. S8).

35 The phylogenetic relationship of the new *Penicillium* species with accepted species in sect.
36 *Charlesia* was determined by analysing single and concatenated sequence datasets of four loci
37 (ITS, *BenA*, *CaM*, and *RPB2*). The concatenated alignment consisted of 2790 characters
38 (including alignment gaps): 732 (ITS), 490 (*BenA*), 550 (*CaM*), and 1018 (*RPB2*). Strains
39 CNUFC JSK815 and CNUFC JSK816 formed a well-supported independent branch (Fig. 3 and
40 Figs. S9–S12). Strain CNUFC CY22014 clustered with the ex-type strain of *P. charlesii* (Fig. 3
41 and Figs. S9–S12). Strains CNUFC JYS28, CNUFC CY22324, and CNUFC U734 grouped with
42 the ex-type strain of *P. ferraniaense* (sect. *Sclerotiorum*, ser. *Sclerotiorum*), *P. glaucoroseum*
43 (sect. *Lanata-Divaricata*, ser. *Janthinella*), and *P. sajarovii* (sect. *Ramosum*, ser. *Raistrickiorum*),
44 respectively (Figs. S13–15).
45

1 3.2. Taxonomy of the Korean isolates

2 ***Aspergillus aculeatinus*** Noonim, Frisvad, Varga, & Samson, Int. J. Syst. Evol. Microbiol. 58:
3 1733 (2008) (Fig. 4A-D)

4 *Colony diameter, 7 days at 25 °C (mm)*: CYA 31–36; MEA, 61–65; YES 31–33.

5 *Colony characteristics*: CYA 25 °C, 7 days: Colonies moderately deep, sulcate; margins
6 low, plane; mycelium white; texture velvety; sporulation absent to sparse; soluble pigments
7 absent; exudates absent; reverse pale yellow. MEA 25 °C, 7 days: Colonies low, plane; mycelia
8 white to light brown; texture floccose; sporulation moderately dense; soluble pigments absent;
9 exudates absent; reverse light brown. YES 25 °C, 7 days: Colonies moderately deep, sulcate;
10 margins low, plane; mycelium white; texture velvety; sporulation absent to sparse; soluble
11 pigments absent; exudates absent; reverse pale yellow.

12 *Micromorphology*: Conidiophores uniseriate, smooth-walled, 197–476 × 7–10.5 µm.
13 Vesicles globose, 31–48 µm. Phialides flask-shaped, covering the entire surface of the vesicle,
14 5.5–8 × 3–4 µm. Conidia subglobose to ellipsoidal, 3–4.5 × 2–3 µm.

15 *Material examined*: Republic of Korea, 2889-5 Saekdal-dong, Seogwipo-si, Jeju-do
16 (33°14'52.7"N 126°24'48.3"E), from soil, April 2021, H.B. Lee (culture CNUFC UDS2111).

17 *Notes*: The morphological characteristics of the isolate were consistent with those of *A.*
18 *aculeatinus*, as described by Noonim et al. (2008). However, CNUFC UDS2111 showed a slower
19 growth in colony diameter on CYA compared to *A. aculeatinus* (CYA >85 mm; MEA 47–50 mm;
20 YES >85 mm) (Noonim et al., 2008). *Aspergillus aculeatinus* was recorded from different
21 samples of Arabica coffee beans, indoor air, vineyard soil, bark of Chinese yew (*Taxus chinensis*
22 var. *mairei*), and sediments from the South China Sea in previous studies (Wu et al., 2021; Zhao
23 et al., 2021; Qiao et al., 2017; Noonim et al., 2008), and from soil in the present study.
24

25 ***Aspergillus aurantiocoflavus*** Hubka, A.J. Chen, Jurjević & Samson, Studies in Mycology 88: 82
26 (2017) (Fig. 4E-I)

27 *Colony diameter, 7 days at 25 °C (mm)*: CYA 8–10; MEA 16–20; YES 27–30

28 *Colony characteristics*: CYA 25 °C, 7 days: Colonies grew slowly, were pale yellow to
29 light brown, flocculent, irregular, no soluble pigments, reverse pale to dark orange. On MEA
30 25 °C, 7 days: Colonies grew with moderate speed, were white to pale yellow, dense sporulation,
31 no soluble pigments, reverse pale yellow. YES 25 °C, 7 days: Colonies grew slowly, texture was
32 floccose, white to bright yellow at the center, sporulation moderate, no soluble pigments, reverse
33 pale yellow.

34 *Micromorphology*: Conidiophores smooth stipes, hyaline, smooth-walled, 55–200 × 4–
35 11 µm; Vesicles globose to subglobose, 9.5–25 µm in diameter; Phialides flask-shaped, 5.8–10.8
36 × 2.8–5.3 µm; Conidia tuberculate, globose, subglobose to ellipsoidal, 5.5–8.1 × 4–7.6 µm;
37 Ascomata eurotium-like, cleistothecial, superficial, globose to subglobose, 60–78 µm.

38 *Material examined*: REPUBLIC OF KOREA, Jisan 2-dong, Dong-gu, Gwangju, Korea
39 (35.152111°N, 126.950056°E), from freshwater (culture CNUFC HSW10).

40 *Notes*: CNUFC HSW10 has similar morphological characteristics with the previously
41 described species of *A. aurantiocoflavus*. However, there were differences observed in the colony
42 diameter in comparison to previously described by Chen et al. (2017) (CYA 8–9 mm vs 2–3 mm;
43 MEA 16–20 mm vs 2–3 mm). *Aspergillus aurantiocoflavus* was isolated from a baby carrier
44 backpack, rubber toys, and cake in previous works (Chen et al., 2016), and from freshwater in the
45 present study.

1 *Aspergillus croceiaffinis* F. Sklenar, Jurjević, & Hubka, Mycologia 112(2): 359 (2020) (Fig. 4J-
2 M)

3 *Colony diameter, 7 days at 25 °C (mm)*: CYA 14–16; MEA, 10–13; YES 17–19.

4 *Colony characteristics*: CYA 25 °C, 7 days: Colonies moderately deep, plane; margins
5 entire to slightly irregular; texture floccose; sporulation moderately dense; soluble pigments very
6 weak; exudates absent; reverse brownish red. MEA 25 °C, 7 days: Colonies slightly raised at the
7 center; margins low, plane; texture velvety, floccose at the center; sporulation moderately dense;
8 soluble pigments absent; exudates absent; reverse deep orange. YES 25 °C, 7 days: Colonies
9 slightly raised at the center, sulcate; margins low, plane; texture floccose; sporulation moderately
10 dense; soluble pigments absent; exudates absent; reverse dark reddish brown.

11 *Micromorphology*: Conidiophores with smooth stipes, 96–147 × 2–3 µm. Vesicles
12 pyriform, 6.5–11 µm in diameter. Metulae cylindrical, 7–8 × 2–2.5 µm. Phialides flask-shaped,
13 7–8(–10) × 2–2.5 µm. Conidia globose to subglobose, 2.5–3.5 µm in diameter.

14 *Material examined*: Republic of Korea, Nari, Buk-myeon, Ulleung-gun, Gyeongsangbuk-
15 do (37°31'28.2"N 130°52'29.4"E), from soil (culture CNUFC 92316).

16 *Notes*: The morphological characteristics of the isolate were consistent with those of *A.*
17 *croceiaffinis*, as described by Sklenář et al. (2020). However, the conidiophore of the isolate
18 CNUFC 92316 was shorter than that of previously reported species [100–200 × 3–4 µm; Sklenář
19 et al. (2020)]. In addition, differences in colony diameter were observed on CYA (10–12 mm)
20 (Sklenář et al., 2020). *Aspergillus croceiaffinis* was isolated from a swab in a previous study
21 (Sklenář et al., 2020), and from soil sample in the present study.

22
23 *Aspergillus pseudoviridinutans* Sugui, S.W. Peterson & Kwon-Chung, in Sugui, Peterson, Figat,
24 Hansen, Samson, Mellado, Cuenca-Estrella & Kwon-Chung, *J. Clin. Microbiol.* 52(10): 3709
25 (2014) (Fig. 4N-Q)

26 *Colony diameter, 7 days at 25 °C (mm)*: CYA agar, 36–40; MEA, 66–70; YES agar, 38–
27 42.

28 *Colony characteristics*: CYA agar 25 °C, 7 days: Colonies centrally raised, surrounded
29 by sunken depression; margins low, plane; texture floccose; mycelia white; sporulation absent to
30 sparse; soluble pigments absent; exudates absent; reverse light yellow. MEA 25 °C, 7 days:
31 Colonies moderately deep; margins low; mycelium white; texture floccose; sporulation absent to
32 sparse; soluble pigments absent; exudates absent; reverse pale greenish yellow. YES agar 25 °C,
33 7 days: Colonies raised at the center, slightly sulcate; margins low; mycelia white; texture floccose;
34 sporulation sparse; soluble pigments absent; exudates absent; reverse vivid yellow.

35 *Micromorphology*: Conidiophores with smooth stipes, 30–83.5 × 3–6 µm. Vesicles
36 subglobose to flask-shaped, 7–14 µm in diameter. Phialides ampuliform, 3–5 × 1.5–2 µm. Conidia
37 in long chains, globose, 2–3.5 µm in diameter.

38 *Material examined*: Republic of Korea, Jeongeup (35°31'34.0"N 126°49'03.7"E),
39 Jeollabuk-do, from *Perilla frutescens* seed, October 2013 (culture CNUFC JPS72).

40 *Notes*: CNUFC JPS72 shared similar morphological characters with *A.*
41 *pseudoviridinutans* (Sugui et al., 2014). However, conidia are bigger than those of *A.*
42 *pseudoviridinutans* (1.75–2 µm) (Sugui et al., 2014). *Aspergillus pseudoviridinutans* was isolated
43 from the mediastinal lymph node of a 14-year-old boy with chronic granulomatous disease, soil,
44 a plant (*Pin. caribaea*), human lungs and eyes, and nostrils of parrots (*Amazona amazonica*) in

1 previous studies (Aneke et al., 2022; Langlois et al., 2021; Watanabe et al., 2021; Hubka et al.,
2 2018; Talbot and Barrs, 2018), and from *Per. frutescens* seed in the current study.

3
4 ***Aspergillus ullungdoensis*** Hyang B. Lee **sp. nov.**

5 *Index Fungorum*: IF 900871; Fig. 4R-V and Table 1

6 *Etymology*: Refers to the isolation location, Ulleungdo (Korea) from where the species
7 was first isolated.

8 *Type specimen*: Republic of Korea, Gitdaebong Peak, Ulleung Island, Korea (37°30'04"
9 N 130°51'23"E), from rhizosphere soil, 30 April 2019 (holotype CNUFC HT19207; ex-type
10 culture CNUFC UD06).

11 *Colony diam, 7 days (mm)*: CYA25°C 23–26 mm, CYA20°C 21–23; CYA30°C 34–37;
12 CYA37°C No growth; MEA25°C 27–29; YES25°C 32–35; OA25°C 23–27; CREA25°C 20–23.

13 *Colony characteristics*. CYA 25°C, 7 days: Colonies raised at center; texture velutinous;
14 mycelium white to beige; soluble pigments absent; exudate absent; sporulation moderate; reverse
15 pale yellow. YES 25°C, 7 days: Colonies radial sulcate; mycelium white; sporulation moderate;
16 exudate absent; soluble pigments absent; reverse moderate yellowish brown. MEA 25°C, 7 days:
17 Colonies low, plain; mycelium white; margins low; exudate absent; soluble pigments absent;
18 reverse pale yellow. OA 25°C, 7 days: Colonies nearly circular, slightly raised at center; texture
19 velutinous; sporulation strong; mycelia white then grayish pink; soluble pigments absent;
20 exudates absent; reverse moderate yellowish brown. CREA 25 °C, 7 days: Colonies slow growth,
21 acid not produced.

22 *Micromorphology*: Conidiophores biseriate. Stipes long, hyaline or light yellowish brown,
23 smooth, not septate, 150–1200 × 4–6 µm. Vesicles hyaline, globose, subglobose to pyriform, 9–
24 14 × 7.5–13 µm. Metulae 5.5–8(–9.5) × 3–3.5 µm. Phialides ampulliform, 5.5–9(–10) × 2.5–3.0
25 µm. Conidia globose, smooth, 2.5–3(–3.5) µm. Hülle cells present.

26 *Additional material examined*: Republic of Korea, Gitdaebong Peak, Ulleung Island,
27 (37°30'04" N 130°51'23"E), from rhizosphere soil, 30 April 2019 (culture CNUFC UD07).

28 *Notes*: In a NCBI BLASTn search based on ITS, *BenA*, *CaM* and *RPB2* sequences, the
29 closest match of *A. ullungdoensis* was 99.87% with *A. lanuginosus* NRRL 4610 (EF669604) (for
30 ITS), 98.16% with *A. alboviridis* FMR 15175 (LT798936) (for *BenA*), 97.67% with *A.*
31 *lanuginosus* NRRL 4610 (EF669562) (for *CaM*), and 99.21% with *A. movilensis* CCF 4410
32 (HG916718) (for *RPB2*).

33 A concatenated phylogenetic tree (ITS, *BenA*, *CaM*, and *RPB2*) revealed that *Aspergillus*
34 *ullungdoensis* forms distinct lineages sister to *A. lanuginosus* but with low statistical support.
35 However, *A. ullungdoensis* differs from *A. lanuginosus* by production Hülle cells. *Aspergillus*
36 *ullungdoensis* grows faster on MEA at 25 °C than *A. lanuginosus* on MEA at 25 °C (20–21 mm).
37 In additional, *A. lanuginosus* can grow at 37 °C, whereas *A. ullungdoensis* cannot grow at 37 °C.

38
39 ***Aspergillus uvarum*** G. Perrone, Varga, & Kozak, Int. J. Syst. Evol. Microbiol. 58(4): 1036 (2008)
40 (Fig. 4W-Z)

41 *Colony diameter, 7 days at 25 °C (mm)*: CYA 30–35; MEA, 30–48; YES 22–33.

42 *Colony characteristics*: CYA 25 °C, 7 days: Colonies raised at the center, sulcate;
43 margins low, plane; mycelium white to pale olive; texture velvety; sporulation moderately dense;
44 soluble pigments absent; exudates absent; reverse light yellow. MEA 25 °C, 7 days: Colonies low,
45 sulcate; margins low, plane; mycelia white; texture velvety; sporulation absent to sparse; soluble

1 pigments absent; exudates absent; reverse pale yellow. YES 25 °C, 7 days: Colonies raised at the
 2 center, strongly sulcate; mycelium white to brown; texture velvety; sporulation dense; soluble
 3 pigments absent; exudates absent; reverse moderate yellow.

4 *Micromorphology*: Conidiophores with smooth stipes, 314–1480 × 5–12 µm. Vesicles
 5 globose to subglobose or elliptical, 35–70 µm in diameter. Phialides ampulliform, 6–10 × 2.5–5
 6 µm. Conidia globose to subglobose, conspicuously spinose at maturity, 3.5–5 µm in diameter.

7 *Material examined*: Republic of Korea, Yongbong Reservoir, Yongbong-dong, Buk-gu,
 8 Gwangju (35°11'00.9"N 126°53'27.4"E), from freshwater, March 2022 (culture CNUFC YB6).

9 *Notes*: The morphological characteristics of the isolate CNUFC YB6 were similar to
 10 those of the previously described *A. uvarum* by Perrone et al. (2008). CNUFC YB6 showed slower
 11 growth on CYA and MEA in comparison to *A. uvarum* (Perrone et al., 2008) (CYA>90 mm;
 12 MEA>90 mm). *Aspergillus uvarum* was isolated from grapes, soil, food, indoor air, and a patient
 13 with onychomycosis in previous studies (Perrone et al., 2017; Qi et al., 2016; Zarei et al., 2015;
 14 Jurjević et al., 2012; Perrone et al., 2008), and from freshwater in the current study.

15
 16 ***Penicillium charlesii*** G. Sm., Trans. Brit. Mycol. Soc. 18 (1): 90 (1933)

17 *Descriptions & Illustrations*: Raper and Thom (1949)

18 *Material examined*: Republic of Korea, in a home garden located on a hill in Kunryang-
 19 ri (36°26'16.2"N 126°46'04.6"E), Cheongyang-eup, Cheongyang, Chungnam Province, from a
 20 woodlouse (*Poc. scaber*), July 2022, H.B. Lee & J.S. Kim (culture CNUFC CY22014).

21 *Notes*: *Penicillium charlesii* has been isolated from mouldy corn (*Zea mays* L.), soil,
 22 surface water, greenhouses used for shiitake cultivation, cream, pasteurised milk, tomato, sputum,
 23 underground sites of Brestovská Cave, body surface and gastrointestinal tract of oribatid mites,
 24 corn kernels (Sun et al., 2021; Ahn et al., 2016; Raghuwanshi et al., 2013; Devi et al., 2012;
 25 Houbraken and Samson, 2011), and from *Porcellio scaber* in the current study. This species has
 26 been reported in UK, Italy, Japan, Portugal, Brazil, and Korea (Sun et al., 2021; Ahn et al., 2016).

27
 28 ***Penicillium ferraniaense*** Houbraken & Di Piazza, Persoonia 46: 491 (2021) (Fig. 5A-E)

29 *Colony diameter, 7 days at 25 °C (mm)*: CYA 19–24; MEA, 25–27; YES 19–25.

30 *Colony characteristics*: CYA 25 °C, 7 days: Colonies raised at the center, radially sulcate;
 31 mycelia white; sporulation strong; soluble pigments absent; exudates absent; reverse yellowish
 32 white to yellowish gray. MEA 25 °C, 7 days: Colonies radially sulcate, concentric rings at the
 33 center; mycelia white; texture velvety; sporulation moderate; soluble pigments absent; exudates
 34 absent; reverse strong orange at the center, pale yellow at the edge. YES 25 °C, 7 days: Colonies
 35 sulcate; texture velvety; sporulation strong; soluble pigments absent; exudates absent; reverse
 36 pale yellow.

37 *Micromorphology*: Conidiophores monoverticillate, minor portion biverticillate; stipes
 38 smooth-walled, 55–80 × 2.5–3.5 µm. Phialides 5–7 per metula, ampulliform, 7–11 × 2–3 µm.
 39 Conidia smooth-walled, globose to subglobose, 2.5–3.5 µm.

40 *Material examined*: Republic of Korea, Chonnam National University, Yongbong-dong,
 41 Buk-gu, Gwangju (35°10'28.8"N 126°54'32.3"E), from the soil, February 2022 (culture CNUFC
 42 JYS28).

43 *Notes*: The morphological descriptions of CNUFC JYS28 matched well with the
 44 previously described species of *P. ferraniaense* (Crous et al., 2021). CNUFC JYS28 differed from
 45 *P. ferraniaense* in colony diameter on CYA (19–24 mm vs 25–28 mm). *Penicillium ferraniaense*

1 was isolated from anaerobic digestate (compost) in a previous study (Crous et al., 2021), and from
2 soil in the present study.

3
4 ***Penicillium glaucoroseum*** Demelius, Verhandlungen der Zoologisch-Botanischen Gesellschaft
5 Wien 72: 2 (1923) (Fig. 5F-I)

6 *Colony diameter, 7 days at 25 °C (mm):* CYA 21–23; MEA 32–37; YES 19–23.

7 *Colony characteristics:* CYA 25 °C, 7 days: Colonies slightly radially sulcate; margins
8 low, narrow; mycelia white; texture floccose; sporulation sparse; soluble pigments absent;
9 exudates absent; reverse yellowish white. MEA 25 °C, 7 days: Colonies raised at the center;
10 margins moderately wide; mycelia white; texture floccose; sporulation sparse; reverse pale
11 greenish yellow. YES 25 °C, 7 days: Colonies radially sulcate, sunken centrally; margins low,
12 wide; mycelia white; texture velvety; soluble pigments absent; exudates absent; reverse pale
13 yellow.

14 *Micromorphology:* Conidiophores biverticillate, few terverticillate; stipes rough walled,
15 105–170 × 2.0–3.0 µm. Phialides 3–4 per metula, ampulliform, 7–12 × 2–2.5 µm. Conidia
16 smooth-walled, globose, 2.5–3.5 µm. Ascospores not observed.

17 *Material examined:* Republic of Korea, Janggok-ri (36°25′00.6″N 126°51′31.8″E),
18 Daechi-myeon, Cheongyang, Chungnam Province, from soil, February 2022, H.B. Lee (culture
19 CNUFC CY22324).

20 *Notes:* The observed characteristics were mostly similar to those previously described for
21 *P. glaucoroseum* (Visagie et al. 2015). However, isolate CNUFC CY22324 grew slower than *P.*
22 *glaucoroseum* (CYA 47–49 mm; MEA 54–56 mm; YES 50–53 mm) (Visagie et al. 2015).
23 *Penicillium glaucoroseum* was isolated from live roots of the coniferous tree kauri (*Agathis*
24 *australis*), soil, worm excreta, farm sludge, and activated human sludge in previous studies
25 (Taghavi et al., 2021; Demelius et al., 1922), and from soil in the current study.

26
27 ***Penicillium jeongsukae*** Hyang B. Lee & T.T.T. Nguyen **sp. nov.**

28 *Index Fungorum:* IF 558124; Fig. 5J-N and Table 2

29 *Etymology:* jeongsukii, named after Jeong Suk Kim. This species was isolated from dry
30 persimmons collected by her.

31 *Type specimen:* Republic of Korea, Kunryang-ri (36°26′16.2″N 126°46′04.6″E),
32 Cheongyang-eup, Cheongyang, Chungnam Province, from Korean dried persimmon, October 25,
33 2021, J.S. Kim (holotype CNUFC HT2102; ex-type culture CNUFC JSK815).

34 *Colony diameter, 7 days (mm):* CYA 25 °C 35–37, CYA20 °C 22–24, CYA30 °C 40–43,
35 CYA 37 °C 30–34, CYA39 °C no growth; MEA25 °C, 40–42; YES25 °C 31–33; OA25 °C 36–
36 38; CREA 25 °C 24–26; CYAS 25 °C 15–18; DG18 25 °C 22–25.

37 *Colony characteristics:* CYA 25 °C, 7 days: Colonies slightly radially sulcate; margins
38 entire; texture floccose; mycelium white; soluble pigments absent; exudates absent; sporulation
39 moderate; conidia green; reverse grayish. YES 25 °C, 7 days: Colonies radially sulcate; mycelium
40 white; sporulation poor, exudates absent; soluble pigments absent; reverse pale yellow. MEA
41 25 °C, 7 days: Sporulation strong; colonies low, plane; mycelium white; margins low; exudates
42 absent; soluble pigments absent; reverse yellowish white. CYA 25 °C, 7 days: Colonies raised
43 centrally; margins entire; texture floccose; sporulation moderate; mycelium white; soluble
44 pigments absent; exudates absent; reverse light yellow. OA 25 °C, 7 days: Colonies low, plane;

1 margins entire; mycelia white; sporulation moderate; soluble pigments absent; exudates absent;
2 reverse light grayish. CREA 25 °C, 7 days: Colonies showing good growth; acid produced.

3 *Micromorphology*: Sclerotia absent. Conidiophores 17.5–40(–45) × 2–3.5 µm, with
4 smooth-walled stipes, monoverticillate. Vesicles (2.5–)3.0–6.0 µm. Phialides ampulliform, 3–15
5 per vesicle, 6.5–10 × 2.0–3.0 µm. Conidia in long chains, smooth-walled, predominantly broadly
6 ellipsoidal or cylindrical, 2.5–3.5 × (1.5–)2–3 µm.

7 *Additional material examined*: Republic of Korea, Kunryang-ri
8 (36°26'16.2"N126°46'04.6"E), Cheongyang-eup, Cheongyang, Chungnam Province, from
9 Korean dried persimmon, October 25, 2021, J.S. Kim (culture CNUFC JSK816).

10 *Notes*: In a NCBI BLASTn search based on ITS, *BenA*, *CaM* and *RPB2* sequences, the
11 closest match of *P. jeongsukae* was 99.15% with *P. lunae* CMV006E6 (MK450725) (for ITS),
12 96.20% with *P. cuddlyae* CMV016A6 (MK951835) (for *BenA*), 96.43% with *P. cuddlyae*
13 CMV016A6 (MK951908) (for *CaM*), and 98.06% with *P. chermesinum* CBS 231.81 (MN969111)
14 (for *RPB2*).

15 Phylogenetic analysis revealed that the new species was distinct and formed a different
16 clade in sect. *Charlesia*. Good growth on Czapek yeast autolysate (CYA) agar at 37 °C, fewer
17 phialides per vesicle, shorter conidiophores, and predominantly broadly ellipsoidal or cylindrical
18 conidia distinguished the new species from other known species.

19
20 ***Penicillium sajarovii*** Quintan., Av. Aliment. Majora Anim.: 539 (1981) (Fig. 5O-R)

21 *Colony diameter, 7 days at 25 °C (mm)*: CYA 26–28; MEA, 26–29; YES 22–24.

22 *Colony characteristics*: CYA 25 °C, 7 days: Colonies plane to slightly sulcate; texture
23 velvety; texture floccose; mycelium white; sporulation dense; soluble pigments absent; exudates
24 absent; reverse light yellowish brown at the center. MEA 25 °C, 7 days: Colonies plane; texture
25 floccose; mycelium white; sporulation moderate; soluble pigments absent; exudates absent;
26 reverse light yellow. MEA 25 °C, 7 days: Colonies sulcate, slightly raised at the center; mycelium
27 white; sporulation sparse; soluble pigments absent; exudates absent; reverse vivid yellow.

28 *Micromorphology*: Conidiophores biverticillate, 130–181 × 2.0–3.5 µm. Phialides
29 ampulliform, 5.5–8.5 × 2.0–3 µm. Conidia globose to subglobose, 2.5–3.0 µm.

30 *Material examined*: Republic of Korea, Gitdaebong Peak, Ulleung Island, Korea
31 (37°30'04"N 130°51'23"E), from soil, May 01, 2019 (culture CNUFC U734).

32 *Penicillium sajarovii* was isolated from rye (*Secale cereale*) as endophyte, the Himalayan
33 herb *Arnebia euchroma*, and the rhizosphere of diseased white lupin (*Lupinus albus*) in previous
34 studies (Jain et al. 2021; Quintanilla 1981), and from soil in the current study.

35 36 **4. Discussion**

37
38 Saprotrophic fungi would be expected in humic soils, but perhaps more surprisingly were also
39 present in freshwater and on seeds. Dried fruits represent a high-sugar, low water-activity habitat
40 for fungi that favours the growth of only xerotolerant and xerophilic species (Lievens et al., 2015).
41 The *Eurotiales* is known to contain a considerable number of very xerotolerant species (see
42 Stevenson et al., 2015; 2017) so it is unsurprising that we isolated such fungi from dried
43 persimmon fruits in the current study. Fungal saprotrophs not only degrade dead plant matter, but
44 in some cases degrade insect exoskeletons, fish scales, and animal hair (Christian et al., 2011).

1 Interesting, freshwater and seawater can also be a source of saprotrophic fungi and can even
2 act to preserve them long-term in an inactive condition (Hallsworth, 2022; Jones et al., 2014). So
3 far, a total of 3,870 species were reported from freshwater, with only 76 species belonging to the
4 order *Eurotiales* (Calabon et al., 2022; Nguyen et al., 2021b). Recently, Torres-Garcia et al. (2022)
5 reported eight *Penicillium* species from Spanish freshwater sediments so the number of known
6 *Eurotiales* species from freshwater has increased to at least 84. Likely, a large number of species
7 are yet to be discovered in freshwater habitats. Whereas we had expected to identify diverse
8 *Eurotiales* fungi in the current study, the discovery of two new species was unexpected. Given
9 the various crises currently facing humanity and the biosphere (habitat destruction including
10 deforestation, wildfires, climate change, biodiversity loss, etc), we believe that it is as important
11 as ever to continue to document species not only for the scientific value, but also for their potential
12 worth in biotechnological solutions to the various problems humanity is facing (e.g., Purchase,
13 2016; Timmis and Ramos 2021; Anand et al., 2023).

14 The new *Aspergillus* species identified in this study was located in sect. *Flavipedes* and ser.
15 *Spelaei* and was found to be related to *A. alboviridis*, *A. lanuginosus* and *A. movilensis* in
16 individual and combined phylogenetic analyses with ITS, *BenA*, *CaM* and *RPB2* sequences.
17 Production of Hülle cells, and the inability to grow on CYA at 37 °C can be used to distinguish
18 between *A. ullungdoensis* and *A. lanuginosus* (that does not produce Hülle cells and can grow on
19 CYA at 37 °C). In addition, *A. ullungdoensis* grows faster than *A. lanuginosus* on MEA at 25 °C
20 (20–21 mm). Like *A. ullungdoensis*, *A. alboviridis*, and *A. movilensis* also produce Hülle cells,
21 but conidiophores of *A. ullungdoensis* are longer than *A. alboviridis* [100–200(–550) µm] and *A.*
22 *movilensis* (usually < 400 µm) (Sklenář et al., 2021). *Aspergillus ullungdoensis* grows faster than
23 *A. alboviridis* on MEA (18–23 mm) and OA (18–19 mm) at 25°C (Sklenář et al., 2021).
24 *Aspergillus alboviridis* and *A. movilensis* grow at 37 °C, but *A. ullungdoensis* cannot grow at
25 37 °C. Sklenář et al. (2021) revealed that length of stipes can be considered taxonomically
26 important and may be used to distinguish *A. alboviridis*, *A. movilensis* and *A. lanuginosus*. Our
27 data demonstrate that *A. ullungdoensis* is morphologically and genetically different from *A.*
28 *lanuginosus*, *A. alboviridis*, and *A. movilensis*. Therefore, it is a newly described species.

29 Species in sect. *Flavipedes* occurs globally in soil, indoor or outdoor environment, as
30 endophytes, food contaminants, dung of herbivores, cave environments including bat guano-
31 contaminated soils, and clinical samples (Sklenář et al., 2021; Visagie et al., 2021). Sect.
32 *Flavipedes* also contains species that are widely used in biotechnology industries (Sklenář et al.,
33 2021; Frisvad and Larsen, 2015). *Aspergillus iizukae* is known to produce oxidative enzymes with
34 an industrial application (Noman et al., 2020) and the potential antiviral activity compounds
35 against influenza virus (H1N1), herpes simplex virus types 1 (HSV-1) and 2 (HSV-2) (Kang et
36 al., 2018). Metabolites 3-hydroxy-2',4,4',6'-tetramethoxyhalocone, 7,3,4,5'-
37 tetramethoxyflavanone, isovitexin, and amodiaquine produced by *A. flavipes* were found effective
38 antimicrobial against *Phytophthora parasitica* (El-Sayed and Ali, 2020). Therefore, new
39 *Aspergillus* species could be promising candidate for product discovery.

40 The new *Penicillium* species belongs to sect. *Charlesia* and ser. *Indica* based on multigene
41 phylogeny data using ITS, *BenA*, *CaM*, and *RPB2* sequences. ITS, *BenA*, *CaM*, and *RPB2* data
42 revealed that the novel species was distinct from previously described *Penicillium* species. The
43 data of independent and combined sequence analyses of the four loci revealed a close relationship
44 among *P. chermesinum*, *P. indicum*, *P. lunae*, and *P. cuddlyae*. *Penicillium jeongsukae* shares
45 several similarities with *P. chermesinum*, *P. cuddlyae*, *P. indicum*, and *P. lunae*, such as the

1 production of monoverticillate sporangiophores and growth on CYA agar at 37 °C (except for *P.*
2 *lunae*) (Crous et al., 2019a, 2019b). However, *P. jeongsukae* differs from *P. lunae* (MEA: 25–26
3 mm) and *P. cuddlyae* (21–23 mm) by its capacity for rapid growth on CYA agar, MEA, and OA
4 (Crous et al. 2019a, 2019b). Species in sect. *Charlesia* have been isolated worldwide from various
5 habitats, such as indoor air, banana fruit, tomato, garlic, rice, beans, dried blue peas, spoiled
6 cereals, other dry foods, rhizosphere of peanut plants, stem and root of mangrove tree, soil,
7 greenhouses used for shiitake cultivation, cream, pasteurised milk, sputum, body surface and
8 gastrointestinal tract of mites (*Heritiera littoralis*), and inner tissues of the marine macroalga
9 *Pterocliadiella tenuis* (Crous et al., 2019a, 2019b; Su et al., 2018; Peterson et al., 2017; Yilmaz et
10 al., 2016). The current study is the first report of the isolation of *P. charlesii* from the woodlouse
11 *Por. scaber*. Members in sect. *Charlesia* are known to. Their ability to produce enzymes that are
12 efficient and potentially useful for anthropogenic purposes make them ideal candidates for
13 biotechnological applications (Raghuwanshi et al., 2013; Batra and Saxena, 2005; Clutterbuck et
14 al., 1935). *Penicillium charlesii* exhibits exo- α -mannosidase, β -glucosidase, protease, and
15 xylanase activities (Yoon et al., 2007; Abbas et al., 1989; Preston et al., 1973). It also exhibits
16 antinematodal and antiparasitic activities (Ondeyka et al., 1990). Moreover, it produces various
17 organic acids, such as carolic acid, itaconic acid, I- γ -methyltetronic acid, ramigenic acid, tartaric
18 acid, tetronic acid, and verticillic acid. Citreoviridin is another known mycotoxin produced by
19 this strain (Preston et al., 1973; Klatt et al., 1969; Bentley et al., 1962; Lybing et al., 1958;
20 Clutterbuck et al., 1935). *Penicillium costaricense* produces extrolites, such as andrastin A and
21 rastin C (Vega et al., 2006) and *Penicillium coffeae* produces δ -lactone (penicoffeazine A) and
22 isocoumarin derivatives (penicoffrazins B and C) (Cao et al., 2019; Peterson et al., 2005).
23 *Penicillium coffeae* was found to tolerate up to 500 mM of arsenate (Bhargavi and Savitha, 2014)
24 which could be a potential for bioremediation applications. Therefore, the new *Penicillium*
25 species isolated in this study could be a potentially useful source for producing secondary
26 metabolites.

27 Fungi of the genera *Aspergillus* and *Penicillium* are involved in humanitarian needs in
28 both positive and negative ways given their various roles in relation to agricultural economics,
29 the food sector, and biotechnology. Some species are plant pathogens, and others can cause food
30 spoilage, so have implications for food security. Some species are pathogenic to humans, and
31 their pathogenicity may become worse as the human population increases, malnutrition is
32 commonplace, and modern medical practices result in immuno-compromised patients.
33 Understanding fungal ecology (including mycotoxin production) is also key to keeping water safe
34 and preventing the proliferation of fungi harmful to human health in the built environment, and
35 fungi are key to soil- and plant health (Timmis and Ramos, 2021). In relation to climate change,
36 within the food-supply chain mycotoxins may become prevalent as weather patterns and
37 conditions for the transport and storage of foods and feeds become increasingly difficult to predict
38 and control (Medina et al., 2015) but microbes can be utilised for the common good including for
39 bioprospecting and biotechnological developments to mitigate the various humanitarian crises
40 (see also Timmis et al., 2024; Timmis and Hallsworth, 2024;). We are in favour of a systematic
41 and coordinated international effort to isolate fungi from local environments within regions
42 world-wide to discover more as-yet-undescribed fungi, bring them into accessible culture
43 collections, and assess them for their biotechnological potential.

44 **Acknowledgments**

1 This work was supported by the Survey and Discovery of Indigenous Fungal Species of Korea
2 Project (NIBR202203204) funded by National Institute of Biological Resources (NIBR) of the
3 Ministry of Environment (MOE), and by the Discovery of Fungi from Freshwater and their
4 Collection for Fungaria Project (NNIBR202201206) funded by Nakdonggang National Institute
5 of Biological Resources (NNIBR) of the Ministry of Environment (Korea). This study was also
6 supported by the Ministry of Science and ICT (Korea) (2022M3H9A1082984). The paper is
7 written in memory of Prof. Naresh Magan at Cranfield University (England, UK) who passed
8 away on April 20, 2023 (Hallsworth et al., 2023).

9 10 **References**

- 11
12 Aneke, C., Youn, J.H., Khil, P., Seyedmousavi, P.A., 2022. P148 Emerging and cryptic
13 *Aspergillus* species isolated from hospitalized patients with underlying primary
14 immunodeficiencies. *Med. Mycol. J.* 60, myac072P148.
- 15 Aveskamp, M.M., Verkley, G.J., de Gruyter, J., Murace, M.A., Perello, A., Woudenberg, J.H.,
16 Crous, P.W., 2009. DNA phylogeny reveals polyphyly of *Phoma* section *Peyronellaea* and
17 multiple taxonomic novelties. *Mycologia* 101(3), 363–82.
- 18 Calabon, M.S., Hyde, K.D., Jones, E.B.G. et al., 2022. Freshwater fungal numbers. *Fungal Divers.*
19 114, 3–235.
- 20 Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. TrimAl: a tool for automated
21 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25, 1972–1973.
- 22 Casadevall, A., Kontoyiannis, D.P., Robert, V., 2019. On the emergence of *Candida auris*:
23 Climate change, azoles, swamps, and birds. *MBio*, 10, e01397-19.
- 24 Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., Bakken, L.R., Baylis, M. et al., 2019.
25 Scientists' warning to humanity: microorganisms and climate change. *Nature Reviews*
26 *Microbiology* 17, 569–586.
- 27 Chen, A.J., Hubka, V., Frisvad, J.C., Visagie, C.M., Houbraken, J., Meijer, M., amson, R.A., 2017.
28 Polyphasic taxonomy of *Aspergillus* section *Aspergillus* (formerly *Eurotium*), and its
29 occurrence in indoor environments and food. *Stud. Mycol.* 88, 37–135.
- 30 Clutterbuck, P.W., Haworth, W.N., Raistrick, H., Smith, G., Stacey, M., 1935. Studies in the
31 biochemistry of micro-organisms: The metabolic products of *Penicillium charlesii* G.
32 Smith. *Biochem. J.* 29(4), 871–83.
- 33 Christian, W., Janice, K., Hans-Peter, G., 2011. Aquatic Fungi.1 In: *The Dynamical Processes of*
34 *Biodiversity: Case Studies of Evolution and Spatial Distribution*, Grillo, O. and G. Venora
35 (Eds.). Chapter 10, InTech Publisher, Rijeka, Croatia, ISBN: 978-953-307-772-7, pp: 227-
36 258.
- 37 Crous, P.W., Wingfield, M.J., Lombard, L., Roets, F., Swart, W.J., Alvarado, P., Carnegie, A.J.,
38 Groenewald, J.Z., 2019a. Fungal planet description sheets: 951–1041. *Persoonia* 43, 223–
39 425.
- 40 Crous, P.W., Carnegie, A.J., Wingfield, M.J., Sharma, R., Mughini, G., Noordeloos, M.E., Silva,
41 B.D.B., 2019b. Fungal Planet description sheets: 868–950. *Persoonia* 42, 291–473.
- 42 Crous, P.W., Cowan, D.A., Maggs-Kölling, G., Yilmaz, N., Thangavel, R., Wingfield, M.J.,
43 Groenewald, J.Z., 2021. Fungal Planet description sheets: 1182–1283. *Persoonia* 46, 313–
44 528.

- 1 Demelius, P., 1922. Beitrag zur Kenntnis der Hyphomyceten Niederösterreichs. Verh Zool-Bot
2 Ges Wien. 72, 66–109.
- 3 Frisvad, J.C., Hubka, V., Ezekiel, C.N., Hong, S.B., Nováková, A., Chen, A.J., Arzanlou, M.,
4 Larsen, T.O., Sklenář, F., Mahakarnchanakul, W., Samson, R.A., Houbraken, J., 2019.
5 Taxonomy of *Aspergillus* section *Flavi* and their production of aflatoxins, ochratoxins and
6 other mycotoxins. Stud. Mycol. 93, 1–63.
- 7 Frisvad, J.C., Samson, R.A., 2004. Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A
8 guide to identification of food and air-borne terverticillate *Penicillia* and their
9 mycotoxins. Stud. Mycol. 49, 1–174.
- 10 Garzon, C.D., Habrylo, O., Lemaire, A., Guillaume, A., Carre, Y., Millet, C., Pelloux, J., 2021.
11 Characterization of a novel strain of *Aspergillus aculeatinus*: from rhamnogalacturonan type
12 I pectin degradation to improvement of fruit juice filtration. Carbohydr. Polym. 262, 117943.
- 13 Glez-Peña, D., Gómez-Blanco, D., Reboiro-Jato, M., Fdez-Riverola, F., Posada, D., 2010.
14 ALTER: program-oriented format conversion of DNA and protein alignments. Nucleic
15 Acids Res. 38, 14–18.
- 16 Glass, N.L., Donaldson, G.C., 1995. Development of primer sets designed for use with the PCR
17 to amplify conserved genes from filamentous ascomycetes. Appl. Environ. Microbiol. 61,
18 1323–1330.
- 19 Hallsworth, J. E., Mswaka, A. Y., Patriarca, A., Verheecke-Vaessen, C., and Medina, A., 2023.
20 The life and works of Professor Naresh Magan. World Mycotoxin Journal 16: 195–197.
- 21 Hallsworth, J.E., 2019. Wooden owl that redefines Earth's biosphere may yet catapult a fungus
22 into space. Environmental Microbiology 21, 2202–2211.
- 23 Hallsworth, J.E., 2022. Water is a preservative of microbes. Microb. Biotechnol. 15(1), 191–214.
- 24 Hong, S.B., Cho, H.S., Shin, H.D., Frisvad, J.C., Samson, R.A., 2006. Novel *Neosartorya* species
25 isolated from soil in Korea. International Journal of Systematic and Evolutionary
26 Microbiology 56, 477–486.
- 27 Houbraken, J., Kocsubé, S., Visagie, C.M., Yilmaz, N., Wang, X.C., Meijer, M., Kraak, B., Hubka,
28 V., Bensch, K., Samson, R.A., Frisvad, J.C., 2020. Classification
29 of *Aspergillus*, *Penicillium*, *Talaromyces* and related genera (Eurotiales): An overview of
30 families, genera, subgenera, sections, series and species. Stud. Mycol. 95, 5–169.
- 31 Houbraken, J., Frisvad, J.C., Samson, R.A., 2011. Fleming's penicillin producing strain is
32 not *Penicillium chrysogenum* but *P. rubens*. IMA Fungus 2, 87–95.
- 33 Houbraken, J., Wang, L., Lee, H.B., Frisvad, J.C., 2016. New sections in *Penicillium* containing
34 novel species producing patulin, pyripyropens or other bioactive compounds. Persoonia 36,
35 299–314.
- 36 Houbraken, J., de Vries, R.P., Samson, R.A., 2014. Modern taxonomy of biotechnologically
37 important *Aspergillus* and *Penicillium* species. Adv. Appl. Microbiol. 86, 199–249.
- 38 Hubka, V., Nováková, A., Kolarík, A., 2014. Revision of *Aspergillus* section *Flavipedes*: seven
39 new species and proposal of section *Jani* sect. nov. Mycologia 107, 169–208.
- 40 Hubka, V., Barrs, V., Dudová, Z., Sklenář, F., Kubátová, A., Matsuzawa, T., Kolařík, M., 2018.
41 Unravelling species boundaries in the *Aspergillus viridinutans* complex (section *Fumigati*):
42 opportunistic human and animal pathogens capable of interspecific hybridization. Persoonia
43 41, 142–174.

- 1 Hyde, K.D., Hongsanan, S., Jeewon, R., Bhat, D.J., McKenzie, E.H.C., Jones, E.B.G., et al., 2016.
2 Fungal diversity notes 367–490: taxonomic and phylogenetic contributions to fungal taxa.
3 Fungal Divers. 80, 1–270.
- 4 Hyde, K.D., Xu, J., Rapior, S., et al., 2019. The amazing potential of fungi: 50 ways we can
5 exploit fungi industrially. Fungal Divers. 97, 1–136.
- 6 Jain, R., Bhardwaj, P., Pandey, S.S., Kumar, S., 2021. *Arnebia euchroma*, a plant species of cold
7 desert in the Himalayas, harbors beneficial cultivable endophytes in roots and leaves. Front
8 Microbiol. 12, 696667.
- 9 Jones, E.B.G., Hyde, K.D., Pang, K.L., 2014. Freshwater fungi and fungal-like organisms. Boston:
10 De Gruyter.
- 11 Jurjević, Ž., Peterson, S.W., Stea, G., Solfrizzo, M., Varga, J., Hubka, V., Perrone, G., 2012. Two
12 novel species of *Aspergillus* section *Nigri* from indoor air. IMA Fungus 3, 159–173.
- 13 Katoh, K., Rozewicki, J., Yamada, K.D., 2019. MAFFT online service: multiple sequence
14 alignment, interactive sequence choice and visualization. Brief. Bioinform. 20, 1160–1166.
- 15 Kedare, S.B., Singh, R.P., 2011. Genesis and development of DPPH method of antioxidant assay.
16 J. Food. Sci. Technol. 48, 412–422.
- 17 Khan, A.L., Hamayun, M., Kim, Y.-H., Kang, S.M., Lee, J.-H., Lee, I.-J., 2011. Gibberellins
18 producing endophytic *Aspergillus fumigatus* sp. LH02 influenced endogenous
19 phytohormonal levels, isoflavonoids production and plant growth in salinity stress. Process
20 Biochem. 46, 440–447.
- 21 Kim, D.Y., Pyo, A., Ji, S., You, S.H., Kim, S.E., Lim, D., Kim, H., Lee, K.H., Oh, S.J., Jung, Y.R.,
22 Kim, U.J., Jeon, S., Kwon, S.Y., Kang, S.R., Lee, H.B., Hyun, H., Kim, S.Y., Moon, K.S.,
23 Lee, S., Kang, S.J., Min, J.J., 2022. In vivo imaging of invasive aspergillosis with 18F-
24 fluorodeoxyisotriton positron emission tomography. Nat. Commun. 13, 1926.
- 25 Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: molecular evolutionary genetics analysis
26 version 7.0 for bigger datasets. Mol. Biol. Evol. 33, 1870–1874.
- 27 Landrigan, P.J., Stegeman, J.J., Fleming, L.E., Allemand, D., Anderson, D.M., Backer, L.C. et
28 al., 2020. Human health and ocean pollution. Annals of Global Health 86, 151.
- 29 Liu, Y.J., Whelen, S., Hall, B.D., 1999. Phylogenetic relationships among ascomycetes: evidence
30 from an RNA polymerase II subunit. Mol. Biol. Evol. 16, 1799–1808.
- 31 Lybing, S., Reio, L., 1958. Degradation of 14C-labelled carolic and carlosic acids from
32 *Penicillium charlesii* G. Smith. Acta Chem. Scand. 12, 1575–1584.
- 33 Lamboni, Y., Nielsen, K.F., Linnemann, A.R., Gezgin, Y., Hell, K., Nout, M.J., Frisvad, J.C.,
34 2016. Diversity in secondary metabolites including mycotoxins from strains of *Aspergillus*
35 section *Nigri* isolated from raw cashew nuts from Benin, West Africa. PLoS One 11,
36 e0164310.
- 37 Langlois, I., Barrs, V.R., Dufresne, P.J., 2021. Corrigendum to ‘Rhinitis due to *Aspergillus*
38 *pseudoviridinutans* in an orange-winged Amazon parrot (*Amazona amazonica*)’ [Med.
39 Mycol. Case Rep. 30 (2020) 46–50]. Med. Mycol. Case. Rep. 33, 38.
- 40 Lee, H.B., Magan, N., 1999. Environmental factors and nutritional utilization patterns affect niche
41 overlap indices between *Aspergillus ochraceus* and other spoilage fungi. Lett. Appl.
42 Microbiol. 28(4), 300–304.
- 43 Machida, M., Yamada, O., Gomi, K., 2008. Genomics of *Aspergillus oryzae*: learning from the
44 history of *Koji* mold and exploration of its future. DNA Res. 15, 173–83.

- 1 Magan, N., 2006. Mycotoxin contamination of food in Europe: early detection and prevention
2 strategies. *Mycopathologia* 162(3), 245–253.
- 3 Magan, N., Aldred, D., 2005. Conditions of formation of ochratoxin A in drying, transport and in
4 different commodities. *Food Addit. Contam.* 22, 10–16.
- 5 Medina, A., Rodríguez, A., Magan N., 2015. Climate change and mycotoxigenic fungi: Impacts
6 on mycotoxin production. *Current Opinion in Food Science* 5, 99–104.
- 7 Meyer, V., Wu, B., Ram, A.F., 2011. *Aspergillus* as a multi-purpose cell factory: current status
8 and perspectives. *Biotechnol. Lett.* 33(3), 469–476.
- 9 Micheluz, A., Pinzari, F., Manente, S., Rivera-Valentín, E.G. & Hallsworth, J.E., 2022.
10 Biophysical manipulation of the extracellular environment by *Eurotium halophilicum*.
11 *Pathogens* 11, 1462.
- 12 Nguyen, T.T.T., Noh, K.J.K., Lee, H.B., 2021a. New species and eight undescribed species
13 belonging to the families Aspergillaceae and Trichocomaceae in Korea. *Mycobiology.* 49,
14 534–550.
- 15 Nguyen, T.T.T., Frisvad, J.C., Kirk, P.M., Lim, H.J., Lee, H.B., 2021b. Discovery and extrolite
16 production of three new species of *Talaromyces* belonging to sections *Helici* and *Purpurei*
17 from freshwater in Korea. *J. Fungi* 7, 722.
- 18 Nguyen, T.T.T., Pangging, M., Bangash, N.K., Lee, H.B., 2020. Five new records of the family
19 Aspergillaceae in Korea, *Aspergillus europaeus*, *A. pragensis*, *A. tennesseensis*, *Penicillium*
20 *fluviserpens*, and *P. scabrosum*. *Mycobiology* 48(2), 81-94.
- 21 Nicoletti, R., Carella, A., Cozzolina, E., 2008. Investigation on fungal antagonists of root rot
22 agents from the rhizosphere of white lupin (*Lupinus albus*). *Dyn. Soil Dyn. Plant.* 2, 69–72.
- 23 Noel, D., Gelhaye, E., Darnet, S., Hallsworth, J.E., Sormani, R. & Morel-Rouhier, M., 2023.
24 Modes-of-action of antifungal compounds: Stressors and (target-site-specific) toxins,
25 toxicants, or *toxin-stressors*. *Microbial Biotechnology* 16, 1438–1455.
- 26 Noonim, P., Mahakarnchanakul, W., Varga, J., Frisvad, J.C., Samson, R.A., 2008. Two novel
27 species of *Aspergillus* section *Nigri* from Thai coffee beans. *Int. J. Syst. Evol. Microbiol.* 58,
28 1727–1734.
- 29 O'Donnell, K., Cigelnik, E., 1997. Two divergent intragenomic rDNA ITS2 types within a
30 monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular and*
31 *Phylogenetic Evolution* 7, 103-116.
- 32 Pangging, M., Nguyen, T.T.T., Lee, H.B., 2022. Seven undescribed *Aspergillus* species from
33 different niches in Korea. *Mycobiology* 50(4), 189–202.
- 34 Paulussen, C., Hallsworth, J.E., Álvarez-Pérez, S., Nierman, W.C., Hamill, P.G., Blain, D. et
35 al., 2017. Ecology of aspergillosis: insights into the pathogenic potency of *Aspergillus*
36 *fumigatus* and some other *Aspergillus* species. *Microbial Biotechnology* 10, 296–322.
- 37 Perrone, G., Gallo, A., 2017. *Aspergillus* species and their associated mycotoxins. *Mycotoxigenic*
38 *fungi: Methods Mol Biol.* 1542, 33–49.
- 39 Perrone, G., Varga, J., Susca, A., Frisvad, J.C., Stea, G., Kocsubé, S., Tóth, B., Kozakiewicz, Z.,
40 Samson, R.A., 2008. *Aspergillus uvarum* sp. nov., an uniseriate black *Aspergillus* species
41 isolated from grapes in Europe. *Int. J. Syst. Evol. Microbiol.* 58, 1032–1039.
- 42 Peterson, S.W., Vega, F., Posada, F., Nagai, C., 2005. *Penicillium coffeae*, a new endophytic
43 species isolated from a coffee plant and its phylogenetic relationship to *P. fellutanum*, *P.*
44 *thiersii* and *P. brocae* based on parsimony analysis of multilocus DNA sequences.
45 *Mycologia* 97, 59–666.

- 1 Pol, D., Laxman, R.S., Rao, M., 2012. Purification and biochemical characterization of
2 endoglucanase from *Penicillium pinophilum* MS 20. Indian J. Biochem. Biophys. 49, 189–
3 194.
- 4 Purchase, D. (Ed.), 2016. Fungal Applications in Sustainable Environmental Biotechnology.
5 Springer Cham
- 6 Qi, T.F., Renaud, J.B., McDowell, T., Seifert, K.A., Yeung, K.C., Sumarah, M.W., 2016.
7 Diversity of mycotoxin-producing black aspergilli in Canadian vineyards. J. Agric. Food
8 Chem. 64, 1583–1589.
- 9 Qiao, W., Ling, F., Yu, L., Huang, Y., Wang, T., 2017. Enhancing taxol production in a novel
10 endophytic fungus, *Aspergillus aculeatinus* Tax-6, isolated from *Taxus chinensis* var. *mairi*.
11 Fungal Biol. 121, 1037–1044.
- 12 Quintanilla, J. A., 1981. Una nueva especie de *Penicillium* aislada en centeno: *P. sajarovii*, sp.
13 nov. A.Y.M.A. XXII: 539–541.
- 14 Ráduly, Z., Szabó, L., Madar, A., Pócsi, I., Csernoch, L., 2020. Toxicological and medical aspects
15 of *Aspergillus*-derived mycotoxins entering the feed and food chain. Front. Microbiol. 10,
16 2908.
- 17 Rambaut, A., 2009. Computer program distributed by the author. FigTree, version 1.3.1 Available
18 online: <http://www.treebioedacuk/software/figtree>.
- 19 Raper, K.B., Thom, C., 1949. A manual of the Penicillia. A manual of the Penicillia. 1–878.
- 20 Sklenář, F., Jurjević, Ž., Peterson, S.W., Kolařík, M., Nováková, A., Flieger, M., Hubka, V., 2020.
21 Increasing the species diversity in the *Aspergillus* section *Nidulantes*: Six novel species
22 mainly from the indoor environment. Mycologia 112, 342–370.
- 23 Sklenář, F., Jurjević, Ž., Zalar, P., Frisvad, J.C., Visagie, C.M., Kolařík, M., Houbraeken, J., Chen,
24 A.J., Yilmaz, N., Seifert, K.A., Cotton, M., Déniel, F., Gunde-Cimerman, N., Samson, R.A.,
25 Peterson, S.W., Hubka, V., 2017. Phylogeny of xerophilic aspergilli (subgenus *Aspergillus*)
26 and taxonomic revision of section *Restricti*. Stud. Mycol. 88, 161–236.
- 27 Sklenář, F., Jurjević, Ž., Houbraeken, J., Kolařík, M., Arendrup, M.C., Jørgensen, K.M., Siqueira,
28 J.P.Z., Gené, J., Yaguchi, T., Ezekiel, C.N., Silva Pereira, C., Hubka, V., 2021. Re-
29 examination of species limits in *Aspergillus* section *Flavipedes* using advanced species
30 delimitation methods and description of four new species. Stud. Mycol. 99, 100120.
- 31 Sandhu, D.K., Sandhu, R.S., 1963. A new species of *Penicillium* isolated from sputum. Can. J.
32 Bot. 41, 1273–1274.
- 33 Susca, A., Tanney, J.B., Varga, J., Kocsubé, S., Sziget, iG., Yaguchi, T., Frisvad, J.C., 2014.
34 Phylogeny, identification and nomenclature of the genus *Aspergillus*. Stud. Mycol. 78, 141–
35 173.
- 36 Taghavi, N., Singhal, N., Zhuang, W.Q., Baroutian, S., 2021. Degradation of plastic waste using
37 stimulated and naturally occurring microbial strains. Chemosphere 263, 127975.
- 38 Talbot, J.J., Barrs, V.R., 2018. One-health pathogens in the *Aspergillus viridinutans*
39 complex. Med. Mycol. J. 56, 1–12.
- 40 Tan, S., Yang, B., Liu, J., Xun, T., Liu, Y., Zhou, X., 2019. Penicillixanthone A, a marine-derived
41 dual-coreceptor antagonist as anti-HIV-1 agent. Nat. Prod. Res. 33(10), 1467–1471.
- 42 Timmis, K., Hallsworth, J. E., 2024. This is the *Age of Microbiology*: crucial roles of learned
43 societies and academies. Microb. Biotechnol. (in press).
- 44 Timmis, K., Hallsworth, J. E., McGenity, T. J., Armstrong, R., Colom Valiente, M. F., Karahan,
45 Z. C., Chavarría, M., et al. 2024. A concept for international societally-relevant microbiology

- 1 education and microbiology knowledge promulgation in society. *Microb. Biotechnol.* (in
2 press).
- 3 Torres-Garcia, D., Gené, J., García, D., 2022. New and interesting species of *Penicillium*
4 (*Eurotiomycetes, Aspergillaceae*) in freshwater sediments from Spain. *MycKeys* 86, 103–
5 145.
- 6 van der Merwe, K.J., Steyn, P.S., Fourie, L., Scott, D.B., Theron, J.J., 1965. Ochratoxin A, a toxic
7 metabolite produced by *Aspergillus ochraceus* Wilh. *Nature* 205(976), 1112–1113.
- 8 Visagie, C.M., Houbraken, J., Frisvad, J.C., Hong, S.B., Klaassen, C.H.W., Perrone, G., Seifert,
9 K.A., Varga, J., Yaguchi, T., Samson, R.A., 2014. Identification and nomenclature of the
10 genus *Penicillium*. *Stud. Mycol.* 78, 343–371.
- 11 Watanabe, K., Yaguchi, T., Hirose, D., 2021. Ubiquitous distribution of azole-resistant
12 *Aspergillus fumigatus*-related species in outdoor environments in Japan. *Med. Mycol. J.* 62,
13 71–78.
- 14 White, T.J., Bruns, T., Lee, S., 1990. Amplification and direct sequencing of fungal ribosomal
15 RNA genes for phylogenetics. *PCR protocols: A Guide to methods and Applications.* 315–
16 22.
- 17 Wierckx, N., Agrimi, G., Lübeck, P.S., Steiger, M.G., Mira, N.P., Punt, P.J., 2020. Metabolic
18 specialization in itaconic acid production: a tale of two fungi. *Curr. Opin. Biotechnol.* 62,
19 153–159.
- 20 Wu, J., Shui, H., Zhang, M., Zeng, Y., Zheng, M., Zhu, K.K., Wang, S.B., Cai, Y.S., 2023.
21 *Aculeaxanthon*es A–E, new xanthones from the marine-derived fungus *Aspergillus*
22 *aculeatinus* WHUF0198. *Front. Microbiol.* 14, 1138830.
- 23 Wu, J., Zhang, H., He, L.M. Xue, Y.Q., Jia, J., Wang, S.B., Cai, Y.S., 2021. A new fusicoccane
24 type norditerpene and a new indone from the marine-derived fungus *Aspergillus*
25 *aculeatinus* WHUF0198. *Chem. Biodivers.* 18, e2100562.
- 26 Zarei, F., Mirhendi, H., Fakhim, H., Geramixhoar, M., 2015. The first case of onychomycosis due
27 to *Aspergillus uvarum* (section *Nigri*). *Mycoses* 58, 239–242.
- 28 Zhao, S., Koffi, D., Latge, J.P. Syll, K., Gibbons, J.G., 2021. Genome sequence of *Aspergillus*
29 *aculeatinus* IC_8, ssolated from an indoor air sample of an urban housing complex in
30 abfidjan, ivory coast. *Microbiol. Resour. Announc.* 10(10):e00096-21.

32 **Table 1.** Morphological characteristics of *Aspergillus ullungdoensis* sp. nov. compared with those
33 of the reference strains *A. lanuginosus*, *A. alboviridis*, and *A. movilensis*

Characteristics	<i>A. ullungdoensis</i> ^(a)	<i>A. lanuginosus</i> ^(b)	<i>A. alboviridis</i> ^(b)	<i>A. movilensis</i> ^(b,c)
Size after 7 days at 25 °C (diameter)	23–26 mm on CYA 27–29 mm on MEA 23–27 mm on OA	26–27 mm on CYA 20–21 mm on MEA 25–26 mm on OA	19–21 mm on CYA 18–23 mm on MEA 18–19 mm on OA	22–25 mm on CYA 25–30 mm on MEA NI
Size after 7 days at 37 °C	No growth	Growth at 37 °C	Growth at 37 °C	10-17 mm
Stipes	Hyaline or light yellowish brown, smooth, not septate, 150–1,200 × 4–6 µm	Smooth, hyaline or light brown, 600–1,100 µm × 3.5–4.5 µm	Smooth, hyaline or brown, 120–200(–550) × 4.5–6(–8) µm	Usually < 400 µm length, 3.5–6 µm width

Vesicle	Globose, subglobose to pyriform, hyaline, 9–14 × 7.5–13 µm	Subglobose to pyriform, hyaline, 10–12 µm	Globose to subglobose, hyaline, 12–16(–20) µm	Pyriform, 5–16 µm
Metulae	5.5–8(–9.5) × 3–3.5 µm	Cylindrical, hyaline, 5–5.5 µm	Cylindrical, hyaline, 6–7(–9) µm	4–8(–10) µm length
Phialides	Ampulliform, 5.5–9(–10) × 2.5–3.0 µm	Flask-shaped, hyaline, 5.5–6.5 µm	Flask-shaped, hyaline, 6–7.5(–8.5) µm	4–7 µm length
Conidia	Globose, smooth, 2.5–3(–3.5) µm	Globose to subglobose, smooth, hyaline, 2.5–3 × 2–2.5 µm	Globose to subglobose, smooth, hyaline, 2.5–3.5 × 2–3 µm	Globose, smooth, 2.5–3.3(–3.5) µm
Hülle cells	Present	Present	Absent	Present

^(a) The present study; ^(b) From the description by Sklenář et al. (2021); ^(c) From the description by Hubka et al. (2015)

Table 2. Morphological characteristics of *Penicillium jeongsukae* sp. nov. compared with those of the reference strains *P. cuddlyae*, *P. chermesinum*, *P. lunae*, and *P. indicum*

Characteristics	<i>P. jeongsukae</i> ^(a)	<i>P. cuddlyae</i> ^(b)	<i>P. chermesinum</i> ^(c)	<i>P. lunae</i> ^(d)	<i>P. indicum</i> ^(e)
Size after 7 days at 25 °C (diameter)	35–37 mm on CYA 31–33 mm on YES 24–26 mm on CREA 40–42 mm on MEA 36–38 mm on OA 21–23 mm on DG18 15–18 mm on CYAS	24–26 mm on CYA 30–32 mm on YES 12–14 mm on CREA 21–23 mm on MEA 28–30 mm on OA 24–25 mm on DG18 19–20 mm on CYAS	40–50 mm on CYA NI on YES NI on CREA 50–55 mm on MEA NI on OA	34–36 mm on CYA 34–35 mm on YES NI on CREA 25–26 mm on MEA 28 mm on OA 24–25 mm on DG18 33–35 mm on CYAS	50 mm on CYA ^(*) NI on YES NI on CREA 55 mm on MEA ^(*) 50 mm on OA ^(*)
Size after 7 days at 37 °C	30–34 mm on CYA	19–21 mm on CYA	Growth at 37 °C	No growth	Growth at 37 °C
Conidiophores	Monoverticillate, stipes smooth, 17.5–40(–45) × 2–3.5 µm	Monoverticillate, stipes smooth-walled, 20–45 × 2–3 µm	Strictly monoverticillate, stipes smooth-walled, mostly 20–40(–50) × 2.0–2.5 µm	Monoverticillate, minor proportion biverticillate; stipes smooth-walled, 13–60 × 2–3(–3.5) µm	Monoverticillate, stipes smooth or finely roughened, 50–100 × 2.0–2.5 µm
Phialides	Ampulliform, 3–15 per vesicle, 6.5–10 × 2–3 µm.	Ampulliform, 10–20 per vesicle, 8–10 × 2–3 µm	Sterigmata crowded, often incurved, 10–15 per vesicle, 6–8 × 2.0–2.5 µm	Ampulliform, 10–20 per vesicle, (7.5–)8–10 × 2–3 µm	Sterigmata mostly in compact clusters, parallel, up to 12 or 15 per vesicle, 6.8–8.4 × 2.0–2.5 µm
Vesicle	(2.5–)3.0–5.5 µm wide	5–6 µm wide	4.0–4.5 µm wide	5–7 µm wide	3.4–6.5 µm wide

Metulae	NA	NA	NA	Two when present, 18–30 × 2–3(–3.5) µm	NA
Conidia	Smooth-walled, predominantly broadly ellipsoidal or cylindrical, 2.5– 3.5 × (1.5–)2–3 µm	Smooth-walled, ellipsoid or cylindrical, 2–3 × 2.5–2 µm	Smooth, elliptical, 2.0–2.5 × 1.5–2.0 µm	Smooth-walled, subglobose to broadly ellipsoid, 2–3(–3.5) × 1.5– 2(–2.5) µm	Smooth, elliptical, globose to subglobose, 1.7– 2.5(–3.0) µm

1 ^(a) The present study. ^(b) From the description by Crous et al. (2019a); ^(c) From the description by
2 Raper and Thom (1949); ^(d) From the description by Crous et al. (2019b). ^(e) From the description
3 by Sandhu and Sandhu (1963). NA: not available. ^(*) 10 days.

4 **Figure legends**

5
6
7 **Fig. 1.** Location of sample collection sites within Korea (A). Samples were collected from
8 different sites including soil at Cheongyang (B), water from waterfall freshwater at Jeju Island
9 (C), water from a freshwater lake at Gwangju (D) and agricultural soil Ulleung Island (E).

10
11 **Fig. 2.** Phylogram generated from maximum likelihood analysis based on combined ITS, *BenA*,
12 *CaM*, and *RPB2* sequences data of *Aspergillus* in ser. *Spelaei*. The numbers above or below
13 branches represent maximum likelihood bootstrap percentages (left) and Bayesian posterior
14 probabilities (right). Bootstrap values ≥ 70% and Bayesian posterior probabilities ≥ 0.95 are
15 indicated above or below branches. Bootstrap values lower than 0.95 and 70% are marked with
16 “-”. *Aspergillus urmiensis* CBS 139558 was used as an outgroup. The newly generated sequence
17 is indicated in blue. ^T = ex-type.

18
19 **Fig. 3.** Phylogram generated from maximum likelihood analysis based on combined ITS, *BenA*,
20 *CaM*, and *RPB2* sequences data of *Penicillium* in sect. *Charlesia*. The numbers above or below
21 branches represent maximum likelihood bootstrap percentages (left) and Bayesian posterior
22 probabilities (right). Bootstrap values ≥ 70% and Bayesian posterior probabilities ≥ 0.95 are
23 indicated above or below branches. Bootstrap values lower than 0.95 and 70% are marked with
24 “-”. *Penicillium chrysogenum* CBS 306.48 was used as an outgroup. The newly generated
25 sequence is indicated in blue. ^T = ex-type.

26
27 **Fig. 4.** Morphology of species in this study. (A, E, J, N, R, W) Colony morphology (top row, left
28 to right: CYA, MEA, YES); (bottom row, left to right: reverse colonies on CYA, MEA, YES).
29 (B, C, F, G, K, L, O, P, S, T, X, Y) Conidiophores. (D, H, M, Q, U, Z) Conidia. (I) Ascomata. (V)
30 Hülle cells. Scale bars = 10 µm.

31
32 **Fig. 5.** Morphology of species in this study. (A, F, J, O) Colony morphology (top row, left to right:
33 CYA, MEA, YES); (bottom row, left to right: reverse colonies on CYA, MEA, YES). (B-D, G,
34 H, K-M, P, Q) Conidiophores. (E, I, N, R) Conidia. Scale bars = 10 µm.

35
36
37
38

1 **Fig. 1**

2



3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19


20

21

22

1 **Fig. 2**

2



3

4

5

6

7

8

9

10


11

12

13

1 **Fig. 3**

2



3

4

5

6

7

8


9

10

11


12

1 Fig. 4



1 **Fig. 5**

2



3

Dear Chief-Editor,

I am pleased to submit the manuscript to renowned Fungal Biology as special issue. This manuscript has not been published elsewhere. All the authors have approved the manuscript and agree with this submission. There were no any interest conflicts among all authors.

The authors have no known competing financial interests or personal relationships that could affect the work reported in this paper.

Best regards,

Hyang Burm Lee PhD

Department of Agricultural Biological Chemistry,

College of Agriculture and Life Sciences,

Chonnam National University, Gwangju 61186, South Korea

+82-062-530-2136 (TEL); +82-062-530-2139 (FAX)

e-mail: hblee@jnu.ac.kr

Aspergillus ullungdoensis sp. nov., Penicillium jeongsukae sp. nov., and other fungi from Korea

Lee, Hyang Burm

2024-12-01

Attribution-NonCommercial-NoDerivatives 4.0 International

Lee HB, Nguyen TT, Noh SJ, et al., (2024) *Aspergillus ullungdoensis* sp. nov., *Penicillium jeongsukae* sp. nov., and other fungi from Korea. *Fungal Biology*, Volume 128, Issue 8, Part B, December 2024, pp. 2479-2492

<https://doi.org/10.1016/j.funbio.2024.05.014>

Downloaded from CERES Research Repository, Cranfield University