

## Synthesis, structure characterization, Hirshfeld surface analysis, and computational studies of 3-nitro-1,2,4-triazol-5-one (NTO): Acridine

Nilgün Şen<sup>\*1,2</sup>, Jean-François Pons<sup>2</sup>, Yunus Zorlu<sup>3</sup>, Eleftheria Dossi<sup>2</sup>, Federica Persico<sup>2</sup>, Tracey Temple<sup>2</sup>, Nazife Aslan<sup>4</sup>, Akachai Khumsri<sup>5</sup>

<sup>1</sup>Institute of Forensic Sciences, Turkish National Police Academy, Ankara, Türkiye

<sup>2</sup>Centre for Defence Chemistry, Cranfield University, Defence Academy of the United Kingdom, Shrivenham, SN6 8LA, UK

<sup>3</sup>Department of Chemistry, Gebze Technical University, 41400, Gebze, Kocaeli, Türkiye

<sup>4</sup>Department of Chemistry, Polatlı Science and Arts Faculty, Ankara Hacı Bayram Veli University, Ankara, Türkiye

<sup>5</sup>School of Chemistry, University of Edinburgh, Edinburgh, EH9 3FJ, UK

Corresponding author e-mail address: nilgunsen2001@gmail.com, nilgun.sen@pa.edu.tr

Telephone number: +90 505 295 5615, Fax number: +90 312 462 85 33

### Abstract

To modify the physical features and extend applications of the 3-nitro-1,2,4-triazol-5-one (NTO), we synthesized NTO with acridine (ACR) at a molar ratio of 1:1, a neutralization reaction. Through altering the chemical composition, it was possible to alter physical properties such as thermal stability, free space (voids), packing coefficient, crystal density, difference in pKa of co-formers, morphology, solubility, impact sensitivity, and calculated detonation parameters. It appears that physical attributes could be entirely altered. Single-crystal and powder X-ray diffraction methods, infrared spectroscopy, mass spectrometry, nuclear magnetic resonance spectroscopy (<sup>1</sup>H-NMR and <sup>13</sup>C-NMR), and thermal analysis were utilized to comprehensively characterize and confirm the formation of the structure of NTO: ACR. The substantial hydrogen bond interactions and planar layered structures observed between the cations and anions generated a complex 3D network, providing insight into the structure-property interrelationship. One intriguing feature discovered is the layered structure present in NTO:ACR, which may be responsible for the low impact sensitivity. According to the experimental results, NTO: ACR showed good thermal stability (Td = 229 °C) and outstanding impact sensitivity (IS = 100 J). Detonation velocity and pressure, were calculated using the EXPLO5 software program and found to be 7006 m·s<sup>-1</sup>, 20.02 G Pa, respectively.

**Keywords:** Energetic salts, 3-nitro-1,2,4-triazol-5-one, Acridine, Non-covalent interactions, NTO

## 1. Introduction

Energetic materials (EMs) include explosives, propellants, and pyrotechnics, are described as those that produce a substantial quantity of heat and/or gaseous products being stimulated by heat, impact, shock, spark, and so on. EMs's widely employed in military and civilian applications including weapons, mining, quarrying, demolition, emergency signalling, breathing device, vehicle safety, space exploration, and entertainment [1-3]. Significant efforts are being made to create high-performance EMs that have not merely outstanding in their combustion and detonation properties, high densities, and positive oxygen balances, but also have good thermal stability, sensitivity to external forces, low-cost synthesis, safe handling, and good environmental compatibility. [4-7]. Despite an almost 1200-year existence, they have gradually progressed in terms of detonation power as a basic feature. For instance, during the previous century, the detonation velocity of TNT (2,4,6-trinitrotoluene), HMX (1,3,5,7-tetranitro-1,3,5,7-tetrazocane) and CL-20 (hexanitrohexaazaisowurtzitane) has increased by just 30% [8-13]. The challenge of balancing high detonation performance with low sensitivity is difficult since the energy and sensitivity of explosives frequently conflict [14-17]. Instead of focusing solely on high energy, we can attempt to maximize the utilization of current compounds or develop new ones with remarkable, all-encompassing qualities.

3-nitro-1,2,4-triazol-5-one (NTO) is a heterocyclic carbonyl insensitive munition that was discovered in 1905, but its explosive characteristics were not studied until the 1980s [18]. It has a higher margin of safety against unanticipated stimulus reactions and outstanding physical qualities [19-22]. It exhibits energetic performance features comparable to those of currently used secondary explosives and has the potential to be used as an explosive and propellant ingredient [23]. This compound has been demonstrated to be less hazardous to human health than conventional explosives [24-25]. Nevertheless, this nitrogen heterocyclic energetic molecule has numerous limitations that limit its future applications, especially negative enthalpy of formation and slightly acidic nature ( $pK_a$  3.67) [26-27]. Furthermore, because of NTO's extremely solubility in water, it is likely to be found in substantial concentrations in effluent, depending on the formulation. In rats, NTO has a high  $LD_{50}$  relative to other EMs of around  $5000 \text{ mg.kg}^{-1}$ , and investigations of water toxicity studies have yielded comparable results, with toxicity levels so low that NTO might be categorized as non-toxic [28-30]. Nevertheless, at amounts greater than  $100 \text{ mg. L}^{-1}$ , NTO contamination of the water supply causes noticeable coloration, which can lead to environmental harm by perception as it displays a bright yellow tinge to water even if no ecotoxicological impact is shown [31].

To address these problems, researchers took two basic strategies: the initial one corresponds to the acidity of NTO-derivative salts, and the second relates to the development of NTO co-crystals. The first is the well-known technique, in which numerous metal and amine salts, as well as other NTO compounds, have been and continue to be made. The second new strategy, co-crystallization, might pave the road for the use of NTO. Co-crystallization is a prominent approach to enhancing the solubility, bioavailability, and physical and chemical stability of pharmaceuticals without affecting their chemical structure [32-35]. As a unique technology, co-crystals are composed of two or more components that are found together by hydrogen bonds,  $\pi - \pi$  stacking, van der Waals forces, and halogen bonds [36-38]. Co-crystallization is used to enhance the physicochemical features and energetic performance of the composition for certain applications. [39-42]. Different NTO co-crystals and salts have lately been studied theoretically and experimentally [19, 43-47]. It is demonstrated that this technology allows for the modification of physicochemical parameters and the creation of improved co-crystals with


enhanced features at the molecular level. ACR, an important nitrogen-containing heterocyclic systems, acting as a weak base ( $pK_b = 5.4$  in water) [48], and its salt display modest detonation performance. It may mitigate the acidity of NTO to some extent by co-crystallizing with mildly basic ACR, without consuming less energy and giving NTO an enhanced integrated performance.

The research disclosed in this paper aims to investigate a concept that has an opportunity to step-change for the control of the properties of EMs, leading in safer and more effective performance. The lack of structure-property relationships presents a challenge for the design of new NTO-based co-crystals and salts. This research thereby intends to exploit the concepts of crystal engineering and co-crystallisation as applied to selected energetic materials in order to accomplish the following objectives: (i) develop an enhanced understanding of how structure influences key properties (ii) control the sensitivity of existing, and (iii) identify new EM with improved energetic properties.

## 2. Experimental section


### 2.1. Chemicals and preparation of the sample

Sigma-Aldrich supplied sulfuric acid ( $H_2SO_4$ , 95-97%), semicarbazide hydrochloride ( $CH_6ClN_3O$ ), formic acid ( $CH_2O_2$ , 85%), nitric acid ( $HNO_3$ , 70%), and ACR ( $C_{13}H_9N$ , 97%). NTO was synthesized in two processes, beginning with the synthesis of TO (1,2,4-triazol-5-one). It was synthesized by the reaction of semicarbazide hydrochloride with formic acid and nitrated with 70%  $HNO_3$  (Scheme 1) [49]. NTO was verified using nuclear magnetic resonance and infrared spectroscopy (Figs. S1-S6).



**Scheme 1.** NTO synthesis process.

Crystallization was accomplished by dissolving an equimolar ratio of NTO (46 mg) with ACR (90 mg) in methanol (about 5-10 mL) at 50 °C and for 30 minutes (Scheme 2).



**Scheme 2.** Crystal formation mechanism of the NTO: ACR

## 2.2. Characterisation techniques

**2.2.a. Optical Microscopy.** Optical micrographs of the NTO, ACR, and NTO: ACR were examined using an Olympus BX51 microscope.

**2.2.b. Single Crystal X-ray Diffraction (SCXRD).** The measurements were carried out with a Bruker APEX II CCD three-circle diffractometer with MoK $\alpha$  ( $\lambda = 0.71073$ ) radiation. APEX2 was used to index the data (Bruker. APEX2. Ver. 2014.11-0, Bruker AXS Inc., Madison, WI 2014). SAINT was used for data integration and reduction (SAINT, version 8.34A, Bruker, Bruker AXS Inc., Madison, W. No. Title, 2013). SADABS' multi-scan approach was used to adjust absorption (SADABS, version 2014/5, Bruker, Bruker AXS Inc., Madison, W. No Title, 2014). NTO: ACR structure was solved by SHELXT [50] in Olex2 Software Package [51], then refined by full-matrix least-squares refinements on  $F^2$  using the SHELXL [52]. Additional crystallographic data with CCDC reference number **2253569** for NTO: ACR has been deposited within the Cambridge Crystallographic Data Centre *via* [www.ccdc.cam.ac.uk/deposit](http://www.ccdc.cam.ac.uk/deposit).

**2.2.c. Powder X-ray Diffraction (PXRD).** PXRD patterns were obtained using the Bruker D8Advance using Cu-K $\alpha$  radiation ( $\lambda = 1.54439$  Å) and an operating voltage and current of 40 kV and 40 mA, respectively. The temperatures ranged from 5 to 80 degrees.

**2.2.d. Nuclear Magnetic Resonance Spectroscopy (NMR) spectral studies.**  $^1\text{H}$ -NMR characterisation of ACR, NTO and NTO: ACR were obtained using a Bruker Ascend 400 MHz spectrometer with a BBFO probe, and the spectra recorded at ambient temperature in dimethyl sulfoxide (DMSO- $d_6$ ) (Typical solution of 10 mg/mL). The solvent peaks were referenced to DMSO- $d_6$  (2.50 ppm). Peak multiplicities were described as follows: singlet (s), triplet (t), broad triplet (brt), doublet of triplet (dt) and multiplet (m).

**2.2.e. Infrared (IR) Spectroscopy.** The IR spectra were obtained with a Bruker Equinox55 with ATR equipment (wavenumber range :4000-400  $\text{cm}^{-1}$ , resolution: 4  $\text{cm}^{-1}$ ).

**2.2.f. Differential Scanning Calorimetry (DSC).** Thermal characterization was performed on the materials using a Mettler Toledo LF1100 TGA/DSC 3+ equipped with a DSC sensor and operated by Stare System software v15.00 (build 8992). The tests were carried out in an inert atmosphere ( $\text{N}_2$  flowing at 50  $\text{cm}^3 \text{min}^{-1}$ ) and at one heating rates: 10  $^\circ\text{C min}^{-1}$  over a range of 25–400  $^\circ\text{C}$ .

**2.2.g. Mass spectrometry (MS).** Mass spectra were attained by an X500 QTOF mass spectrometer system equipped with a direct inlet (DI) unit and an electron impact ionizer. And the data was analysed with Bruker Compass Data Analysis systems.

**2.2.h. pH measurement.** Thermo Scientific's Orion 720Aplus and pH/mV/ISE Benchtop Meters were used to measure the pH values of NTO, ACR, and NTO:ACR. pH values were measured with combined glass pH electrode and found to be 3.48, 5.34 and 4.28, respectively (Fig S7).

Thermo Scientific™ Orion™ 720Aplus and pH/mV/ISE Benchtop Meters was used to measured pH solutions of NTO, ACR, and NTO: ACR.

### **2.3. Impact sensitivity measurement test**

The impact sensitivity was measured using a BAM fall hammer (BFH-12) and used 30-trial Bruceton method [54, 55].

### **2.4. Explosive performance analysis**

The EXPLO5 software program was used to calculate all explosive properties presented in this work, including detonation velocity, pressure, and oxygen balance [56].

### **2.5. Hirshfeld surface analysis**

Hirshfeld surface and 2D fingerprint calculations were carried out on crystal structures imported from CIF files using Crystal Explorer software version 3.1 [57].

## **3. Theoretical Calculations**

### **3.1. Calculation of heat of formation (HOF)**

In this study, ab initio calculations were performed using the program package Gaussian 09 (Revision-D.01) [58] and visualized with GaussView 5.0.8 [59]. SCXRD data was used to initial input data. For those computations, the hybrid B3LYP density functional with the def2-TVZP basis set in the gas phase was applied. Enthalpy values in the gas and solid phases were calculated using the complete basis set method (CBS-4M; M refers to the method of using minimum population localization).

### **3.2. Non-covalent interactions (NCIs)**


The NCI index is an excellent tool for studying non-covalent interactions. It has been frequently used to investigate hydrogen, halogen, and pi-pi stacking interactions [60, 61]. The basic density gradients were calculated using Multiwfn software (v.3.8) [62]. These gradients were then used for generating the demonstration graphs using the VMD (Visual Molecular Dynamics) program (v.1.9.3) [63].

## **4. Results and Discussion**

### **4.1. The outcomes of the experiments**

#### **4.1.1. The microscope images of NTO, ACR, NTO:ACR**

The microscope images of the NTO, ACR, and the NTO: ACR are presented in **Fig. 1**. The morphological distinctions between them are obvious. With the comparison of NTO and ACR, it came to light that the NTO: ACR was a yellow prism-like clear crystal with a well-defined morphology. The crystal shapes and colours are distinct from those of their co-formers (NTO in **Fig. 1a**, ACR in **Fig. 1b**, NTO: ACR in **Fig. 1c**).




**Fig. 1.** Microscope images of NTO (a), ACR (b), NTO: ACR (c)

#### **4.1.2. Characterization of the NTO: ACR**

##### ***SCXRD Analysis***


A new energetic compound, NTO: ACR crystals, which are appropriate for SCXRD analysis measurements, was attained by the slow solvent evaporation method. NTO: ACR was acquired in high yields and stable in an open atmosphere under ambient conditions. The compound (NTO: ACR) SCXRD analysis was performed by choosing good-quality crystals of sufficient size ( $0.15 \times 0.13 \times 0.12 \text{ mm}^3$ ). **Table S1** summarizes the crystallographic data attained and the structural refinement consequences of NTO: ACR. The ORTEP drawing and crystal structure of NTO: ACR are indicated in **Figs. 2a–b**. NTO: ACR forms crystals in the triclinic crystal system and corresponds to the *P*-1 space group with four molecules. (**Fig. 2c**). Four of the two molecules per unit cell have face-to-face connection, and each side is stacked with another molecule to form an edge-to-edge configuration (**Fig. 2d**). In line with our design concept, the significant to building a layer-by-layer structure in EMs is strategically opt for H-bond donor-acceptor units. The acridinium cation is the hydrogen bond donor, while NTO is the acceptor. The deprotonation of NTO and protonation of ACR result in a hydrogen bond between the carbonyl of NTO and the N of the acridine ( $\text{N}^+ \dots \text{H}^- \dots \text{O}$ ).



**Fig. 2.** ORTEP drawing of A) NTO: ACR, B) Crystal structure of NTO: ACR c) Structure of NTO: ACR in unit cell, d) Packing diagram in the unit cell of NTO: ACR.

The introduction of one hydrogen to the ACR, it altered the planer structure of the NTO precursor (**Fig.2D**). The heterodimers are held together to create a molecular layer. Given the rising quantity of eutectic molecules identified through this technology, co-crystallization to get compounds with unique characteristics has been intensively investigated. NTO act as anions and ACR (N-base) as cation (**Fig.3**). The presence of a (mildly) basic nitrogen of ACR helps forming the interactions seen, in particular the hydrogen bonding network. This is not surprising since the  $pK_a$  of NTO is  $\sim 3.7$  and the  $pK_a$  for ACR is  $\sim 5.4$  [48]. It has two ionizable hydrogen sites, which are crucial for salt formation; also, it is soluble in water, making it excellent for evaporative crystallisation. Furthermore, it is nitrogen-rich and contains functional groups that are favorable for interactions such as hydrogen bonding. For acid–base complexes whose  $\Delta pK_a$  values lie between -1 and 4, a linear relationship between the possibility of salt formation and their  $\Delta pK_a$  value has been established. This relationship has enabled us to quantify prior formulations of the  $pK_a$  rule, and it can be a useful tool for scientists involved in co-crystal design and salt selection strategies [64]. There are two moderate hydrogen bonds at the NTO: ACR ( $N_4-H_{4A} \dots O_3$  and  $N_5-H_5 \dots O_3$ ), which is already transferred onto the nitrogen atom (ACR) (**Fig.4b** and **Table 1**). Other interaction in **Fig.4a** are not hydrogen bonds; they should be regarded as necessary interactions that occur in the absence of stacking [65]. The formation of hydrogen bonds is a thermodynamically advantageous and exothermic reaction. Inter-molecular bonds of hydrogen decrease molecular volume and vapor pressure while increasing molar mass, viscosity, thermal conductivity, dissolving power, phase transition temperatures, dielectric constant, dipole moment, and surface tension [66].





**Fig. 3.** Calculated planes indicating NTO- stacking a), Calculated planes indicating zig-zag relationship of NTO- packing in NTO: ACR b)


Given the nature of electrostatic interactions, these interactions are extremely weak. The  $\pi \dots \pi$  stacking interaction occurs between two parallel ACR cations (center to center distance of 3.66 Å, Fig. 4a). IR spectrum data,  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra, and mass spectrometry were used to further characterize and support the SCXRD analysis of NTO: ACR structure.

**Table 1.** Hydrogen bond geometries for NTO: ACR [Å, °].

| D<br>(Donor) | H      | A<br>(Acceptor) | [ARU] <sup>a</sup> | D-H [Å] | H...A<br>[Å] | D...A<br>[Å] | D-H...A<br>[°] |
|--------------|--------|-----------------|--------------------|---------|--------------|--------------|----------------|
| N (4)        | H (4A) | O (3)           | [1]                | 0.86    | 1.9          | 2.759 (4)    | 174            |
| N (5)        | H (5)  | O (3)           | [2]                | 0.86    | 1.75         | 2.606 (3)    | 174            |

<sup>a</sup> Translation of ARU-code to CIF and equivalent position code: [1] = -x, 1-y, 3-z, [2] = x, y, -1+z, Note 1: Analysis of Potential Hydrogen Bonds and Schemes with  $d(\text{D} \dots \text{A}) < R(\text{D}) + R(\text{A}) + 0.50$ ,  $d(\text{H} \dots \text{A}) < R(\text{H}) + R(\text{A}) - 0.12$  Ang.,  $\text{D-H} \dots \text{A} > 100.0$  Deg. Note 2: \*HBs lengths and angles are measured on Mercury 4.1.3 [47].

N-H...O interactions are seen in the IR spectrum, but some signals are shifted. The N...H bands of the NTO at 3200  $\text{cm}^{-1}$  shifts around 3300  $\text{cm}^{-1}$ . This could also be the ACR N-H salt formation, or even an indication of the N-H...O=C hydrogen bond. Owing to formation of hydrogen bonds, stretching vibration of some signals shift about 100  $\text{cm}^{-1}$  to the low energy levels [67].




**Fig. 4.** The  $\pi \dots \pi$  stacking interaction and hydrogen-bonded pattern and in NTO:ACR.



### PXRD Analysis


The PXRD examination confirmed the formation of NTO: ACR, and the PXRD patterns of NTO, ACR, and NTO: ACR are presented in **Fig. 5**. The PXRD pattern of the salt (NTO: ACR) is clearly distinct from its co-formers, identifying the NTO: ACR as a novel compound rather than product of crystal transformation.



**Fig. 5.** PXRD patterns for ACR, NTO, and NTO: ACR.


#### 4.1.3. NMR spectral studies

To confirm the type of interaction of ACR with NTO molecules,  $^1\text{H}$ -NMR spectra of the single components and NTO:ACR were recorded in  $\text{DMSO-d}_6$  solution. The  $^1\text{H}$ -NMR spectrum of ACR (**Fig. 7**), shows all absorptions between 9.11 ppm and 7.60 ppm; a singlet at 9.11 ppm is attributed to H-7 proton, a multiplet centred at 8.17 ppm assigned to the overlapping signals of H-2, H-2', H-5 and H-5' and two sets of doublets of triplets assigned to H-3 and H3', H-4 and H-4' (**Fig.6**) at 7.86 ppm and 7.65 ppm respectively. The  $^1\text{H}$ -NMR spectrum of NTO shows a singlet at 12.83 ppm attributed to the NH protons.



**Fig. 6.** Chemical structure of ACR (left) with assigned numbered carbon atoms and NTO (right)

$^1\text{H}$ -NMR spectrum of the NTO: ACR (DMSO- $d_6$ ) shows the deshielding of all ACR protons (**Fig.7**, red line). In details, the singlet attributed to H-7 is deshielded by 0.03 ppm from 9.12 ppm to 9.15 ppm, and the multiplet assigned to H-2, H-2', H-5, and H-5' coalesces into a broad triplet and is deshielded by 0.02 ppm from 8.20–8.15 ppm to 8.22–8.15 ppm. The two multiplets assigned to H-3, H3', and H-4, H-4', are deshielded by 0.01 ppm to 7.90–7.85 ppm and by 0.02 ppm to 6.67–7.62 ppm, respectively. The small deshielding shifts observed by  $^1\text{H}$  NMR supports the formation of the weak acridinium-NTO salt (NTO: ACR) (**Scheme 2**), which is also confirmed by SCXRD analysis.



**Fig. 7.**  $^1\text{H}$ -NMR spectra of the ACR, NTO:ACR and ACR.HCl.

**Acridinium chloride:**  $^1\text{H}$ -NMR (DMSO  $d_6$  + DCl,  $\delta$ ) 7.93 (2H, m), 8.30 (2H, m), 8.51 (4H, brq,  $J=5.45$  Hz), 10.04 (1H, s).

**NTO: ACR:**  $^1\text{H}$ -NMR (DMSO  $d_6$ ,  $\delta$ ): 7.62 (1H, m), 7.86 (1H, m), 8.17 (1H, s), 9.12 (1H, s), 12.82 (1H, s).

**ACR:**  $^1\text{H}$ -NMR (DMSO  $d_6$ ,  $\delta$ ): 7.64 (2H, m), 7.86 (2H, m), 8.17 (4H, m), 9.11 (1H, s).

#### 4.1.3. Mass spectra (MS) analysis

The base peaks correspond to the ACR and NTO compounds. It can be displayed in **Fig. S8** at  $m/z$  180.08  $[\text{ACR}+\text{H}]^+$ ,  $[\text{C}_{13}\text{H}_{10}\text{N}_1]^+$  and at  $m/z$  129.00  $[\text{NTO}-\text{H}]^-$ ,  $[\text{C}_2\text{H}_1\text{N}_4\text{O}_3]^-$ . MS analysis only supports the presence of the two entities (NTO and ACR).

#### 4.1.4. DSC analysis

In order to estimate the thermal stabilities of NTO: ACR, DSC testing was performed. The DSC traces in **Fig.8** show that NTO: ACR salt show only one exothermic peak with onset temperature 226  $^{\circ}\text{C}$  and peak maximum 229  $^{\circ}\text{C}$ . The decomposition temperature has a lower value than that of pure NTO (279  $^{\circ}\text{C}$ ). The curves show that the thermal behaviour of NTO: ACR differs from that of their co-formers, and the variations in thermal stability of these substances imply the formation of a new compound. The formation of new compound could

lead to in inter molecular hydrogen bonding. Inter-molecular hydrogen bonds have been found to increase thermal conductivity, resulting in greater hot spot energy transmission, dissipation and greater insensitivity [2, 68, 69,70].




Fig. 8. Differential scanning calorimetry plots for NTO, ACR, and NTO: ACR [°C]

#### 4.1.5. Infrared (IR) Spectroscopy analysis

The IR spectra of NTO, ACR, and NTO: ACR are indicated in **Figs. S6, S9, S10**. The assignments for the most characteristic vibrational bands are listed in **Table 2**. It can be determined from the **Table 2** that a group of low intensity bands assigned to N–H stretching vibration of NTO decreased from the region of  $3203\text{ cm}^{-1}$  to  $3047\text{ cm}^{-1}$ . While the C=O stretching of NTO shifted from the region of  $1699\text{ cm}^{-1}$  to  $1650\text{ cm}^{-1}$ , and the C–N stretching vibration decreased from  $1188\text{ cm}^{-1}$  to  $1047\text{ cm}^{-1}$ . At the same time, some peak shift occurs for both ACR and NTO bonds. The hydrogen bond increases the X–H bond length, causing the corresponding stretching vibrations band in the IR spectrum to move towards lower frequencies [67].


Table 2. The main bands of the IR spectra of NTO, ACR, and NTO: ACR [ $\text{cm}^{-1}$ ]

| Assignment                                      | NTO  | NTO:ACR | ACR  |
|-------------------------------------------------|------|---------|------|
| C-H stretching aromatic vibration               |      |         | 3053 |
| N-H stretching vibration                        | 3203 | 3047    |      |
| C=O stretching vibration                        | 1699 | 1650    | 1316 |
| NO <sub>2</sub> asymmetric stretching vibration | 1541 | 1591    |      |
| C-C stretching aromatic vibration               |      | 1487    | 1556 |
| NO <sub>2</sub> symmetric stretching vibration  | 1346 | 1340    |      |
| C-N stretching vibration                        | 1188 | 1047    | 1313 |
| C=N stretching aromatic amine                   |      | 1629    | 1616 |

#### 4.2. Outcomes of the theoretical calculations

#### 4.2.1. Hirshfeld surface (HS) analysis

Hirshfeld surface (HS) studies were carried out in order to acquire a better knowledge of the packing motifs and inputs of the key intermolecular interactions that determine the molecular design of substances. **Fig.9a** depicts HS projected in two orientations on the  $d_{\text{norm}}$  property. The Crystal Explorer 3.1 software program was utilized to assess the strength and relevance of hydrogen bonds along with additional intermolecular interactions, including their effect on crystal form, 2D fingerprints (FP), and the relevant HS [71]. It was conducted according with the standards outlined in the literature and the instances presented [72], demonstrating the existence of the interactions identified by Mercury software [53] and illustrated in **Tables 1**. The colour scale is utilized to differentiate the HS's various intermolecular interaction areas. The HS analyses for NTO: ACR are shown in **Fig. 9a**, which show surfaces that have been mapped over  $d_{\text{norm}}$  (-0.7844 to 1.4763 Å). The consequences of the details are shown in **Fig. 9a** are plainly visible in these locations, with considerable circular depressions (deep red) visible on the surface, demonstrating  $\text{N}_4\text{-H}_{4\text{A}}\dots\text{O}_3$  and  $\text{N}_5\text{-H}_5\dots\text{O}_3$  intermolecular interactions. The red colour dots in the FP plots represent short contacts for the  $\text{O}\dots\text{H}$ ,  $\text{H}\dots\text{H}$ ,  $\text{N}\dots\text{H}$ ,  $\text{C}\dots\text{H}$ ,  $\text{O}\dots\text{O}$ , and  $\text{C}\dots\text{C}$  interactions (**Fig. 10**), emphasizing their importance in the molecular packing between NTO and ACR. The FP graphs in **Fig. 9b** show two separate "tails" for  $\text{N-H}\dots\text{O}$ , hydrogen bonds extending down to approximately  $d_i=0.75$  Å,  $d_e=1.16$  Å for NTO: ACR. The 30%  $\text{O}\dots\text{H}$ , 24%  $\text{H}\dots\text{H}$ , and 14%  $\text{N}\dots\text{H}$  interactions remain the most significant contributors to total HS and can be seen as the "ridge" of the FP. The arrangement of red-blue triangles in **Fig. 11b** is typical of  $\pi\dots\pi$  stacking (ACR...ACR), showing in the FP as a characteristic value approximately  $d_e = d_i = 1.8$  Å [73]. Moderate hydrogen bonds ( $\text{N-H}\dots\text{O}$ ) were some of the most critical intermolecular interactions, as seen in **Figs.4a-b**; this is dependable with the HS calculations and crystal structure data.



**Fig. 9.** a) HS mapped with  $d_{\text{norm}}$  b) FP plots c) Populations of inter-molecular and interatomic contacts of NTO: ACR.




Fig. 10. FP plots for NTO:ACR, displaying contributions from various contacts.




Fig. 11. a) curvedness, b) shape index and c)  $d_e$ .

#### 4.2.2. Theoretical calculation of heat of formation (HOF)

HOF is a significant measure for evaluating the explosive performance of EMs and can be determined using Eq. (1) [74–76]. where  $\Delta H_f^\circ(g, M)$  symbolizes the enthalpy of formation of the investigated molecule (M) in gas phase;  $H_{(molecule)}^\circ$  represents the calculated enthalpy value of the molecule M with the CBS-4M method ( $H^{298}$  in **Table 3**);  $\Sigma H_{(atoms)}^\circ$  symbolizes of calculated enthalpies for the separate atoms (H, C, N, O) and  $\Sigma \Delta H_f^\circ(atoms)$  stands for experimental values of the enthalpy of formation ( $\Delta H_f^\circ$ ) for the corresponding atoms H, C, N, and O to be 217, 716, 472, and 249  $\text{kJ}\cdot\text{mol}^{-1}$ , respectively [77].

$$\Delta H_f^\circ(\mathbf{g}, \mathbf{M}) = H_{(molecule)}^\circ - \Sigma H_{(atoms)}^\circ + \Sigma \Delta H_f^\circ(atoms) \quad 1$$

**Table 3.** Gas-phase enthalpy of formation results

| Entry    | P.G. <sup>1</sup> | NIMAG <sup>2</sup> | -H <sup>298</sup><br>(a. u.) <sup>3</sup> | $\Delta H_f^\circ(\mathbf{g}, \mathbf{M})$<br>(kJ.mol <sup>-1</sup> ) <sup>4</sup> |
|----------|-------------------|--------------------|-------------------------------------------|------------------------------------------------------------------------------------|
| NTO      | C <sub>i</sub>    | 0                  | 519                                       | 38.00                                                                              |
| NTO: ACR | C <sub>i</sub>    | 0                  | 1069                                      | 168.90                                                                             |

<sup>1</sup>Point group, <sup>2</sup>Number of imaginary frequencies.

<sup>3</sup>Enthalpy values obtained from the CBS-4M calculations,

<sup>4</sup>Enthalpy of formation (in gas phase).

Furthermore, it is widely accepted that, to precisely examine the detonation properties of EMs, the calculation of the solid-phase heat of formation  $\Delta H_f^\circ(\mathbf{s}, \mathbf{M})$  is necessary, and this is easily performed done by applying Hess's law (Eq. 2) [78].

$$\Delta H_f^\circ(\mathbf{s}, \mathbf{M}) = \Delta H_f^\circ(\mathbf{g}, \mathbf{M}) - \Delta H_f^\circ(\mathbf{sub}, \mathbf{M}) \quad 2$$

where  $\Delta H_f^\circ(\mathbf{sub}, \mathbf{M})$  indicates the heat of sublimation, which may be determined through the Rice and Politzer equation (Eq. 3) [79].

$$\Delta H_f^\circ(\mathbf{sub}, \mathbf{M}) = aA^2 + b(\nu\sigma_{tot}^2)^{0.5} + c \quad 3$$

where A denotes the surface area of the 0.001 electron/Bohr<sup>3</sup> iso-surface of the electronic density,  $\nu$  represents the degree of balance between positive and negative potentials on the isosurface, and finally  $\sigma_{tot}^2$  is an index of evaluating the variability of the electrostatic potential on the molecular surface (Table 4). A,  $\nu$  and  $\sigma_{tot}^2$  values were calculated by using Multiwfn program (v.3.8 – submenu 12: quantitative analysis of molecular surface) [80], other coefficients ( $a= 2.670 \times 10^{-4}$  kcal.mol<sup>-1</sup>,  $b=1.650$  kcal.mol<sup>-1</sup>, and  $c=2.966$  kcal.mol<sup>-1</sup>) were obtained from the literature [81].

**Table 4.** Calculated  $\Delta H_f^\circ(\mathbf{sub}, \mathbf{M})$ ,  $\Delta H_f^\circ(\mathbf{s}, \mathbf{M})$ , and  $\Delta H_f^\circ(\mathbf{g}, \mathbf{M})$  results of NTO and NTO: ACR


| Entry   | gas phase                                  | solid phase                                 |                                              |
|---------|--------------------------------------------|---------------------------------------------|----------------------------------------------|
|         | $\Delta H_f^\circ(\mathbf{g}, \mathbf{M})$ | $-\Delta H_f^\circ(\mathbf{s}, \mathbf{M})$ | $\Delta H_f^\circ(\mathbf{sub}, \mathbf{M})$ |
| NTO     | 38                                         | 58                                          | 96                                           |
| NTO:ACR | 168                                        | 37                                          | 206                                          |

#### 4.2.3. Electrostatic potential (ESP) maps

ESP is essentially significant physical property of compounds since it offers information on charge density distribution and chemical reactivity. In accordance with Fig. 12, positive ESP (red regions) is mainly dispersed across the main skeleton, while negative ESP (blue areas)



focuses on the molecules' borders, particularly around the nitrogen and oxygen atoms, due to their higher electronegativity. The entirety of the following alterations points to variances in NTO: ACR charge dispersion may result in changes in sensitivity.



**Fig.12.** ESP of NTO, ACR, and NTO: ACR

#### 4.2.4. Non-covalent interaction (NCI) plots

In order to show the intermolecular interactions of the NTO: ACR in real space, a reduced density gradient (RDG) study utilizing electron density was performed. RDG can identify non-covalent interactions (NCIs), which can provide important information on the source of molecular change, crystal packing, and stability [82, 83]. Erin et al. suggested the following relationship between the functions RDG and  $\rho(r)$ : [84].

$$RDG = \frac{1}{2 (3 \pi^2)^{1/3}} \frac{|\nabla \rho(r)|}{\rho(r)^{4/3}} \quad 4$$

$$(\Omega) = \text{sign}(\lambda_2(r)) \rho(r) \quad 5$$

where  $\rho(r)$  is the electron density,  $\text{sign}(\lambda_2(r))$  is the sign of the second largest Hessian eigenvalue. The spikes are classified into three groups in RDG maps: the hydrogen bond area, the vdW interaction area, and the steric area. Initially, we investigate a case with  $\rho$  ranging from -0.04 to -0.03 a.u., whose occurs within the hydrogen bond area. (**Figs. 13a-b**). It is obvious that NTO: ACR salt is formed through intermolecular interactions including  $\text{N}_4\text{-H}_{4\text{A}}\dots\text{O}_3$  and  $\text{N}_5\text{-H}_5\dots\text{O}_3$ . Following that, in the vdW interaction region with values ranging from -0.01 to 0.005 a.u., it is shown that NTO: ACR exhibits greater spikes, confirming significant vdW interactions (**Figs. 13a-b**). As shown in **Figs.13a-b**, the scatter points and electron density of NTO: ACR in the steric area (varying from 0.01 to 0.05 a.u) has sharper spikes and more points, confirming significant strong repulsion interactions. It is straightforward to deduce that moderate hydrogen bonds, vdW forces, and steric effects account for a significant portion of NTO:ACR intermolecular interactions.






Fig. 13. RDG map and non-covalent interaction plots of low-gradient iso-surfaces for NTO: ACR.

### 4.3. Detonation Parameters for NTO: ACR

#### 4.3.1. Impact sensitivity

Impact sensitivity was executed on NTO and NTO:ACR. NTO:ACR are insensitive, with no signs of initiation up to the testing limit of 100 J (**Table 5**). NTO:ACR is substantially less impact sensitive than NTO, and this is attributed to the layered structure of NTO:ACR. NTO:ACR has layered structures akin to TATB (triaminotrinitrobenzene), one of the most insensitive traditional EMs.

#### 4.3.2. Free space in the crystal lattice and packing coefficient

By applying Mercury CSD Ver. 4.1, the free space, i.e., voids, and packing coefficient of NTO and NTO: ACR were calculated [53]. As a consequence of the formation of NTO: ACR, the packing coefficient altered. Despite NTO: ACR is not dense than NTO, the new compound's making drastically altered the packing coefficient and voids (**Table 5**) **Figs. S11-S12**. Furthermore, higher compactness indicates less free volume in the crystal, causing molecular breakdown and the occurrence of hotspots, resulting in greater insensitivity [69, 85, 86].

**Table 5.** Packing coefficient and free space of NTO, and NTO: ACR.

| Entry    | Packing Coefficient (%) | Free space (voids) (%) | Impact sensitivity (J) |
|----------|-------------------------|------------------------|------------------------|
| NTO      | 74.13                   | 5.9                    | 100                    |
| NTO: ACR | 69.04                   | 12.7                   | 100                    |

#### 4.3.3. Calculation of detonation velocity, pressure, and oxygen balance

The characteristics of detonation are crucial for acquiring a fundamental comprehension of EMs performance. The oxygen balance, detonation velocity and pressure were calculated using EXPLO5 V6.04. It uses the Becker-Kistiakowsky-Wilson's equation of state for gaseous detonation products and Cowan-Fickett's equation of state for solid carbon [56]. The

detonation parameters were calculated at the C-J point. The primary detonation products were thought to consist of C, N<sub>2</sub>, H<sub>2</sub>O, CH<sub>4</sub>, NH<sub>3</sub>, CO<sub>2</sub>, C<sub>2</sub>H<sub>6</sub>, H<sub>2</sub>, CO, CH<sub>2</sub>O<sub>2</sub>, C<sub>2</sub>H<sub>4</sub>, HCN, CH<sub>3</sub>OH (**Figs. S13-S15.**) Calculated heat of formation ( $\Delta H^\circ_f$  (g, M),  $\Delta H^\circ_f$  (s, M)), and detonation parameters are listed in **Table 6**.

**Table 6.** Calculated detonation properties of NTO, and NTO: ACR.

| Entry    | $\Omega$ [%] <sup>a</sup> | IS [J] <sup>b</sup> | Gas phase                          |                     | Solid phase                         |                     |
|----------|---------------------------|---------------------|------------------------------------|---------------------|-------------------------------------|---------------------|
|          |                           |                     | D[m.s <sup>-1</sup> ] <sup>c</sup> | P[GPa] <sup>c</sup> | D [m.s <sup>-1</sup> ] <sup>c</sup> | P[GPa] <sup>d</sup> |
| NTO      | -24.6                     | 100                 | 8747                               | 33.01               | 7200                                | 21.04               |
| NTO: ACR | -168.1                    | 100                 | 7006                               | 20.02               | 5480                                | 8.75                |

<sup>a</sup>Oxygen balance, <sup>c</sup>Detonation velocity, and <sup>d</sup>Detonation pressure was calculated using EXPLO5 V6.04

<sup>b</sup>Impact sensitivity was measured by BAM hammer instrument

## 5. Conclusion

The co-crystallization method was utilized to form a novel energetic compound 3-nitro-1,2,4-triazol-5-one (NTO): acridine (ACR), which was then comprehensively characterized by IR spectroscopy, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR spectroscopy, mass spectrometry, and DSC study. The structure attributes of NTO: ACR were investigated using SCXRD and PCXRD. Through changing the chemical composition, it was possible to alter physical properties such as thermal stability, free space (voids), packing coefficient, crystal density, difference in pKa of co-formers, morphology, solubility, impact sensitivity, and calculated detonation parameters were also gathered and compared. Appears it was possible to completely modify physical properties. The crystal structure and crystal packing features of NTO: ACR demonstrate that the anions and cations in this energetic compound have modest hydrogen-bond intermolecular interactions (N<sub>4</sub>-H<sub>4A</sub>...O<sub>3</sub> and N<sub>5</sub>-H<sub>5</sub>...O<sub>3</sub>). Furthermore, Hirshfeld surface analysis and non-covalent interaction studies gave theoretical support to better investigate its structure-property relationship. The detonation velocity and pressure of NTO: ACR were calculated with the assistance of the EXPLO5 program and found to be 7006 ms<sup>-1</sup> and 20.02 G Pa, respectively, slightly less than those of NTO. Although non-energetic components usually decrease the detonation velocity and pressure, they can significantly enhance the number of energetic co-crystals and salts, enabling a rich database for subsequent investigations in a data-driven or crystal engineering approach.

## Acknowledgments

The authors would like to thank Dr Emily Arnold from Cranfield University for providing the data from the power X-Ray diffraction experiments.

## Author contribution

All authors contributed to the study conception and design. Nilgün Şen: Conceptualisation, Investigation, Methodology, Data curation, Formal Analysis, Writing-Editing, Funding acquisition. Jean-François Pons: Investigation, Methodology, Data curation, Formal Analysis, Writing-review and editing. Yunus Zorlu: Data curation, Visualization. Eleftheria Dossi:

Investigation, Methodology, Data curation, Formal Analysis, Writing-review and editing. Federica Persico: Formal Analysis, Writing-review and editing. Tracey Temple: Formal Analysis, Writing-review and editing, Nazife Aslan: Investigation, Methodology, Data curation, Akachai Khumsri: Data curation, Formal Analysis, Validation, Software. All authors read and approved the final manuscript.

### **Funding**

This study was funded by the Scientific and Technological Research Council of Türkiye-TUBITAK-BIDEB 2219-International Postdoctoral Research Fellowship Program (Grant number:1059B192100006).

### **Data availability**

The corresponding CIF files are available in the Cambridge Structural Database (CSD); for 1,2,4-triazol-5-one (NTO): Acridine, the CCDC number is 2253569.

### **Code availability**

Not applicable.

### **Declarations**

Competing interests, the authors declare no competing interests.

### **References**

- [1] Thottempudi V, Yin P, Zhang J, Parrish DA, Shreeve JM (2014) 1,2,3-Triazolo[4,5-e]furazano[3,4,-b]pyrazine 6-Oxide—A Fused heterocycle with a roving hydrogen forms a new class of insensitive energetic materials. *Chem Eur J* 20:542-548. <https://doi.org/10.1002/chem.201303469>
- [2] Gao H, Joo YH, Twamley B, Zhou Z (2011) Azole-based energetic salts. *Chem Rev* 111 (11):7377-7436. <https://doi.org/10.1021/cr200039c>
- [3] Dippold AA, Izsák D, Klapötke TM (2013) A Study of 5-(1, 2, 4-triazol-c-yl) tetrazol-1-ols: combining the benefits of different heterocycles for the design of energetic materials. *Chem Eur J* 19:12042-12051. <https://doi.org/10.1002/chem.201301339>
- [4] Millar RW, Hamid J, Endsor R, Swinton PF, Cooper J (2008) Selection and synthesis of energetic heterocyclic compounds suitable for use in insensitive explosive and propellant compositions. *Propel Explos Pyrotech* 33:66-72. <https://doi.org/10.1002/prop.200800211>
- [5] Badgujar DM, Talawar MB, Asthana SN, Mahulikar PP (2008) Advances in science and technology of modern energetic materials: an overview. *J Hazard Mater* 151:289-305. <https://doi.org/10.1016/j.jhazmat.2007.10.039>
- [6] Fischer D, Klapötke TM, Stierstorfer J (2014) Oxalylhydrazinium nitrate and dinitrate—efficiency meets performance. *J Energ Mater* 32:37–49. <https://doi.org/10.1080/07370652.2012.750697>
- [7] Hanafi S, Trache D, Abdous S, Bensalem Z, Mezroua A (2019) 5-Nitro-1, 2, 4-triazole-3-one: A review of recent advances. *Chin J Ener Mater* 27:326-347.

<https://doi.org/10.11943/CJEM2018371>

- [8] Pagoria PF, Lee GS, Mitchell AR, Schmidt RD (2002) A review of energetic materials synthesis. *Thermochim Acta* 384:187–204. [https://doi.org/10.1016/S0040-6031\(01\)00805-X](https://doi.org/10.1016/S0040-6031(01)00805-X)
- [9] Sabatini JJ, Oyler KD (2016) Recent advances in the synthesis of high explosive materials. *Crystals* 6:1-22. <https://doi.org/10.3390/cryst6010005>
- [10] Chen J, Tang J, Xiong H, Yang H, Cheng G (2020) Combining triazole and furazan frameworks via methylene bridges for new insensitive energetic materials. *Ener Mater Front* 1:34-39. <https://doi.org/10.1016/j.enmf.2020.07.001>
- [11] Zheng Y, Zhao X, Qi X, Wang K, Liu T (2020) Synthesis of 5-(1H-pyrazol-1-yl)-2H-tetrazole-derived energetic salts with high thermal stability and low sensitivity. *Ener Mater Front* 1:83–89. <https://doi.org/10.1016/j.enmf.2020.08.004>
- [12] Yao W, Xue Y, Qian L, Yang H, Cheng G (2021) Combination of 1,2,3-triazole and 1,2,4-triazole frameworks for new high-energy and low-sensitivity compounds. *Ener Mater Front* 2. <https://doi.org/10.1016/j.enmf.2021.05.002>
- [13] Bu R, Jiao F, Liu G, Zhao J, Zhang C (2021) Categorizing and understanding energetic crystals. *Cryst Growth Des* 21:3–15. <https://doi.org/10.1021/acs.cgd.0c01300>
- [14] Zhang C (2018) On the energy safety contradiction of energetic materials and the strategy for developing low-sensitive high-energetic materials. *Chin J Energy Mater* 26:2–10. <https://doi.org/10.11943/j.issn.1006-9941.2018.01.001>
- [15] Zhang C, Jiao F, Li H (2018) Crystal engineering for creating low sensitivity and highly energetic materials. *Cryst Growth Des* 18:5713–5726. <https://doi.org/10.1021/acs.cgd.8b00929>
- [16] Jiao F, Xiong Y, Li H, Zhang C (2018) Alleviating the energy & safety contradiction to construct new low sensitivity and highly energetic materials through crystal engineering. *Cryst Eng Comm* 20:1757–1768. <https://doi.org/10.1039/C7CE01993A>
- [17] Teipel U (2005) *Energetic materials: In particle processing and characterization*. Wiley-VCH
- [18] Fawcett-Hirst W, Temple TJ, Ladyman MK, Coulon F (2021) A review of treatment methods for insensitive high explosive contaminated wastewater. *Heliyon* 7:e07438. <https://doi.org/10.1016/j.heliyon.2021.e07438>
- [19] Wu JT, Zhang JG, Li T, Li ZM, Zhang TL (2015) A novel cocrystal explosive NTO/TZTN with good comprehensive properties. *Rsc Advances* 5:28354-28359. <https://doi.org/10.1039/C5RA01124H>
- [20] Migliaro FW, Rothgery EF (1993) Hydroxylammonium salts of 5-nitro-1,2,4-triazol-3-one. US Pat. no. 5274105
- [21] Botcher TR, Beardall DJ, Wight CA, Fan L, Burkey TJ (1996) Thermal decomposition mechanism of NTO. *J Phys Chem* 100:8802–8806. <https://doi.org/10.1021/jp952984y>
- [22] Zhao Y, Chen S, Jin S, Li Z, Zhang X, Wang L, Mao Y, Guo H, Li L (2017) Heat effects

- of NTO synthesis in nitric acid solution. *J Therm Anal Calorim* 128:301–310. <https://doi.org/10.1007/s10973-016-5912-x>
- [23] Pourmortazavi SM, Rahimi-Nasrabadi M, Kohsari I, Hajimirsadeghi SS (2012) Non-isothermal kinetic studies on thermal decomposition of energetic materials : KNF and NTO. *J Therm Anal Calorim* 110:857–863. <https://doi.org/10.1007/s10973-011-1845-6>
  - [24] Sweeney LM, Phillips EA, Goodwin MR, Bannon DI (2015) Toxicokinetic model development for the insensitive munitions component 3-nitro-1,2,4-triazol-5-one. *Int J Toxicol* 34:408–416. <https://doi.org/10.1177/1091581815589000>
  - [25] 3-Nitro-1,2,4-triazol-5-one (2014). (2018)*Toxicology and Industrial Health* 34 (1):8–14. <https://doi.org/10.1177/0748233717719688>.
  - [26] Mark N, Arthur J, Dontsova K, Brusseau M, Taylor S, Šimůnek J (2017) Column transport studies of 3-nitro-1,2,4-triazol-5-one (NTO) in soils. *Chemosp* 171:427–434. <https://doi.org/10.1016/j.chemosphere.2016.12.067>
  - [27] Şen N, Dursun H, Hope KS, Nazir H, Acar N, Atakol O (2020) Towards low-impact-sensitivity through crystal engineering: New energetic co-crystals formed between picric acid, trinitrotoluene and 9-vinyanthracene. *J Mol Struct* 1219. <https://doi.org/10.1016/j.molstruc.2020.128614>
  - [28] Haley MV, Kuperman RG, Checkai RT (2009) Aquatic toxicity of 3-nitro-1,2,4-triazol-5-one. In: Report: ECBC-TR-726. US Army Center for Health Promotion and Preventative Medicine, Directorate of Toxicology
  - [29] Quinn MJ, Bannon DI, Jackovitz AM, Hanna TL, Shiflett AA, Johnson MS (2014) Assessment of 3-nitro-1,2,4-triazol-5-one as a potential endocrine disrupting chemical in rats using the hershberger and uterotrophic bioassays. *Int J Tox* 33:367–372. <https://doi.org/10.1177/1091581814548729>
  - [30] O'Bryan TR, Ross RH (1988) Chemical scoring system for hazard and exposure identification. *J Toxicol Environ Health* 25:119–134. <https://doi.org/10.1080/15287398809531193>
  - [31] Tennant M, Chew SC, Krämer T, Mai N, McAteer D, Pons JF (2019) Practical colorimetry of 3-nitro-1,2,4-triazol-5-one. *Propel Explos Pyrotech* 44:198–202. <https://doi.org/10.1002/prop.201800239>
  - [32] Shen JP, Duan XH, Luo QP, Zhou Y, Bao Q, Ma YJ, Pei CH (2011) Preparation and characterization of a novel cocrystal explosive. *Cryst Growth Des* 11:1759–1765. <https://doi.org/10.1021/cg1017032>
  - [33] Remenar JF, Morissette SL, Peterson ML, Moulton B, MacPhee JM, Guzmán HR, Almarsson O (2003) Crystal engineering of novel cocrystals of a triazole drug with 1, 4-dicarboxylic acids. *J Am Chem Soc* 125:8456–8457. <https://doi.org/10.1021/ja035776p>
  - [34] Weyna DR, Shattock T, Vishweshwar P, Zaworotko MJ (2009) Synthesis and structural characterization of cocrystals and pharmaceutical cocrystals: mechanochemistry vs slow

- evaporation from solution. *Cryst Growth Des* 9:1106–1123. <https://doi.org/10.1021/cg800936d>
- [35] Almarsson Ö, Zaworotko MJ (2004) Crystal engineering of the composition of pharmaceutical phases. Do pharmaceutical co-crystals represent a new path to improved medicines?. *Chem Commun* 1889–1896. <https://doi.org/10.1039/B402150A>
- [36] Çelik GG, Dedeoglu B, Gürek AG, Zorlu Y, Ayhan MM (2023) Enhancing supramolecular assembly in bodipy derivatives: harnessing halogen bonding for cocrystal design. *Cryst Growth Des* 23:7285–7294. <https://doi.org/10.1021/acs.cgd.3c00663>
- [37] Schütrumpf A, Kirpi E, Bulut A, Morel FL, Ranocchiari M, Lork E, Zorlu Y, Grabowsky S, Yücesan G, Beckmann J (2015) Tetrahedral tetraphosphonic acids. new building blocks in supramolecular chemistry. *Cryst Growth Des* 15:4925–4931. <https://doi.org/10.1021/acs.cgd.5b00811>
- [38] Alshahateet SF (2011) Synthesis and X-ray crystallographic analysis of pharmaceutical model rac-ibuprofen cocrystal. *J Chem Crystallogr* 41:276–279. <https://doi.org/10.1007/s10870-010-9872-x>
- [39] Zhu W, Xiao J, Zhao F, Ji G, Ma X, Xiao H (2007) Molecular Dynamics Simulation of Elastic Properties of HMX/TATB. *Acta Chim Sin* 65:1223–1228
- [40] Wei C, Huang H, Duan X, Pei C (2011) Structures and Properties Prediction of HMX/TATB Co-Crystal. *Propel Explos Pyrotech* 36:416–423. <https://doi.org/10.1002/prop.201000022>
- [41] Landenberger KB, Matzger AJ (2012) Cocrystals of 1, 3, 5, 7-Tetranitro-1, 3, 5, 7-tetrazacyclooctane (HMX). *Cryst Growth Des* 12:3603–3609. <https://doi.org/10.1021/cg3004245>
- [42] Sun T, Xiao JJ, Liu Q, Zhao F, Xiao HM (2014) A Comparative study on structure, energetic and mechanical properties of a  $\epsilon$ -CL-20/HMX cocrystal and its composite with molecular dynamics simulation. *J Mater Chem A* 2:13898–13904. <https://doi.org/10.1039/C4TA01150C>
- [43] Li JC, Jiao QJ, Gong YG, Wang YY, Liang T, Sun J (2018) Explosive performance of HMX/NTO co-crystal. *IOP Conference Series: Materials Science and Engineering* 292:012032. <https://doi.org/10.1088/1757-899X/292/1/012032>
- [44] Hope KS, Lloyd HJ, Ward D, Michalchuk AA, Pulham CR (2015) Resonant acoustic mixing and its applications to energetic materials. *New Trends in Research of Energetic Materials*, University of Pardubice, Pardubice, Czech Republic
- [45] Lloyd HJ (2017) Co-crystallisation of energetic materials-a step-change in the control of properties and performance of munitions. Dissertation of doctorate. University of Edinburgh
- [46] Ren YH, Zhao FQ, Yi JH, Xu KZ, Ma HX, Hu RZ, Song JR (2012) Studies on an ionic compound  $(3\text{-ATz})^+(\text{NTO})^-$ : crystal structure, specific heat capacity, thermal behaviors and thermal safety. *J Iran Chem Soc* 9:407–414. <https://doi.org/10.1007/s13738-011-0051-6>

- [47] Ma H, Song J, Hu R, Zhai G, Xu K, Wen Z (2004) Molecular structure, the quantum chemical investigation and the thermal behavior of the dimethylamine salt of 3-nitro-1,2,4-triazol-5-one,  $(\text{CH}_3)_2\text{NH}_2^+\text{C}_2\text{N}_4\text{O}_3\text{H}^-$ . *J Mol Struct: THEOCHEM* 678:217-222. <https://doi.org/10.1016/j.theochem.2004.03.018>
- [48] AAT Bioquest, Inc. (2023, November 7). Quest Database™ pKa and pKb Reference Table. AAT Bioquest. <https://www.aatbio.com/data-sets/pka-and-pkb-reference-table>
- [49] Ledgard JB (2007) *The Preparatory Manual of Explosives*, 3<sup>rd</sup> edn. Lulu Publishing, Morrisville, Washington, USA
- [50] Sheldrick GM (2015) SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr Sect A Found Adv* 71:3–8. <https://doi.org/10.1107/S2053273314026370>
- [51] Dolomanov OV, Bourhis LJ, Gildea RJ, Howard JAK, Puschmann H (2009) OLEX2: a complete structure solution, refinement and analysis program. *J Appl Crystallogr* 42:339–341. <https://doi.org/10.1107/S0021889808042726>
- [52] Sheldrick GM (2015) Crystal structure refinement with SHELXL. *Acta Crystallogr Sect C Struct Chem* 71:3–8. <https://doi.org/10.1107/S2053229614024218>
- [53] Mercury CSD 4.1.0 (2022) (Build 235316). <http://www.ccdc.cam.ac.uk/mercury/>
- [54] Kosanke KL, Kosanke BJ, Sturman BT, Winokur RM (2012) *Encyclopedic dictionary of pyrotechnics*, 1<sup>st</sup> edn. J Pyrotechnics, USA
- [55] Fuh C, Lee J, Liaw C (2003) The design aspect of the Bruceton test for pyrotechnics sensitivity analysis. *J Data Sci* 1:83–101.
- [56] Sucéska M, Sucéska M (2020) EXPLO5, Version 6.01; Brodarski Institute, Zagreb, Croatia, EXPLO5 Version 6.04
- [57] Spackman MA, Turner MJ, McKinnon JJ, Wolff SK, Grimwood DJ, Jayatilaka D, Spackman PR (2017) Crystal Explorer 17. Univ West Aust <http://hirshfeldsurface.net>
- [58] Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Petersson GA, Nakatsuji H, Li X, Caricato M, Marenich A, Bloino J, Janesko BG, Gomperts R, Mennucci B, Hratchian HP, Ortiz JV, Izmaylov AF, Sonnenberg JL, Williams-Young D, Ding F, Lipparini F, Egidi F, Goings J, Peng B, Petrone A, Henderson T, Ranasinghe D, Zakrzewski VG, Gao J, Rega N, Zheng G, Liang W, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven T, Throssell K, Montgomery Jr. JA, Peralta JE, Ogliaro F, Bearpark M, Heyd JJ, Brothers E, Kudin KN, Staroverov VN, Keith T, Kobayashi R, Normand J, Raghavachari K, Rendell A, Burant JC, Iyengar SS, Tomasi J, Cossi M, Millam JM, Klene M, Adamo C, Cammi R, Ochterski JW, Martin RL, Morokuma K, Farkas O, Foresman JB, Fox DJ (2016) *Gaussian 09*. Rev. D.01, Gaussian Inc., Wallingford, CT
- [59] Dennington R, Keith T, Millam J (2009) *GaussView*. Version 5.0.8, GaussView, Version 5.0.8.



- [60] Jomaa L, Issaoui N, Roisnel T, Marouani H (2021) Insight into non-covalent interactions in a tetrachlorocadmate salt with promising NLO properties: experimental and computational analysis. *J Mol Struct* 1242:130730. <https://doi.org/10.1016/j.molstruc.2021.130730>
- [61] Tian L, Xu Y, Lin Q, Wang P, Lu M (2019) Syntheses of energetic cyclo -pentazolate salts. *Chem Asian J* 14:2877 – 2882. <https://doi.org/10.1002/asia.201900776>
- [62] Lu T, Chen F (2012) Multiwfn: A multifunctional wavefunction analyzer. *J Comput Chem* 33:580–592. <https://doi.org/10.1002/jcc.22885>
- [63] Humphrey W, Dalke A, Schulten K (1996) VMD: Visual molecular dynamics. *J Mol Graph* 14:33–38. [https://doi.org/10.1016/0263-7855\(96\)00018-5](https://doi.org/10.1016/0263-7855(96)00018-5)
- [64] Cruz Cabeza AJ (2012) Acid-base crystalline complexes and the pKa rule. *Cryst Eng Comm* 14:6362–6365, <https://doi.org/10.1039/C2CE26055G>
- [65] Steiner T (2002) The hydrogen bond in the solid state. *Angew Chem Int Ed* 41:48-76. [https://doi.org/10.1002/1521-3773\(20020104\)41:1<48::AID-ANIE48>3.0.CO;2-U](https://doi.org/10.1002/1521-3773(20020104)41:1<48::AID-ANIE48>3.0.CO;2-U)
- [66] Vereshchagin AL, Morozova EA (2022) Hydrogen bonding in insensitive high explosives. *Russ J Appl Chem* 95:1069–1084. <https://doi.org/10.1134/S1070427222080018>
- [67] Freedman HH (1961) Intramolecular H-bonds. I. A spectroscopic study of the hydrogen bond between hydroxyl and nitrogen. *J Am Chem Soc* 83:2900-2905
- [68] Klapotke TM, Mieskes F, Stierstorfer J, Weyrauther M (2016) Studies on energetic salts based on (2,4,6-trinitrophenyl) guanidine. *Propel Explos Pyrotech* 41:217-222. <https://doi.org/10.1002/prop.201500338>
- [69] Politzer P, Murray JS (2016) High performance, low sensitivity: conflicting or compatible?. *Propel Explos Pyrotech* 41:414–425, doi:10.1002/prop.201500349
- [70] Wu Q, Pan Y, Xia X, Shao Y, Zhu W, Xiao H (2013) Theoretic design of 1,2,3,4-tetrazine-1,3-dioxide-based high-energy density compounds with oxygen balance close to zero. *Struct Chem* 24:1579–1590. doi:10.1007/s11224-012-0190-0
- [71] Spackman PR, Turner MJ, McKinnon JJ, Wolff SK, Grimwood DJ, Jayatilaka D, Spackman MA (2021) Crystal explorer: A program for hirshfeld surface analysis, visualization and quantitative analysis of molecular crystals. *J Appl Cryst* 54:1006-1011, <https://doi.org/10.1107/S1600576721002910>
- [72] Panini P, Mohan TP, Gangwar U, Sankolli R, Chopra D (2013) Quantitative crystal structure analysis of 1,3,4-thiadiazole derivatives. *Cryst Eng Comm* 15:4549-4564. <https://doi.org/10.1039/C3CE40278A>
- [73] Bondi A (1964) Van der Waals volumes and radii. *J Phys Chem* 68:441–447. <https://pubs.acs.org/doi/10.1021/j100785a001P>
- [74] Curtiss LA, Raghavachari K, Redfern PC, Pople JA (1997) Assessment of gaussian-2 and density functional theories for the computation of enthalpies of formation. *J Chem Phys* 106:1063-1079. <https://doi.org/10.1063/1.473182>

- [75] Byrd EFC, Rice BM (2009) Improved prediction of heats of formation of energetic materials using quantum mechanical calculations. *J Phys Chem A* 113:5813-5813. <https://doi:10.1021/jp806520b>
- [76] Rice BM, Pai SV, Hare J (1999) Predicting heats of formation of energetic materials quantum mechanical calculations. *J Com Flame* 118:445-458. [https://doi:10.1016/S0010-2180\(99\)00008-5](https://doi:10.1016/S0010-2180(99)00008-5)
- [77] Johnson III RD (2015) NIST computational chemistry comparison and benchmark data base (Release 17b): NIST Standard Reference Database Number 101
- [78] Byrd EFC, Rice BM (2009) A comparison of methods to predict solid phase heats of formation of molecular energetic salts. *J Phys Chem A* 113:345-352. <https://doi.org/10.1021/jp807822e>
- [79] Politzer P, Murray JS (2011) Some perspectives on estimating detonation properties of C, H, N, O compounds. *Cen Eur J Energ Mat* 8:209-220
- [80] Jin XH, Hu BC, Jia HQ, Liu ZL, Lu CX (2014) DFT studies on two novel explosives based on the guanidine-fused bicyclic structure. *Bull Korean Chem Soc* 35:1043–1049. [doi:10.5012/bkcs.2014.35.4.1043](https://doi.org/10.5012/bkcs.2014.35.4.1043)
- [81] Xie Z, Hu S, Cao X (2016) Theoretical insight into the influence of molecular ratio on the binding energy and mechanical property of HMX/2-picoline-N-oxide co-crystal, cooperativity effect and surface electrostatic potential. *Mol Phys* 114:2164–2176. <https://doi.org/10.1080/00268976.2016.1190038>
- [82] Lin H, Zhu SG, Li HZ, Peng XH (2013) Structure and detonation performance of a novel HMX/LLM-105 cocrystal explosive. *J Phys Org Chem* 26:898–907. <https://doi.org/10.1002/poc.3188>
- [83] Chen T, Zhang Y, Guo SF, Zhao LM, Chen W, GZ Hao, Xiao L, Ke X, Jiang W (2019) Preparation and property of CL-20/BAMO-THF energetic nanocomposites. *Def Tech* 15:306–312. <https://doi.org/10.1016/J.DT.2018.08.013>
- [84] Johnson ER, Keinan S, Mori-Sánchez P, Contreras-García J, Cohen AJ, Yang W (2010) Revealing non-covalent interactions. *J Am Chem Soc* 132:6498. <https://doi.org/10.1021/JA100936W>
- [85] Xu Z, Yang H, Cheng G (2016) Studies on the synthesis and properties of polynitro compounds based on esteryl backbones. *New J Chem* 40:9936–9944. <https://doi.org/10.1039/C6NJ02198K>



**DSpace**

<https://dspace.org/>

## **Synthesis, structure characterization, Hirshfeld surface analysis, and computational studies of 3-nitro-1,2,4-triazol-5-one (NTO):acridine**

**en, Nilgün; Pons, Jean-François; Zorlu, Yunus; Dossi, Eleftheria; Persico, Federica ...**

**2024-12**

Springer

<https://doi.org/10.1007/s11224-024-02326-0>

en N, Pons JF, Zorlu Y, et al., (2024) Synthesis, structure characterization, Hirshfeld surface analysis, and computational studies of 3-nitro-1, 2, 4-triazol-5-one (NTO): acridine. Structural Chemistry, Volume 35, Issue 6, December 2024, pp. 1865–1879