

Small-scale indirect plant responses to insect herbivory could have major impacts on canopy photosynthesis and isoprene emission

Kristiina Visakorpi^{1,2}, Sofia Gripenberg¹, Yadvinder Malhi², Conor Bolas³, Imma Oliveras², Neil Harris⁴, Sami Rifai² and Terhi Riutta²

1.) Department of Zoology, University of Oxford, The John Krebs Field Station, Wytham, Oxford OX2 8QJ, UK.

2.) Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, OX1 3QY, UK.

3.) Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK

4.) Centre for Atmospheric Informatics and Emissions Technology, Cranfield University, Cranfield MK43 0AL, UK.

Corresponding author:

Kristiina Visakorpi

Email: kristiina.visakorpi@zoo.ox.ac.uk

Tel: +447468887679

Summary

- Insect herbivores cause substantial changes in the leaves they attack, but their effects on the ecophysiology of neighbouring, non-damaged leaves have never been quantified in natural canopies. We studied how winter moth (*Operophtera brumata*), a common herbivore in temperate forests, affects the photosynthetic and isoprene emission rates of its host plant, the pedunculate oak (*Quercus robur*).
- Through a manipulative experiment, we measured leaves on shoots damaged by caterpillars or mechanically by cutting, or left completely intact. To quantify the effects at the canopy scale, we surveyed the extent and patterns of leaf area loss in the canopy.
- Herbivory reduced photosynthesis both in damaged leaves and in their intact neighbours. Isoprene emission rates significantly increased after mechanical leaf damage. When scaled up to canopy-level, herbivory reduced photosynthesis by $48 \pm 10\%$.
- The indirect effects of herbivory on photosynthesis on undamaged leaves (40%) were much more important than the direct effects of leaf area loss (6%). If widespread across other plant-herbivore systems, these findings suggest that insect herbivory has major and previously underappreciated influences in modifying ecosystem carbon cycling, with potential effects on atmospheric chemistry.

Keywords: canopy, carbon cycling, herbivory, isoprene, photosynthesis, *Quercus robur*

Introduction

Interactions between plants and insect herbivores are among the most common ecological interactions (Strong *et al.*, 1984; Schoonhoven *et al.*, 2005). By influencing plant distribution, abundance and evolution, insect herbivores can have major impacts on community composition, primary productivity and biosphere–atmosphere interactions (Belovsky & Slade, 2000; Karl *et al.*, 2008; Metcalfe *et al.*, 2014).

By removing plant tissue (*a direct effect* of herbivory), insect herbivores can substantially reduce photosynthesis. The loss of tissue often changes both primary (basic metabolic processes like respiration) and secondary (e.g. production of defensive chemicals) plant metabolism (Herms & Mattson, 1992; Kerchev *et al.*, 2012). This can lead to changes in the nutrient content or toxicity of the plant. Plants can also respond to herbivory by emitting volatile organic compounds (“VOCs”, Rowen & Kaplan, 2016). These changes, often triggered as defensive reactions, can spread to systemic undamaged tissue and affect all parts of the plant (Agrawal, 2000; Staudt & Lhoutellier, 2007; Wu & Baldwin, 2009).

Insect-induced changes in chemistry and metabolism can further alter the photosynthetic capacity of the remaining leaf tissue (*an indirect effect* of herbivory, Zangerl *et al.*, 2002; Nykänen & Koricheva, 2004; Nability *et al.*, 2009). Leaf damage often triggers upregulation of defence-related genes and down-regulation of genes related to photosynthesis (Bilgin *et al.*, 2010). Nevertheless, previous studies have found both increased (“compensatory photosynthesis”) and decreased photosynthetic rate as a response to herbivory (Zangerl *et al.*, 2002; Nykänen & Koricheva, 2004; Nability *et al.*, 2009). Similarly, VOC emission can either increase (as defensive reaction through plant-predator communication or plant-plant signalling) or decrease after leaf damage (Loreto & Sharkey, 1993; Dicke & Baldwin, 2010; Rowen & Kaplan, 2016). The exact plant response to herbivory depends on the characteristics of the specific species interaction, for example on the diet

breath (e.g. specialist vs. generalist) or feeding guild (e.g. chewing vs sap-sucking) of the herbivore (Nykänen & Koricheva, 2004; Kessler & Halitschke, 2007; Rowen & Kaplan, 2016).

Isoprene is one of the most abundant plant-emitted hydrocarbons (Guenther *et al.*, 1995; Wang & Shallcross, 2000), produced by many long-lived woody species (Dani *et al.*, 2014). It is often emitted in small quantities alongside photosynthesis (Rasulov *et al.*, 2009), but also plays a key role as a stress chemical helping the plant to cope with high temperature (Sharkey & Singsaas, 1995; Rasulov *et al.*, 2010). Because isoprene influences the formation and lifetime of lower tropospheric pollutants (Fehsenfeld *et al.*, 1992; Fuentes *et al.*, 2000), changes in isoprene emissions can influence atmospheric chemistry (Mentel *et al.*, 2013; Kravitz *et al.*, 2016). For estimating the effects of insect herbivory on atmospheric chemistry, quantifying herbivory-induced changes in isoprene emissions is of key interest.

To date, most studies assessing the link between herbivory and photosynthesis or isoprene emission have used cultivated model plant species (mostly species in the Brassicaceae or Solanaceae), simulated herbivory (Portillo-Estrada *et al.*, 2015), or controlled greenhouse environments (Kessler & Halitschke, 2007). The effect of herbivory (including its *indirect effects*) on photosynthesis or isoprene emissions in natural systems thus remains largely unknown. In addition, these effects have often been studied at the scale of individual plants or plant parts, and remain poorly quantified at larger scales. This prevents us from drawing conclusions about the large-scale influence of insect herbivory on carbon cycling and atmospheric chemistry.

Using a manipulative experiment, we investigated how a common insect herbivore affects photosynthesis and isoprene emission rate of its host plant in a natural broadleaf deciduous forest. As a study system, we used the pedunculate oak (*Quercus robur* L.) and caterpillars of the winter moth (*Operophtera brumata* L.), both of which are common species throughout temperate woodlands. We measured rates of photosynthesis and isoprene emissions in intact leaves, leaves eaten by herbivores, intact leaves close to eaten leaves (to quantify the systemic effects), and leaves

subject to mechanical damage (to gain insights into how the potential herbivory-induced responses are triggered). Specifically, we addressed the following questions: 1.) Do photosynthetic and/or isoprene emission rates of oak leaves change following leaf damage? 2.) Is the effect different between herbivore-induced damage versus mechanical wounding? 3.) Are damage-induced responses restricted to damaged leaves, or can changes in photosynthetic and/or isoprene emission rates be observed on intact leaves close to their damaged neighbour? 4.) What are the total effects of herbivory-induced leaf area loss (*direct effect*) and changes in the remaining leaf tissue (*indirect effect*) at the canopy scale?

Materials and methods

Experimental setup

The study was carried out during the springs and summers 2015-2016 on ten oak trees (*Quercus robur* L.) in Oxfordshire, UK. Five of the oaks were mature trees (mean diameter at breast height, “dbh” $67.2 \text{ cm} \pm 5.4 \text{ cm SEM}$) located in Wytham Woods ($51^{\circ}.46' 27.48'' \text{ N}$, $1^{\circ} 20' 16.44'' \text{ W}$, 160 m.a.s.l), and the remaining five were young (mean dbh $13.6 \text{ cm} \pm 1.8 \text{ cm SEM}$) planted oaks by the John Krebs field station in Wytham ($51^{\circ} 47' 1.32'' \text{ N}$, $1^{\circ} 19' 1.2'' \text{ W}$, 63 m.a.s.l). Oak is a strong isoprene emitter (Lehning *et al.*, 1999). On both sites, the oaks are naturally infested by caterpillars of the winter moth, which is a common generalist early-spring herbivore. The caterpillars emerge in synchrony with the budburst, and feed on the newly flushed leaves until June (Hunter, 1992). Relatively few herbivore species feed on the mature oak leaves later in the season (Feeny, 1970). Oaks in our study area do not reach their full photosynthetic capacity until late June, (Morecroft *et al.*, 2003), creating a time lag between the peak herbivory and the peak photosynthesis. For herbivores to have substantial impact on photosynthesis in this system, their effect should carry over until the oak has reached its full photosynthetic capacity.

Between 11th and 15th May 2015 and 9th and 11th May 2016, when most leaves were still newly flushed, we identified 15 shoots (of ~ 8 leaves) with only intact leaves from each study tree and enclosed each shoot in a small mesh fabric bag (see Supplementary Information, Methods S1). We randomly assigned each bag into one of the three treatments: 1) *herbivore addition*, 2) *mechanical damage*, or 3) *control*, so that each tree had five bags of each treatment. For each of the *herbivore addition* bags we added a locally collected winter moth caterpillar, and let it feed on the leaves for 3-5 days until at least two of the leaves showed signs of feeding damage. Because the effect of damage often depends on its type and amount (Wu & Baldwin, 2009; Portillo-Estrada *et al.*, 2015), each *herbivory addition* shoot was paired with a *mechanical damage* shoot immediately after the caterpillars had been removed from the mesh bags. The damage on the herbivory shoots was then replicated by tearing or punching holes with a cork borer in the leaves in the mechanical damage treatment. *Control* shoots were left intact. The timing of the manipulations coincided with the peak herbivory in the area (Charmantier *et al.*, 2008). The mesh bags were left around the shoots to prevent additional herbivory until 25th June 2015 or 28th June 2016, when the amount of insect herbivory had levelled off.

One month after the application of the treatments, we randomly chose three shoots from each tree (one *herbivory addition* shoot, one *mechanical damage* shoot, and one *control* shoot) for gas exchange measurements. The few control shoots (n=6) that showed signs of damage were excluded from further measurements. From each *herbivory addition* and *mechanical damage* shoot we measured two leaves: one damaged and one intact. From each *control* shoot we measured one intact leaf. This setup allowed us to measure five leaf-level treatments: damaged leaf in herbivory treatment, undamaged leaf in herbivory treatment, damaged leaf in mechanical treatment, undamaged leaf in mechanical treatment, and intact control leaf. We constructed photosynthetic light response curves (over the period of 28th July - 25th August 2015) for 49 leaves from ten trees and photosynthesis-CO₂ (A/C_i) -curves (over the periods of 26th August - 10th September 2015 and

11th July - 11th August 2016) for 79 leaves from ten different trees (six of the trees were measured on both years) belonging to all the five leaf-level treatments. The timing of the gas exchange measurements corresponded to the peak photosynthetic activity of oak in the study area (Morecroft *et al.*, 2003).

On each leaf, we measured an intact part of an area of 2.5 cm² of the leaf with an infra-red gas analyser (CIRAS-2, PP-Systems, Hitchin, UK). For the light response curves, we took five point measurements on 15 different light levels between 2000 and 0 $\mu\text{mol m}^{-2}\text{s}^{-1}$ of photosynthetically active radiation (PAR). For the A/C_i curves, we measured the photosynthetic rate under ten different CO₂ concentrations between 1300 and 30 ppm. All the raw photosynthesis measurements were processed using the protocol provided by PP-Systems (ppsystems.com) for the CIRAS-2 to apply corrections for the measured variables. The resultant variable used in the analyses was photosynthetic rate per unit leaf area, expressed as $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$.

To study how herbivory and leaf damage affect the production of isoprene by the oak, we measured isoprene emission rate of 32 leaves from seven trees, using the same leaves (and thus the same five leaf-level treatments) as for the A/C_i curves with a portable gas chromatograph (iDirac, see Supporting Information, Methods S2), 21st July - 9th August 2016. iDirac is a novel gas chromatograph, designed for *in-situ* use. Here we report its use for the first time in a field study. We attached the iDirac directly into the CIRAS-2 system to allow for simultaneous measurements of isoprene production and photosynthetic rate. See Supporting Information, Methods S1 for details on all the gas exchange measurements.

After measurements were taken the leaves were photographed to estimate the leaf area lost to herbivory. To estimate the natural level of insect herbivory on the study trees throughout the growing season, we collected 15 additional shoots from each tree on four time points (16-28th May, 25th June, 14th July - 10th August and 18th August 2015), and pressed and scanned the leaves. The area lost to herbivory of the photographed and scanned leaves were estimated as the percentage of

missing area from the side of the leaf, from the tip, or as holes, using the ImageJ software (NIH, MD, USA).

Extracting response parameters.

To calculate the light-saturated photosynthesis, we fitted a Michaelis-Menten equation to the light response data for each leaf separately to estimate the parameters for the maximum light-saturated photosynthetic rate (A_{sat}) and the light intensity at which the gross photosynthetic rate is half of its maximum, K (Marino *et al.*, 2010). To obtain a measure of the mean dark respiration (R_d) for each leaf, we calculated the average photosynthetic rate on the light response curves when the light level was zero. To analyse the photosynthetic response to experimental treatments under different CO_2 concentrations, we constructed A/C_i response curves, where the photosynthetic rate (A) is modelled against the intercellular CO_2 mole fraction (C_i) (Farquhar *et al.*, 1980; Sharkey *et al.*, 2007), allowing us to estimate three important photosynthetic parameters: maximum carboxylation rate, describing the activity of Rubisco (V_{cmax}), rate of photosynthetic electron transport (J_{max}) and triose phosphate use efficiency (TPU). See Supporting Information, Methods S2 for details on model fitting.

After fitting, all the parameters were normalized to 25 °C (Harley *et al.*, 1992) (Sharkey *et al.*, 2007) to reduce variation caused by different ambient temperatures. For most leaves ($n = 65$) the Farquhar *et al.* (1980) model could be fitted to the data. For some leaves ($n = 14$) the model failed to estimate at least one of the parameters. These leaves were omitted from the further analyses of the treatment effects on A/C_i parameters. To study possible changes in leaf conductance, we extracted the mean stomatal conductance (g_s) recorded by the gas analyser during the A/C_i curve measurements. From those leaves of which only light response was measured (24 leaves), we used mean stomatal conductance of the light response curve. Single outlier values of

stomatal conductance, K and isoprene emission were removed from further analyses. See Fig. 2 for final sample sizes per parameter

To estimate isoprene emissions, the height of each isoprene peak in the gas chromatogram was measured and converted into mixing ratios (ppb) by using calibration measurements with known isoprene concentrations. The mixing ratios were scaled with the known air volume, area of leaf measured and flow rate to yield emission rates as $\text{nmol m}^{-2} \text{s}^{-1}$. Because isoprene emission is strongly influenced by temperature, we corrected the measured emission values for temperature (Guenther *et al.*, 1993, 1995), yielding the standard emission factor of isoprene (as $\mu\text{g m}^{-2} \text{h}^{-1}$), I_s (in 303 K and $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation). See Supporting Information, Methods S1 for details on the temperature correction.

To describe the photosynthetic rate of the study leaves in natural conditions, we extracted values from the light-response and A/C_i curves for photosynthetic rates at ambient CO_2 concentration (400 ppm) and in light intensity that corresponds to typical full light conditions ($1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation). This parameter (A_{1000}), was used to assess the correlation between photosynthesis and isoprene emission rate, and to scale up the effects of herbivory from leaf scale to the canopy level.

Statistical analyses. To test for effects of our experimental treatments on photosynthesis and isoprene emission, we built a separate linear mixed effects model for each of the key response parameters described above. Each photosynthesis-related response parameter (A_{sat} , K , R_d , V_{cmax} , J_{max} , TPU , g_s) was modelled as a function of leaf-level treatment (a categorical variable with five levels), site (Wytham Woods or John Krebs field station), mean leaf temperature (to account for any remaining variation by the ambient temperatures), year (2015 or 2016, for the parameters that had been measured in both years), and the percentage of leaf damage as explanatory variables. Time of the day was assumed to have a non-linear effect, and was added as general additive smoother. To

avoid spurious treatment effects due to small sample sizes, interactions were not included (Zuur, 2009). Tree identity and shoot identity (nested within tree identity), were included as random factors (random intercepts) to account for non-independence of leaves on the same shoots and trees. Isoprene emissions (I_s) were modelled using the same approach, except that variance structure was allowed to vary between the leaf treatments to allow for unequal variances across these groups. For each response variable, the full model was simplified by dropping one explanatory variable at a time. The change in the model fit was assessed using likelihood ratio tests. Fixed factors that did not improve model fit were dropped from the final model (Crawley, 2007). Where leaf type was significant, a post-hoc Tukey's test was applied to assess which of the five leaf treatments differed significantly from one other. Because of the adjusted variance structure in the isoprene model, the pairwise leaf treatment comparisons were carried out estimating least square means.

To analyse the relationship between isoprene emission and the photosynthetic parameters measured simultaneously (A_{1000} , V_{cmax} , J_{max} and TPU), we built linear, exponential and quadratic models in which the isoprene emission rate was modelled as a function of each selected photosynthetic parameter. We then estimated the model fit by comparing the adjusted r^2 -values between the different models (linear, exponential and quadratic), and selected the model with the highest r^2 value for each of the parameters.

To test for the differences in the amount of leaf damage between the two damage treatments (mechanical and herbivory) and naturally occurring damaged leaves, we built a linear model with proportion of damage as a function of damage type (herbivore addition, mechanical, natural). To test for patterns in natural herbivory levels, we built a linear model of proportion of damage as a function of the site and the collection date. Proportions were arcsine-square root –transformed in order not to violate model assumptions (Crawley, 2007). For all models, the model assumptions were tested by visually examining plots of residuals against fitted values for the homoscedasticity of residuals, and a Quantile-Quantile plot for the normal distribution of the residuals. All analyses

were conducted using R version 3.4.1 (R Core Team, 2017) and the packages lme4 (Bates *et al.*, 2015), multcomp (Hothorn *et al.*, 2008), nlme (Pinheiro *et al.*, 2017), gamm4 (Wood & Scheipl, 2017) and lsmeans (Lenth, 2016).

Quantifying the effects of herbivory on leaf and canopy scales. To estimate the effects of herbivory on photosynthesis and isoprene emission at the canopy scale, we combined three types of data: 1) the proportion of leaf area loss per leaf under natural conditions (direct effect), 2) the effect of insect herbivory on the photosynthetic rate (A_{sat}) or isoprene emission rate (I_s) per unit leaf area (indirect effect), and 3) information on natural patterns of herbivory in the oak canopy. Control leaves, which were intact leaves on intact shoots were set as a reference point to describe photosynthesis and isoprene emission in the absence of herbivory. To estimate the leaf-scale effect of herbivory on the light-saturated photosynthesis or isoprene emission rate, we first multiplied the per leaf unit area rate of a leaf damaged by herbivores with the proportion of remaining leaf area in the corresponding leaf type, yielding a “per leaf” - rate. We then compared this to a “per leaf” -rate of an intact control leaf:

$$\text{light saturated leaf scale effect}_t = \frac{A_t * (1 - D_t)}{A_{t=1}} - 1$$

(Eq. 1.)

where A is the light-saturated assimilation rate (A_{sat}) or the isoprene emission rate, D is the proportion of leaf area loss per leaf type (= direct effect, between 0 and 1) and t denotes the three different leaf types (1 = intact leaf in a completely intact shoot, 2 = intact leaf in an herbivory treatment, 3 = damaged leaf). For the intact leaves in the herbivory treatment, the leaf scale effect

was simply the percentage change in the photosynthetic or isoprene emission rate, indicating a “shoot-level effect” of herbivory spreading from the damaged leaves to the intact neighbours.

We estimated the effect of herbivory at the level of the canopy with two different methods. Firstly, to estimate the herbivory effect at the level of the canopy for the maximum light-saturated photosynthesis and isoprene emission rate, we multiplied the light saturated leaf-scale effect of each leaf type by the proportion of the respective leaf type in the canopy, and then summed these values over the three leaf types:

$$light\ saturated\ canopy\ effect = \sum_{t=1}^3 leaf\ scale\ effect_t * l_t$$

(Eq. 2.)

where t denotes the three different leaf types and l is the proportion of leaf type t in the canopy. For photosynthesis, this model estimates the maximum potential photosynthesis in full light (as $\mu\text{mol m}^{-2} \text{s}^{-1}$ of leaf_area), without considering light transmission through the canopy.

Secondly, because photosynthesis is strongly affected by the amount of available light, we estimated the effect of herbivory on canopy photosynthesis when the diffusion of light through the canopy is taken into account. To estimate this, we used the Big Leaf approach of The Joint UK Land Environment Simulator (“JULES”, Clark *et al.*, 2011) to estimate canopy assimilation, combined with an estimate for canopy respiration (Mercado *et al.*, 2007). The reduction of direct light through the canopy was calculated by Beer's law (Monsi & Saeki, 1953). As a result, our model estimates instantaneous big-leaf approximated net CO₂ assimilation rate. Assimilation is reduced proportional to the transmission of light through the canopy, while leaf respiration increases as light decreases:

$$NPC = \int_0^{LAI} A_{sat} * \left(\frac{PAR}{K + PAR} \right) * (e^{-k*LAI}) - (0.5 - 0.05 * \ln(PAR * e^{-k*LAI})) * R_d$$

(Eq 3.)

where NPC is canopy net photosynthesis (as $\mu\text{mol m}^{-2} \text{s}^{-1}$ of ground area), A_{sat} is the light-saturated photosynthetic rate, k is a light extinction coefficient, LAI is a canopy leaf area index, PAR is the light intensity (“photosynthetically active radiation”) at the top of the canopy and R_d is the dark respiration rate estimated from the Michaelis-Menten equation (Supporting Information Methods S1, Eq. S2). The light extinction coefficient (k) was set to 0.5 as a previously used estimate for broadleaf forests (Clark *et al.*, 2011), leaf area index (LAI) was set to 7.8 as previously measured for this field site (Fenn *et al.*, 2015) and PAR was set to $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ as a standard daytime light intensity at the top of the canopy. We estimated canopy net photosynthesis for each leaf type (i.e. canopy consisting of only that leaf type), multiplied the estimates with the proportion of the respective leaf type observed in the canopy, and then summed these values over the three leaf types. This estimate was then compared to an estimate of a canopy with intact leaves only. Finally, we included the direct effect of leaf area loss by subtracting the proportion of leaf area loss at canopy level:

$$\text{canopy effect at diffused light} = \left(\frac{\sum_{t=1}^3 NPC_t * l_t}{NPC_{t=1}} - D_c \right) - 1$$

(Eq. 4.)

where t denotes the three different leaf types, l is the proportion of leaf type t in the canopy and D_c is the proportion of leaf area loss (=direct effect) at the canopy scale.

Results

Herbivory under natural and experimental settings. There was no difference between the natural levels of herbivory between the two study sites ($t = -0.55$, $df = 2$, 1461 , $p = 0.58$) and no change throughout the growing season ($t = -1.65$, $df = 2$, 1461 , $p = 0.10$), indicating that early-season herbivory is the dominant type of insect herbivory in the study system. Almost all shoots surveyed for natural herbivory levels had at least one damaged leaf: of the 175 shoots surveyed, only three (1.7%) were completely intact.

The mesh bags successfully prevented herbivores from colonizing the experimental shoots (94 of 100 control shoots remained intact). The amount of leaf damage did not differ between the two damage treatments ($10.88\% \pm 1.84\%$ in mechanical and $14.13\% \pm 1.91\%$ in herbivore addition, $t = -0.90$, $df = 2$, 1086 , $p = 0.37$), but was higher in leaves with experimental herbivory compared to naturally occurring herbivory ($8.45\% \pm 0.39\%$, $t = 3.04$, $p = 0.002$ for herbivore addition and $t = 1.72$, $p = 0.09$ for mechanically damaged). Most leaf damage occurred at sides and tips, and only a small portion as holes (Supporting Information, Table S1).

Treatment-effects on photosynthesis and isoprene emission. Leaf treatment significantly influenced the light-saturated photosynthetic rate A_{sat} ($\chi^2 = 17.31$, $p = 0.002$, $df = 4,8$; Supporting Information, Table S2; Fig. 1a. and 2a), the mean carboxylation rate V_{cmax} ($\chi^2 = 9.51$, $p = 0.05$, $df = 4,11$, Table S2; Fig. 1b and 2d), the mean electron transport rate J_{max} ($\chi^2 = 11.23$, $p = 0.02$, $df = 4,10$, Table S2; Fig. 1c and 2e), the mean stomatal conductance g_s ($\chi^2 = 10.48$, $p = 0.03$, $df = 4,10$, Table S2. Fig. 2g) and the isoprene emission rate I_s ($L_{\text{ratio}} = 23.15$, $p < 0.001$, $df = 4,9$, Table S2; Fig. 2h). Both damaged and undamaged leaves in the herbivore addition shoots experienced a significant reduction in their A_{sat} and J_{max} compared to control leaves ($z = -4.26$, $p < 0.001$ damaged leaves and $z = -4.26$, $p < 0.001$ undamaged leaves for A_{sat} , $z = -38.92$, $z = -2.84$, $p = 0.03$ damaged leaves and $z = -3.24$, $p = 0.01$ undamaged leaves for J_{max}). V_{cmax} was different mainly

between leaves damaged mechanically and intact leaves in the herbivory treatment, but the difference (revealed by the Tukey's test) was only marginally significant ($z = 2.55$, $p = 0.08$). Stomatal conductance (g_s) was different between control and the undamaged leaf in the herbivory treatment ($z = -2.73$, $p = 0.049$). The light intensity at which the gross photosynthetic rate is half of its maximum (K, Fig. 2b), dark respiration (R_d , Fig. 2c), and triose phosphate use efficiency (TPU, Fig. 1d and 2f), on the other hand, were not influenced by leaf treatment. Mean leaf temperature significantly increased V_{cmax} ($\chi^2 = 4.21$, $p = 0.04$, $df = 1$, 11), J_{max} ($\chi^2 = 9.98$, $p = 0.002$, $df = 1$, 10), TPU ($\chi^2 = 9.93$, $p = 0.002$, $df = 1$, 6), R_d ($\chi^2 = 8.11$, $p = 0.004$, $df = 1$, 5) and g_s ($\chi^2 = 5.34$, $p = 0.02$, $df = 1$, 10). V_{cmax} , J_{max} , TPU and g_s were significantly different between the two sites ($\chi^2 = 5.07$, $p = 0.02$, $df = 1$, 11 for V_{cmax} ; $\chi^2 = 5.58$, $p = 0.02$, $df = 1$, 10 for J_{max} ; $\chi^2 = 5.34$, $p = 0.02$, $df = 1$, 6 for TPU and $\chi^2 = 5.95$, $p = 0.01$, $df = 1$, 10 for g_s), and V_{cmax} differed between the two measuring years ($\chi^2 = 8.82$, $p = 0.03$, $df = 1$, 11).

Leaves damaged mechanically had significantly higher isoprene emission rate compared to control leaves and undamaged leaves in the herbivory treatment ($t = -6.57$, $p < 0.007$ and $t = -7.16$, $p < 0.004$, respectively). The isoprene emission rate per unit leaf area decreased with increasing percentage of leaf damage ($L_{ratio} = 8.32$, $p = 0.004$, $df = 1$, 9). Isoprene emission rate correlated positively and significantly with the photosynthetic parameters (Supporting Information, Fig. S4).

The effects of herbivory on leaf and canopy scales. Leaf area loss (the *direct effect* of herbivory) per leaf was $8.5\% \pm 0.4\%$. The *indirect effect* of herbivory, i.e. the herbivory-induced change in photosynthesis in the remaining leaf tissue, accounted for a $45.5\% \pm 10.1\%$ reduction in the leaf-scale light-saturated photosynthesis (A_{sat} , Table 1). Hence, the indirect effect of herbivory was several magnitudes larger than the direct effect of leaf area loss. Within the shoots that had herbivory damage, the reduction in photosynthesis was almost identical between damaged leaves and their undamaged neighbors. When the *direct* and *indirect effects* and the proportion of damaged

and undamaged leaves in the canopy were combined, $45.6\% \pm 7.6\%$ of the light-saturated photosynthetic potential and $47.9\% \pm 9.5\%$ of the net photosynthesis under diffused light was lost to herbivores at the canopy-scale (Table 1). The first estimate represents a canopy consisting only of sun leaves at full light, (see Supporting Information, Table S3 for estimates on canopy-scale effects of herbivory on photosynthesis at lower light intensity), whereas the second estimate represent a canopy where light is reduced with increasing leaf area index due to shading. Despite the different assumptions of these estimates, the proportional change in photosynthesis due to herbivory is effectively the same.

In contrast to the photosynthesis results, isoprene emission rates increased in the damaged leaves by $85.4 \pm 115.6\%$ compared to the intact control leaves, though the small number of samples and the associated large error makes drawing conclusions difficult. The shoot-level effect, where shoot-level herbivory affects undamaged leaves within the same shoot, was small ($29.8 \pm 32.1\%$) for isoprene. At the canopy-scale, the total effect of herbivory corresponded to a $52.5 \pm 82.6\%$ increase in isoprene emissions, but with large variation (Table 1).

Discussion

In this study herbivory substantially reduced photosynthesis in damaged leaves and in their intact neighbours. Isoprene emission rates significantly increased after mechanical leaf damage. At the canopy-scale, these results indicate that even a relatively moderate level of herbivory (6% of canopy leaf area), leads to a 48% reduction in the potential photosynthesis and a 53% increase in isoprene emission rate, although the effect on isoprene emission was not statistically significant at the canopy-scale. Below, we will discuss each of our findings in turn.

Why does the photosynthetic rate change following leaf damage? Previous studies on the indirect effects of herbivory on photosynthesis have reported increases (Oleksyn *et al.*, 1998;

Nykänen & Koricheva, 2004), decreases (Oleksyn *et al.*, 1998; Nabity *et al.*, 2009) and no changes (Peterson *et al.*, 2004) in the assimilation rates after leaf damage. In this study, leaf damage by herbivores lowered the maximum light-saturated photosynthetic rate (A_{sat}), maximum carboxylation rate (V_{cmax}) and the maximum electron transport rate (J_{max}). As stomatal conductance (g_s) correlates with photosynthesis (Wong *et al.*, 1979; Gago *et al.*, 2016), its responses to the treatments were similar to that of photosynthesis. These effects were visible several months after the initial damage. It is unclear whether photosynthesis had remained low during the entire period, or whether the reduction became observable only late in the season. Other studies have reported delayed effects of herbivory on plant physiology, which can be visible several weeks (Gibberd *et al.*, 1988; Meyer, 1998) or even seasons (Kaitaniemi *et al.*, 1998) after the initial damage.

One possibility is that physical injury is inhibiting photosynthesis. Severed vein network can disrupt the transport of water and nutrients with long-lasting effects (Sack & Holbrook, 2006), simultaneously reducing stomatal conductance. Ruptures in the leaf can cause diffusion of CO_2 before it is used in the carbon-fixing reactions, lowering the efficiency of carbon assimilation (Oleksyn *et al.*, 1998; Nabity *et al.*, 2006, 2009, 2013). Furthermore, repairing the damaged tissue uses valuable resources. Trade-offs in resource use might also occur between growth (and hence photosynthesis) and defence (Herms & Mattson, 1992). Defensive reactions against herbivores require synthesis of complex chemical compounds, which act as repellents or additional signalling molecules, using the same resources or molecular pathways than photosynthesis (Herms & Mattson, 1992; Taiz & Zeiger, 2010; Zhou *et al.*, 2015). Build-up of defensive compounds in the plant tissue might also cause the problem of auto-toxicity, lowering photosynthetic efficiency (Baldwin & Callahan, 1993; Nabity *et al.*, 2009). Damage early in the season could also “prime” the plant (Conrath *et al.*, 2002), making it more resistant to future herbivory by activating long-lasting defences. The cost of maintaining a primed state could alter primary metabolism over long-term (van Hulst *et al.*, 2006; Frost *et al.*, 2008).

Why does the photosynthetic rate differ between leaves damaged mechanically or by herbivores? In this study, the mechanically damaged leaves experienced a significantly smaller reduction in their photosynthetic rate than leaves damaged by caterpillars. In previous studies, mechanical damage alone has failed to produce a response in the plant, whereas application of herbivore oral secretions, even without any physical damage, have done so (Korth & Dixon, 1997; Alborn, 1997). The herbivore-induced defensive responses depend on the species identity, specifically on the chemical make-up of the insect saliva (Alborn, 1997; Erb *et al.*, 2012). These herbivory-specific effects are usually mediated through hormonal pathways including jasmonic and salicylic acids, the activation of which also switches off photosynthesising reactions (Wasternack & Hause, 2013). These results suggest that the herbivory-inflicted photosynthetic reduction in our study is a response to the presence of herbivores specifically, instead of leaf damage alone, and possibly actively triggered by the defence machinery of the plant (Kerchev *et al.*, 2012; Zhou *et al.*, 2015).

How does leaf damage affect intact neighbouring leaves? In this study, intact and damaged leaves on the same shoots showed an almost identical degree of reduction in photosynthesis. Damage-triggered defence reactions can travel to intact plant parts through shared vasculature (Jones *et al.*, 1993), as electric signals (Sukhov, 2016), or to neighbour plants through volatile organic compounds (Arimura *et al.*, 2000). This systemic signalling can subsequently affect photosynthesis of intact plant parts (Agrawal, 2000; Barron-Gafford *et al.*, 2012; Meza-Canales *et al.*, 2017). Especially jasmonic acid can travel to systemic tissues (Baldwin & Zhang, 1997; Stratmann, 2003), and accumulate in them (Leitner *et al.*, 2005). Because in our study the systemic changes were detected within individual shoots, the signal has probably travelled through within-shoot vascular connections, which might have also restricted it from reaching the intact control

shoots, or dampened the effect (Orians, 2005). The reduction in photosynthesis in neighbouring leaves might prepare the leaf for the forthcoming herbivory, either by increasing the level of defences at the expense of assimilation, or by actively shutting down the production of further carbohydrates, to provide less nutrition for herbivores (Zhou *et al.*, 2015). Herbivore-specific signalling might also explain why the mechanical treatment responded less than the herbivore addition. Our study thus shows that naturally occurring herbivory can have a considerable effect also on systemic intact leaves. These kinds of shoot-level effects have not been previously taken into account in ecosystem-scale studies.

Why did the isoprene emission rate increase after leaf damage? We observed a significant positive relationship between photosynthesis and isoprene emission, concurrent with previous studies (Rasulov *et al.*, 2009; Copolovici *et al.*, 2017). Nevertheless, the treatment-specific effects on isoprene were opposite to the effects on photosynthesis. The isoprene emission rates per unit leaf area were significantly higher in the mechanically damaged leaves than in non-damaged leaves on the intact control shoots, suggesting that the observed change might not be a response to herbivory specifically. Because the effect was not visible in the surrounding intact leaves, the damage-triggered change in isoprene emission seems to be a leaf-level response. Contrary to our results, previous studies have found *a reduction* in isoprene emission immediately after leaf damage (Loreto & Sharkey, 1993; Portillo-Estrada *et al.*, 2015; Copolovici *et al.*, 2017), but see Ferrieri *et al.*, 2005). VOC emission profile emitted immediately after damage can substantially differ from longer-term emissions (Maja *et al.*, 2014). Nevertheless, most herbivore-induced VOCs are studied immediately after the damage occurs.

Oak could be actively increasing its isoprene emission over a longer period after the damage. Physical injury to the leaf venation network could lead to increased water loss lasting for several days (Aldea *et al.*, 2005). Drought, and a release from it, have been shown to increase

isoprene emissions (Sharkey & Loreto, 1993; Tattini *et al.*, 2015). If mechanical damage caused water stress at the time of the injury, this might have led to an increased isoprene emission later, once the damage had been repaired. Long-term monitoring of damaged-induced isoprene emission is needed to fully understand its response to herbivory.

Canopy scale effect of insect herbivory. At our study site, the *direct effect* of insect herbivory was small: insect herbivores removed 6.0% ($\pm 3.8\%$) of the oak leaf area, consistent with global estimates of average herbivory rates (Cyr & Pace, 1993). The *indirect effect* of herbivory on the remaining leaf tissue of the damaged leaf, and on the neighbouring intact leaves, on the other hand, was several magnitudes larger, reducing the light-saturated photosynthesis by 46% ($\pm 10\%$) and 37% ($\pm 12\%$) on average, respectively. This supports the previous results on the importance of indirect effects over direct ones (Zangerl *et al.*, 2002; Barron-Gafford *et al.*, 2012). Nevertheless, in many ecosystem-scale studies the effects of herbivory are quantified only as the amount of leaf area loss (Metcalf *et al.*, 2014).

By combining indirect effects with the leaf area loss ($8.5\% \pm 0.4\%$ per leaf), we estimate that every damaged leaf has its photosynthetic rate reduced by 50% ($\pm 10\%$). Surveying the natural level of herbivory in the area, only 1.7% of shoots per tree were completely intact. Therefore, most of the oak canopy (98.3%) is photosynthesising below its potential. Effectively no tree in natural settings is devoid of this herbivory-influenced suppression of photosynthesis. On a scale of the canopy, then, only 52% ($\pm 10\%$) of the photosynthesis is realised. Previous studies have not considered the combined direct and indirect effects on ecosystem-level carbon cycle. We show that herbivores can reduce the canopy-scale carbon sequestration considerably, and the shoot-level effect observed in the intact neighbour leaves is a major contributor to this reduction.

Similarly, herbivory had a large effect on isoprene emission, causing an 85% ($\pm 116\%$) increase in the leaf-scale isoprene emission rate and an 53% ($\pm 83\%$) increase on the canopy-scale.

The large error margin makes it difficult to draw firm conclusions on the role of herbivory on canopy-level isoprene emissions. However, if our estimates are correct, this increase would be enough to counteract the predicted reduction in isoprene emissions due to climate change, increasing atmospheric CO₂ concentrations and land-use changes combined (Squire *et al.*, 2014). Despite their potential importance, biotic interactions are usually lacking from the global isoprene emission models (Müller *et al.*, 2008; Arneth *et al.*, 2008; Squire *et al.*, 2014). Previous studies have recorded higher forest-scale isoprene emissions than expected by models (Geron *et al.*, 1997; Gu *et al.*, 2017), and changes in species composition have been shown to affect forest-scale isoprene emissions (Wang *et al.*, 2017). Our study suggests that enhanced emissions resulting from leaf damage might be leading to underestimates of the actual forest-scale isoprene emissions, which could have significant knock-on effects on calculations of ozone and particle formation.

Because emission of isoprene is temperature-sensitive, measurements of temperature change through the different canopy layers would be needed for a more realistic estimate on canopy-level isoprene emissions. Also, further studies on differences between sun and shade leaves and herbivory rates across the canopy, and direct canopy measurements are needed to improve the estimates on canopy photosynthesis and isoprene emissions under herbivory.

With the predicted climate change, species distributions, abundances and hence the frequencies of specific species interactions are projected to shift, and in many cases, have already shifted (Jepsen *et al.*, 2008; Kurz *et al.*, 2008). Nevertheless, insect herbivory is rarely addressed in biosphere and climate models (Kurz *et al.*, 2008). Our results clearly demonstrate that for predicting the responses of forest ecosystems to climate change, including the effects of herbivory on the carbon cycle and atmospheric chemistry is crucial. Ignoring the role of insect herbivory might thus overestimate the role of forests as carbon sinks (Kurz *et al.*, 2008; Schäfer *et al.*, 2010), or underestimate their role as isoprene emitters. We have demonstrated the importance of indirect

herbivory effects for a single plant-herbivore system; there is a clear need to replicate such studies in other systems.

Conclusions. Moth caterpillars reduce the per unit leaf area photosynthetic rate of their host plant, both in the remaining leaf tissue of the damaged leaf, and in the intact neighbour leaves. The reduction by natural herbivory is greater than that by mechanical damage alone. This indicates the host plant can differentiate between these two types of damage, pass on the signal to undamaged parts, and respond accordingly. Isoprene emission rate is increased by mechanical leaf damage, and does not seem to be an herbivory-specific reaction. These responses expressed on a scale of individual leaves and shoots have large-scale consequences on the carbon dynamics on the scale of the forest. On a scale of a canopy, the indirect effects of herbivory emerge several times more important than the direct effect of leaf area removed. Including these effects in estimates of the interactions between biosphere and the atmosphere is crucial for better prediction of the effects of changing climate on forest ecosystems.

Acknowledgements. We thank Nikos Fyllas for help with the photosynthetic models, Andrew Robinson for the help with the isoprene measurements, Angelica Martinez Bauer for help in the field and Nico Raab for the help with the CIRAS-2. Martijn Slot and three anonymous reviewers provided helpful comments on the manuscript. KV was funded by Oskari Huttunen Foundation and the Finnish Cultural Foundation. NERC sponsored iDirac development in BALI (NE/K016377/1) and the Doctoral Training Partnership for CB. YM is supported by a European Research Council Advanced Investigator Award GEM-TRAITS (321131) and the Jackson Foundation. SG is a Royal Society University Research Fellow.

Author contributions. KV, SG and TR designed the research. KV carried out the data collection, analysed the data and wrote the first version of the paper. SG, TR and YM supervised the writing and analysing. CB, IO and SR contributed to the data analyses. CB and NH designed the isoprene measuring system. All authors contributed substantially to revisions.

References

Agrawal AA. 2000. Specificity of induced resistance in wild radish: causes and consequences for two specialist and two generalist caterpillars. *Oikos* **89**: 493–500.

Alborn HT. 1997. An elicitor of plant volatiles from beet armyworm oral secretion. *Science* **276**: 945–949.

Aldea M, Hamilton JG, Resti JP, Zangerl AR, Berenbaum MR, DeLucia EH. 2005. Indirect effects of insect herbivory on leaf gas exchange in soybean. *Plant, Cell and Environment* **28**: 402–411.

Arimura G, Ozawa R, Shimoda T, Nishioka T, Boland W, Takabayashi J. 2000. Herbivory-induced volatiles elicit defence genes in lima bean leaves. *Nature* **406**: 512–515.

Arneth A, Monson RK, Schurgers G, Niinemets Ü, Palmer PI. 2008. Why are estimates of global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)? *Atmospheric Chemistry and Physics* **8**: 4605–4620.

Baldwin IT, Callahan P. 1993. Autotoxicity and chemical defense: nicotine accumulation and carbon gain in solanaceous plants. *Oecologia* **94**: 534–541.

Baldwin IT, Zhang Z-P. 1997. Transport of [2- ¹⁴C] jasmonic acid from leaves to roots mimics wound-induced changes in endogenous jasmonic acid pools in *Nicotiana sylvestris*. *Planta* **203**: 436–441.

- Barron-Gafford GA, Rascher U, Bronstein JL, Davidowitz G, Chaszar B, Huxman TE. 2012.** Herbivory of wild *Manduca sexta* causes fast down-regulation of photosynthetic efficiency in *Datura wrightii*: an early signaling cascade visualized by chlorophyll fluorescence. *Photosynthesis Research* **113**: 249–260.
- Bates D, Mächler M, Bolker B, Walker S. 2015.** Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of Statistical Software* **67**.
- Belovsky GE, Slade JB. 2000.** Insect herbivory accelerates nutrient cycling and increases plant production. *Proceedings of the National Academy of Sciences* **97**: 14412–14417.
- Bilgin DD, Zavala JA, Zhu J, Clough SJ, Ort DR, DeLucia EH. 2010.** Biotic stress globally downregulates photosynthesis genes: Biotic stress downregulates photosynthesis. *Plant, Cell & Environment* **33**: 1597–1613.
- Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC. 2008.** Adaptive phenotypic plasticity in response to climate change in a wild bird population. *Science* **320**: 800–803.
- Clark DB, Mercado LM, Sitch S, Jones CD, Gedney N, Best MJ, Pryor M, Rooney GG, Essery RLH, Blyth E, et al. 2011.** The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics. *Geoscientific Model Development* **4**: 701–722.
- Conrath U, Pieterse CMJ, Mauch-Mani B. 2002.** Priming in plant–pathogen interactions. *Trends in Plant Science* **7**: 210–216.
- Copolovici L, Pag A, Kännaste A, Bodescu A, Tomescu D, Copolovici D, Soran M-L, Niinemets Ü. 2017.** Disproportionate photosynthetic decline and inverse relationship between constitutive and induced volatile emissions upon feeding of *Quercus robur* leaves by large larvae of gypsy moth (*Lymantria dispar*). *Environmental and Experimental Botany* **138**: 184–192.
- Crawley MJ. 2007.** *The R book*. Chichester, England; Hoboken, N.J.: Wiley.

- Cyr H, Pace ML. 1993.** Magnitude and patterns of herbivory in aquatic and terrestrial ecosystems. *Nature* **361**: 148–150.
- Dani KGS, Jamie IM, Prentice IC, Atwell BJ. 2014.** Evolution of isoprene emission capacity in plants. *Trends in Plant Science* **19**: 439–446.
- Dicke M, Baldwin IT. 2010.** The evolutionary context for herbivore-induced plant volatiles: beyond the ‘cry for help’. *Trends in Plant Science* **15**: 167–175.
- Erb M, Meldau S, Howe GA. 2012.** Role of phytohormones in insect-specific plant reactions. *Trends in Plant Science* **17**: 250–259.
- Farquhar GD, von Caemmerer S, Berry JA. 1980.** A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* **149**: 78–90.
- Feeny P. 1970.** Seasonal changes in oak leaf tannins and nutrients as a cause of spring feeding by Winter moth caterpillars. *Ecology* **51**: 565–581.
- Fehsenfeld F, Calvert J, Fall R, Goldan P, Guenther AB, Hewitt CN, Lamb B, Liu S, Trainer M, Westberg H, et al. 1992.** Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry. *Global Biogeochemical Cycles* **6**: 389–430.
- Fenn K, Malhi Y, Morecroft M, Lloyd C, Thomas M. 2015.** The carbon cycle of a maritime ancient temperate broadleaved woodland at seasonal and annual scales. *Ecosystems* **18**: 1–15.
- Ferrieri RA, Gray DW, Babst BA, Schueller MJ, Schlyer DJ, Thorpe MR, Orians CM, Lerdau M. 2005.** Use of carbon-11 in *Populus* shows that exogenous jasmonic acid increases biosynthesis of isoprene from recently fixed carbon. *Plant, Cell and Environment* **28**: 591–602.
- Frost CJ, Mescher MC, Carlson JE, De Moraes CM. 2008.** Plant Defense Priming against Herbivores: Getting Ready for a Different Battle. *Plant Physiology* **146**: 818–824.

- Fuentes JD, Gu L, Lerdau M, Atkinson R, Baldocchi D, Bottenheim JW, Ciccioli P, Lamb B, Geron C, Guenther A, et al. 2000.** Biogenic hydrocarbons in the atmospheric boundary layer: a review. *Bulletin of the American Meteorological Society* **81**: 1537–1575.
- Gago J, Daloso D de M, Figueroa CM, Flexas J, Fernie AR, Nikoloski Z. 2016.** Relationships of leaf net photosynthesis, stomatal conductance, and mesophyll conductance to primary metabolism: a multispecies meta-analysis approach. *Plant Physiology* **171**: 265–279.
- Geron CD, Nie D, Arnsts RR, Sharkey TD, Singaas EL, Vanderveer PJ, Guenther A, Sickles JE, Kleindienst TE. 1997.** Biogenic isoprene emission: Model evaluation in a southeastern United States bottomland deciduous forest. *Journal of Geophysical Research: Atmospheres* **102**: 18889–18901.
- Gibberd R, Edwards PJ, Wratten SD. 1988.** Wound-induced changes in the acceptability of tree-foliage to Lepidoptera: within-leaf effects. *Oikos* **51**: 43.
- Gu D, Guenther AB, Shilling JE, Yu H, Huang M, Zhao C, Yang Q, Martin ST, Artaxo P, Kim S, et al. 2017.** Airborne observations reveal elevational gradient in tropical forest isoprene emissions. *Nature Communications* **8**: 15541.
- Guenther A, Hewitt CN, Erickson D, Fall R, Geron C, Graedel T, Harley P, Klinger L, Lerdau M, Mckay WA, et al. 1995.** A global model of natural volatile organic compound emissions. *Journal of Geophysical Research* **100**: 8873.
- Guenther AB, Zimmerman PR, Harley PC, Monson RK, Fall R. 1993.** Isoprene and monoterpene emission rate variability: Model evaluations and sensitivity analyses. *Journal of Geophysical Research* **98**: 12609.
- Harley PC, Thomas RB, Reynolds JF, Strain BR. 1992.** Modelling photosynthesis of cotton grown in elevated CO₂. *Plant, Cell and Environment* **15**: 271–282.

- Hermes DA, Mattson WJ. 1992.** The dilemma of plants: to grow or defend. *The Quarterly Review of Biology* **67**: 283–335.
- Hothorn T, Bretz F, Westfall P. 2008.** Simultaneous inference in general parametric models. *Biometrical Journal* **50**: 346–363.
- van Hulten M, Pelsler M, van Loon LC, Pieterse CMJ, Ton J. 2006.** Costs and benefits of priming for defense in Arabidopsis. *Proceedings of the National Academy of Sciences* **103**: 5602–5607.
- Hunter MD. 1992.** Interactions within herbivore communities mediated by the host plant: the keystone herbivore concept. In: Effects of resource distribution on animal–plant interactions. Elsevier, 287–325.
- Jepsen JU, Hagen SB, Ims RA, Yoccoz NG. 2008.** Climate change and outbreaks of the geometrids *Operophtera brumata* and *Epirrita autumnata* in subarctic birch forest: evidence of a recent outbreak range expansion. *Journal of Animal Ecology* **77**: 257–264.
- Jones CG, Hopper RF, Coleman JS, Krischik VA. 1993.** Control of systemically induced herbivore resistance by plant vascular architecture. *Oecologia* **93**: 452–456.
- Kaitaniemi P, Ruohomäki K, Ossipov V, Haukioja E, Pihlaja K. 1998.** Delayed induced changes in the biochemical composition of host plant leaves during an insect outbreak. *Oecologia* **116**: 182–190.
- Karl T, Guenther A, Turnipseed A, Patton EG, Jardine K. 2008.** Chemical sensing of plant stress at the ecosystem scale. *Biogeosciences* **5**: 1287–1294.
- Kerchev PI, Fenton B, Foyer CH, Hancock RD. 2012.** Plant responses to insect herbivory: interactions between photosynthesis, reactive oxygen species and hormonal signalling pathways: Redox-mediated crosstalk in insect herbivory. *Plant, Cell & Environment* **35**: 441–453.
- Kessler A, Halitschke R. 2007.** Specificity and complexity: the impact of herbivore-induced plant responses on arthropod community structure. *Current Opinion in Plant Biology* **10**: 409–414.

- Korth KL, Dixon RA. 1997.** Evidence for chewing insect-specific molecular events distinct from a general wound response in leaves. *Plant Physiology* **115**: 1299–1305.
- Kravitz B, Guenther AB, Gu L, Karl T, Kaser L, Pallardy SG, Peñuelas J, Potosnak MJ, Seco R. 2016.** A new paradigm of quantifying ecosystem stress through chemical signatures. *Ecosphere* **7**: e01559.
- Kurz WA, Dymond CC, Stinson G, Rampley GJ, Neilson ET, Carroll AL, Ebata T, Safranyik L. 2008.** Mountain pine beetle and forest carbon feedback to climate change. *Nature* **452**: 987–990.
- Lehning A, Zimmer I, Steinbrecher R, Brüggemann N, Schnitzler J-P. 1999.** Isoprene synthase activity and its relation to isoprene emission in *Quercus robur* L. leaves. *Plant, Cell & Environment* **22**: 495–504.
- Leitner M, Boland W, Mithöfer A. 2005.** Direct and indirect defences induced by piercing-sucking and chewing herbivores in *Medicago truncatula*. *New Phytologist* **167**: 597–606.
- Lenth RV. 2016.** Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software* **69**: 1–33.
- Loreto F, Sharkey TD. 1993.** Isoprene emission by plants is affected by transmissible wound signals. *Plant, Cell and Environment* **16**: 563–570.
- Maja MM, Kasurinen A, Yli-Pirila P, Joutsensaari J, Klemola T, Holopainen T, Holopainen JK. 2014.** Contrasting responses of silver birch VOC emissions to short- and long-term herbivory. *Tree Physiology* **34**: 241–252.
- Marino G, Aqil M, Shipley B. 2010.** The leaf economics spectrum and the prediction of photosynthetic light-response curves: Leaf economics spectrum and light-response. *Functional Ecology* **24**: 263–272.
- Mentel TF, Kleist E, Andres S, Dal Maso M, Hohaus T, Kiendler-Scharr A, Rudich Y, Springer M, Tillmann R, Uerlings R, et al. 2013.** Secondary aerosol formation from stress-induced biogenic emissions and possible climate feedbacks. *Atmospheric Chemistry and Physics* **13**: 8755–8770.

- Mercado LM, Huntingford C, Gash JHC, Cox PM, Jogireddy V. 2007.** Improving the representation of radiation interception and photosynthesis for climate model applications. *Tellus B: Chemical and Physical Meteorology* **59**: 553–565.
- Metcalfe DB, Asner GP, Martin RE, Silva Espejo JE, Huasco WH, Farfán Amézquita FF, Carranza-Jimenez L, Galiano Cabrera DF, Baca LD, Sinca F, et al. 2014.** Herbivory makes major contributions to ecosystem carbon and nutrient cycling in tropical forests. *Ecology Letters* **17**: 324–332.
- Meyer GA. 1998.** Pattern of defoliation and its effect on photosynthesis and growth of Goldenrod. *Functional Ecology* **12**: 270–279.
- Meza-Canales ID, Meldau S, Zavala JA, Baldwin IT. 2017.** Herbivore perception decreases photosynthetic carbon assimilation and reduces stomatal conductance by engaging 12-oxo-phytodienoic acid, mitogen-activated protein kinase 4 and cytokinin perception: Photosynthesis regulation by herbivore perception. *Plant, Cell & Environment* **40**: 1039–1056.
- Monsi M, Saeki T. 1953.** Ueber den Lichtfaktor in den Pflanzengesellschaften und seine Bedeutung fuer die Stoffproduktion. *Journal of Japanese Botany*: 22–52.
- Morecroft MD, Stokes VJ, Morison JIL. 2003.** Seasonal changes in the photosynthetic capacity of canopy oak (*Quercus robur*) leaves: the impact of slow development on annual carbon uptake. *International Journal of Biometeorology* **47**: 221–226.
- Müller J-F, Stavrakou T, Wallens S, De Smedt I, Van Roozendaal M, Potosnak MJ, Rinne J, Munger B, Goldstein A, Guenther AB. 2008.** Global isoprene emissions estimated using MEGAN, ECMWF analyses and a detailed canopy environment model. *Atmospheric Chemistry and Physics* **8**: 1329–1341.
- Nabity PD, Heng-Moss TM, Higley LG. 2006.** Effects of insect herbivory on physiological and biochemical (oxidative enzyme) responses of the halophyte *Atriplex subspicata* (Chenopodiaceae). *Environmental Entomology* **35**: 1677–1689.

- Nabity PD, Zavala JA, DeLucia EH. 2009.** Indirect suppression of photosynthesis on individual leaves by arthropod herbivory. *Annals of Botany* **103**: 655–663.
- Nabity PD, Zavala JA, DeLucia EH. 2013.** Herbivore induction of jasmonic acid and chemical defences reduce photosynthesis in *Nicotiana attenuata*. *Journal of Experimental Botany* **64**: 685–694.
- Nykänen H, Koricheva J. 2004.** Damage-induced changes in woody plants and their effects on insect herbivore performance: a meta-analysis. *Oikos* **104**: 247–268.
- Oleksyn J, Karolewski P, Giertych MJ, Zytowskiak R, Reich PB, Tjoelker MG. 1998.** Primary and secondary host plants differ in leaf-level photosynthetic response to herbivory: evidence from *Alnus* and *Betula* grazed by the alder beetle, *Agelastica alni*. *New Phytologist* **140**: 239–249.
- Orians C. 2005.** Herbivores, Vascular Pathways, and Systemic Induction: Facts and Artifacts. *Journal of Chemical Ecology* **31**: 2231–2242.
- Peterson RKD, Shannon CL, Lenssen AW. 2004.** Photosynthetic responses of legume species to leaf-mass consumption injury. *Environmental Entomology* **33**: 450–456.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2017.** *nlme: Linear and Nonlinear Mixed Effects Models*.
- Portillo-Estrada M, Kazantsev T, Talts E, Tosens T, Niinemets Ü. 2015.** Emission timetable and quantitative patterns of wound-induced volatiles across different leaf damage treatments in aspen (*Populus tremula*). *Journal of Chemical Ecology* **41**: 1105–1117.
- R Core Team. 2017.** *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

- Rasulov B, Huve K, Bichele I, Laisk A, Niinemets Ü. 2010.** Temperature response of isoprene emission in vivo reflects a combined effect of substrate limitations and isoprene synthase activity: a kinetic analysis. *Plant Physiology* **154**: 1558–1570.
- Rasulov B, Huve K, Valbe M, Laisk A, Niinemets Ü. 2009.** Evidence that light, carbon dioxide, and oxygen dependencies of leaf isoprene emission are driven by energy status in hybrid aspen. *Plant Physiology* **151**: 448–460.
- Rowen E, Kaplan I. 2016.** Eco-evolutionary factors drive induced plant volatiles: a meta-analysis. *New Phytologist* **210**: 284–294.
- Sack L, Holbrook NM. 2006.** Leaf hydraulics. *Annual Review of Plant Biology* **57**: 361–381.
- Schäfer KVR, Clark KL, Skowronski N, Hamerlynck EP. 2010.** Impact of insect defoliation on forest carbon balance as assessed with a canopy assimilation model. *Global Change Biology* **16**: 546–560.
- Schoonhoven LM, Loon JJA van, Dicke M. 2005.** *Insect-plant biology*. Oxford ; New York: Oxford University Press.
- Sharkey TD, Bernacchi CJ, Farquhar GD, Singaas EL. 2007.** Fitting photosynthetic carbon dioxide response curves for C₃ leaves. *Plant, Cell & Environment* **30**: 1035–1040.
- Sharkey TD, Loreto F. 1993.** Water stress, temperature, and light effects on the capacity for isoprene emission and photosynthesis of kudzu leaves. *Oecologia* **95**: 328–333.
- Sharkey TD, Singaas EL. 1995.** Why plants emit isoprene. *Nature* **374**: 769–769.
- Squire OJ, Archibald AT, Abraham NL, Beerling DJ, Hewitt CN, Lathière J, Pike RC, Telford PJ, Pyle JA. 2014.** Influence of future climate and cropland expansion on isoprene emissions and tropospheric ozone. *Atmospheric Chemistry and Physics* **14**: 1011–1024.

- Staudt M, Lhoutellier L. 2007.** Volatile organic compound emission from holm oak infested by gypsy moth larvae: evidence for distinct responses in damaged and undamaged leaves. *Tree Physiology* **27**: 1433–1440.
- Stratmann J. 2003.** Long distance run in the wound response – jasmonic acid is pulling ahead. *Trends in Plant Science* **8**: 247–250.
- Strong DR, Lawton JH, Southwood R. 1984.** *Insects on Plants*. Southampton, United Kingdom: Blackwells Scientific Publications.
- Sukhov V. 2016.** Electrical signals as mechanism of photosynthesis regulation in plants. *Photosynthesis Research* **130**: 373–387.
- Taiz L, Zeiger E. 2010.** *Plant physiology*. MA, U.S.A: Sinauer Associates Inc.
- Tattini M, Loreto F, Fini A, Guidi L, Brunetti C, Velikova V, Gori A, Ferrini F. 2015.** Isoprenoids and phenylpropanoids are part of the antioxidant defense orchestrated daily by drought-stressed *Platanus × acerifolia* plants during Mediterranean summers. *New Phytologist* **207**: 613–626.
- Wang K-Y, Shallcross D. 2000.** Modelling terrestrial biogenic isoprene fluxes and their potential impact on global chemical species using a coupled LSM–CTM model. *Atmospheric Environment* **34**: 2909–2925.
- Wang B, Shugart HH, Lerdau MT. 2017.** An individual-based model of forest volatile organic compound emissions—UVAFME-VOC v1.0. *Ecological Modelling* **350**: 69–78.
- Wasternack C, Hause B. 2013.** Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Annals of Botany* **111**: 1021–1058.
- Wong SC, Cowan IR, Farquhar GD. 1979.** Stomatal conductance correlates with photosynthetic capacity. *Nature* **282**: 424–426.
- Wood S, Scheipl F. 2017.** *gamm4: Generalized Additive Mixed Models using ‘mgcv’ and ‘lme4’*.

Wu J, Baldwin IT. 2009. Herbivory-induced signalling in plants: perception and action. *Plant, Cell & Environment* **32**: 1161–1174.

Zangerl AR, Hamilton JG, Miller TJ, Crofts AR, Oxborough K, Berenbaum MR, DeLucia EH. 2002. Impact of folivory on photosynthesis is greater than the sum of its holes. *Proceedings of the National Academy of Sciences* **99**: 1088–1091.

Zhou S, Lou Y-R, Tzin V, Jander G. 2015. Alteration of plant primary metabolism in response to insect herbivory. *Plant Physiology* **169**: 1488–1498.

Zuur AF (Ed.). 2009. *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.

Supporting Information:

Methods S1: Details on the experimental set up and on extracting the gas exchange parameters

Figure S1: Example of a mesh bag

Figure S2: Experimental leaves in herbivory addition and mechanical damage -treatments

Table S1: Leaf area loss at the study area and in the experiment

Figure S3: The average A/Ci response curves per leaf treatment

Figure S4: Correlation between the isoprene emission rate and photosynthetic parameters

Table S2: Coefficient estimates for mixed effects models

Table S3: Effects of herbivory on A₁₀₀₀ on leaf and canopy scales

Methods S2: iDirac overview and operation

Figure 1. The average model predicted response curves. Panel a) shows photosynthetic response to light, b) the maximum carboxylation rate (V_{cmax}), c) the maximum electron transport rate (J_{max}) and d) the maximum triose phosphate use efficiency (TPU). The original measurements are shown as points, and average model fitted parameters per treatment are shown as lines. For panels b-d, the solid points represent measurements used to estimate the corresponding parameter (*i.e.* when $[\text{CO}_2] < 25 \text{ Pa}$ for V_{cmax} , $[\text{CO}_2] > 45 \text{ Pa}$ for J_{max} , and assimilation at its maximum for TPU, see Supporting Information, Methods S1 for details), and the circles show the remaining measurements. The data represent measures from both field sites, and in panels b-d during both measuring years. Note that the effect of site and year has been taken into account in the statistical analyses.

Figure 2. The average parameter values per leaf treatment. Panel a) shows the average maximum model-fitted light-saturated photosynthetic rate (A_{sat}), b) the average light intensity at which the model-fitted photosynthetic rate is half of its maximum (K), c) the average dark respiration rate (R_d), d) the temperature-corrected average maximum carboxylation rate (V_{cmax}), e) the temperature-corrected average maximum electron transport rate (J_{max}), f) the temperature-corrected average triose phosphate use efficiency (TPU), g) the average stomatal conductance (g_s) and h) the average standard isoprene emission rate (I_s). $n=10$ per leaf treatment for the figures in the panels a-c, except $n=9$ for the mechanically damaged leaf and $n=9$ for herbivore undamaged leaf for panel b. For figures in the panels d-f, $n=15$ for control, $n=13$ for the herbivory treatments and $n=12$ for the mechanical treatments. For panel g, $n=19$ for control, $n=18$ for damaged leaf in herbivore treatment and intact leaf in mechanical treatment, and $n=17$ for intact leaf in the herbivore treatment and damaged leaf in the mechanical treatment. For panel h, $n=7$ for control and damaged leaf in the mechanical treatment, $n=6$ for undamaged leaf in the mechanical treatment and intact leaf in the herbivory treatment, and $n=4$ for the damaged leaf in the herbivory treatment. Error bars are ± 1 SEM. Means not sharing a letter are statistically significantly different from one another, e.g. AB

and C in panel a (Tukey's test, $p < 0.05$). Note that the y-axis for respiration (panel c) is expressed as positive values (instead of the negative assimilation rates) to make the graph more intuitive. The data represent measures from both field sites, and in panels d-g during both measuring years. Note that the effect of site and year has been taken into account in the statistical analyses.

Table 1. Total effect of the herbivory from the leaf to the canopy scale. The average percentage of leaf area loss per leaf (D_t , direct effect), the average proportion of different leaf types ($t=1,2,3$) in the canopy, the effect of insect herbivory on the light-saturated photosynthetic rate (A_{sat}) and on the isoprene emission rate per unit leaf area (indirect effect) of the different leaf types, the estimates of the combined (direct + indirect) effects of these at leaf and canopy scales, and the canopy-scale estimates when change in the light intensity through the canopy is taken into account. The effects are expressed relative to the control treatment values (intact leaves in intact shoots). Errors are ± 1 SEM derived through error propagation. See Supporting Information, Table S3 for values for photosynthetic rate in $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (A_{1000}).

Table 1

	Intact leaf, intact shoot (t=1)	Intact leaf, damaged shoot (t=2)	Damaged leaf, damaged shoot (t=3)	Canopy scale total effect
Direct effect				
Leaf area loss (%) (D_i)	0	0	-8.5 ± 0.4	
% of leaves in canopy (l_i)	1.7	27.3 ± 1.9	71.0 ± 1.9	
Canopy scale effect % (D_c)				-6.0 ± 3.8
Light saturated photosynthesis (A_{sat})				
Rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ of leaf area)	19.8 ± 2.2	12.5 ± 1.9	10.8 ± 1.6	
Rate (% of intact)	100	63.1 ± 11.9	54.5 ± 10.1	
Indirect effect per unit leaf area %	0	-36.9 ± 11.9	-45.5 ± 10.1	
Leaf scale effect % (direct + indirect) ^{Eq 1.}	0	-36.9 ± 11.9	-50.1 ± 9.5	
Canopy scale effect % (direct + indirect) ^{Eq 2.}				-45.6 ± 7.60
Isoprene				
Rate ($\mu\text{g m}^{-2} \text{ h}^{-1}$ of leaf)	871.7 ± 257.6	612.1 ± 213.5	1766.0 ± 967.0	
Rate (% of intact)	100	70.2 ± 32.1	202.6 ± 126.0	
Indirect effect per unit leaf area %	0	-29.8 ± 32.1	102.6 ± 126.0	
Leaf scale effect % (direct + indirect) ^{Eq 1.}	0	-29.8 ± 32.1	85.4 ± 115.6	
Canopy scale effect % (direct + indirect) ^{Eq 2.}				52.5 ± 82.6
Light diffused photosynthesis				
Canopy net rate per leaf type ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ of ground area, NPC) ^{Eq 3}	29.96 ± 3.19	17.87 ± 2.59	16.92 ± 2.28	
Canopy net rate combined, weighted with the leaf type proportions ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ of ground area)				17.4 ± 1.83
Canopy net rate (% of intact)				58.1 ± 8.70
Canopy scale effect % (direct + indirect) ^{Eq 4.}				-47.9 ± 9.50




Figure 1.




Figure 2.

New *Phytologist* Supporting Information

Article title: Small-scale indirect plant responses to insect herbivory could have major impacts on canopy photosynthesis and isoprene emission

Authors: Kristiina Visakorpi, Sofia Gripenberg, Yadvinder Malhi, Conor Bolas, Imma Oliveras, Neil Harris, Sami Rifai and Terhi Riutta

Article acceptance date: 11 June 2018

The following Supporting Information is available for this article:

Methods S1 Details on experimental design

Mesh bags were sown from white “tutu mesh”, with a mesh size $\ll 1$ mm. Bags were approximately 20 cm x 10 cm in size, and were attached around the shoots either with a piece of thin metal wire (2015, Fig. S1), or with a piece of thread sown around the mouthpiece of the bag (2016). This setup allowed bags to be attached tightly around the shoots without causing damage to the branch while preventing herbivores from leaving or entering the mesh bag. Since the bags were in place only during the time when the caterpillars were active (9th May – 28th June 2016 and 11th May – 25th June 2015), we estimate that any effects of altered light penetration or microclimate on leaf functioning by the time of the measurements (from 11th July 2016 and 28th July 2015) was small, and consistent across the treatments. After removing the bags, the shoots were marked with a small piece of red (2015) or orange (2016) tape.

To create the *herbivory addition* treatment, we collected caterpillars of winter moth on different oak trees around Wytham Wood during early May in 2015 and 2016. Different larval instars were used for creating the treatments, and specific instar used varied randomly between the mesh bags. For each mesh bag belonging to the *herbivory addition* treatment, we carefully placed one caterpillar on a haphazardly chosen leaf inside the bag, and waited until the caterpillar attached itself to the leaf before closing the bag. We checked each bag three days after the caterpillar addition. If feeding marks were seen on at least two leaves, caterpillar was removed, otherwise it was left in the bag for another 1-2 days. Dead caterpillars were

replaced by fresh ones. All caterpillars were removed after maximum of five days, by which time all shoots in the *herbivory addition* treatment had experienced sufficiently damage.

The *mechanical damage* treatment was created by first pairing up each shoot belonging to that treatment with an *herbivory addition* –shoot (Fig. S2) immediately after removing the caterpillars from the *herbivory addition* -shoots. The amount and type of damage caused by the caterpillar on the leaves in the shoot belonging to the *herbivory addition* -treatment was estimated visually as percentage of leaf area loss from the sides, from the tip, or as holes. The location of the holes was noted (e.g. near the midrib, close to the side). The damage was then carefully replicated on the leaves in the *mechanical damage* -shoot. Damage on the sides and leaf tip was created by tearing, and holes were created by punching with a cork borer. The mesh bags were left around the shoots (including the *Control* shoots) until 25th June 2015 or 28th June 2016 to protect the leaves from further herbivory until the amount of insect herbivory had levelled off.

Details on the gas exchange measurements

We constructed photosynthetic light response curves during the period of 28th July - 25th August 2015 for leaves from all the ten study trees (49 leaves). We constructed photosynthesis- CO_2 (A/C_i) -curves during two years, over the periods of 26th August - 10th September 2015 and 11th July - 11th August 2016. During the measuring period of 2015, we measured A/C_i curves for leaves from six trees (28 leaves). During 2016, we measured A/C_i curves for leaves from all the ten experimental trees (51 leaves). Each response curve was measured once per leaf. Leaves on treatment and control shoots not used for gas exchange measurements were later collected for other analyses (Visakorpi et al., in prep.)

For all gas exchange measurements, relative humidity was kept between 60 - 80 %, temperature as ambient, and flow rate at 200 ml min^{-1} . Measurements were taken throughout the day between 9:00 h and 20:00 h, except when leaves showed signs of stomatal closure, thus inhibiting photosynthesis (on particularly sunny and dry days). The average daytime air temperature during the measuring period 2015 was $15.7 \text{ }^\circ\text{C}$ ($\pm 0.1 \text{ }^\circ\text{C SEM}$) and 2016 $17.1 \text{ }^\circ\text{C}$ ($\pm 0.2 \text{ }^\circ\text{C SEM}$) (Rennie *et al.*, 2017), which are within the range of normal summer temperatures

in the area (Morecroft *et al.*, 2003). Leaf temperatures ranged between 15 °C and 36 °C with a mean of 23 °C (± 0.1 °C SEM).

For the light response curves, we took five point measurements on 15 different light levels (2000, 1500, 1000, 750, 500, 400, 300, 250, 200, 150, 100, 75, 50, 25 and 0 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (PAR)), starting from the highest level. We allowed the leaf to settle to each new light level for 2 minutes (after testing this was a sufficient time for the leaf to settle to new light conditions) and kept CO₂ concentration at 400 ppm.

For the A/C_i curves, we measured the photosynthetic rate under ten different CO₂ concentrations. Since the gas-analyser was slow to settle to each new exact CO₂ concentration (~15 min with each change in concentration), we set the analyser to find “an approximate” concentration near each target concentration (1500, 1000, 750, 500, 400, 300, 200, 150, 100 and 50 ppm). The realized CO₂ concentrations on average (\pm SEM) across the different measurements were 1336 \pm 4.8, 885 \pm 2.9, 702 \pm 2.2, 513 \pm 1.7, 423 \pm 1.3, 328 \pm 1.0, 234 \pm 0.7, 142 \pm 0.5, 63 \pm 0.3 and 28 \pm 0.1 ppm. This set-up considerably shortened the time it took for the CO₂ concentration to settle (5 min), and allowed the CO₂ concentration to stay stable during each measurement. Since the variation in the realized CO₂ concentrations between the samples was small and random across the treatments, its effect is most likely small. For each leaf, we started at the highest concentration and took three measurements in each concentration. Light intensity was kept at 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$. For both the light and A/C_i measurements, the leaves were allowed to settle for the highest light or CO₂ concentration until there was no consistent change in the photosynthetic rate (usually after 30 min).

For the isoprene measurements the light was kept at 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of PAR, and temperature at the ambient level (leaf temperatures ranging between 18 °C and 35 °C). Since some of the isoprene measurements were taken simultaneously with the A/C_i curves, the CO₂ concentration was again set to approach 400 ppm approximately, in order to keep the measuring conditions similar between the different isoprene samples. The average realized CO₂ concentration was 321 ppm (± 2.6 ppm SEM). Each leaf was measured three times in a row. To record the ambient level of isoprene, we measured the air surrounding the leaf before and after every set of three measurements. The three replicate measurements per leaf were

averaged to obtain a single value for each measured leaf. To account for the ambient isoprene concentration, the average of the ambient measurements taken before and after each measured leaf was subtracted from the estimated leaf emission values.

Extracting the gas exchange parameters

To calculate the light-saturated photosynthesis (A_{sat}), we fitted a Michaelis-Menten equation to the light response data for each leaf separately:

$$A(I) = \frac{G_{\text{max}}I}{K + I} - R_d$$

(Eq. S1.)

where A is the photosynthetic (assimilation) rate per given light intensity (I). From these variables, the model estimates the parameters for maximum gross photosynthetic rate (G_{max}), the leaf respiration rate (R_d , a model fitted parameter, hence not used in subsequent analyses as the respiration rate) and the light intensity at which the gross photosynthetic rate is half of its maximum (K , Marino *et al.*, 2010). The five point measurements per light level were averaged before fitting the curves.

To analyse the photosynthetic response to experimental treatments under different CO_2 concentrations, we first calculated an average of the three repeated measures per CO_2 concentration per leaf. We then constructed A/C_i response curves for each leaf, where the photosynthetic rate (A) is modelled against the intercellular CO_2 mole fraction (C_i). We fitted the model for photosynthesis as described by Farquhar *et al.* (1980) and Sharkey *et al.* (2007). In this model, the photosynthetic reactions are assumed to be in one of the three steady states: in Rubisco-limited photosynthesis (normally on low C_i), in RuBP regeneration-limited photosynthesis, or in triose phosphate use limited photosynthesis (Farquhar *et al.*, 1980; Sharkey *et al.*, 2007). Fitting this model allowed us to estimate three important parameters describing the photosynthetic efficiency: maximum carboxylation rate, describing the activity of

Rubisco (V_{cmax}), rate of photosynthetic electron transport (J_{max}) and triose phosphate use efficiency (TPU).

When photosynthesis is limited by the availability of Rubisco, the response to CO_2 concentrations can be described as:

$$A = V_{cmax} \left[\frac{C_c - \Gamma^*}{C_c + K_c \left(1 + \frac{O}{K_o}\right)} \right] - R_d$$

(Eq. S2.)

where the parameter V_{cmax} is the maximum velocity of Rubisco for carboxylation, and R_d is the daytime respiration rate. The model variables are as follows: C_c is the CO_2 partial pressure at Rubisco, transformed from C_i (Sharkey *et al.*, 2007). K_c is the Michaelis constant of Rubisco for carbon dioxide (set to 40.4 Pa at 25 °C and then adjusted to the actual leaf temperature) and K_o is the Michaelis constant of Rubisco for oxygen (set to 24.8 kPa at 25 °C and then adjusted to the actual leaf temperature). These variables describe the kinetic properties of Rubisco, and their values were taken from previous literature on experiments with tobacco leaves (von Caemmerer *et al.*, 1994; Dreyer *et al.*, 2001). O is the partial pressure of oxygen at Rubisco (set to 21 kPa, as defined by the altitude), and Γ^* is the CO_2 concentration of the photorespiratory compensation point, i.e. the point where CO_2 uptake by photosynthesis is exactly compensated by the release of CO_2 by photorespiration (set to 3.7 Pa at 25 °C and then adjusted to the actual leaf temperature; (Manter & Kerrigan, 2004; Sharkey *et al.*, 2007). Since the model parameters (K_c , K_o and Γ^*) have their own temperature responses, the variables were adjusted to the actual leaf temperature (Sharkey *et al.*, 2007). We assumed the Rubisco-limited state to occur with C_i below 25 Pa, and that the transition between the Rubisco and RuBP limited states occurs between the C_i of 25 Pa and 45 Pa, as these are commonly used and conservative estimates of the upper and lower limits of the transitional stage (Sharkey *et al.*, 2007).

When photosynthesis is limited by the regeneration of RuBP, it can be described with the following equation:

$$A = J \frac{C_c - \Gamma^*}{4C_c + 8\Gamma^*} - R_d$$

(Eq. S3.)

where J is the rate of electron transport and other variables and parameters are as above. This equation allows the estimation of the maximum electron transport rate J_{\max} that could be obtained in saturating light. We assumed this state to occur with C_i was above 45 Pa.

When the triose phosphate use is the limiting factor, the photosynthetic rate is:

$$A = 3TPU - R_d$$

(Eq. S4.)

where TPU is the rate of use of triose phosphates. TPU-limited photosynthesis describes CO_2 saturated state, where photosynthetic rate (A) stays stable or even decreases with increasing CO_2 concentration. We assumed this state to occur when the CO_2 concentration was at its maximum level.

The temperature correction for isoprene

Since isoprene emission is strongly influenced by temperature and light, the emission values were corrected for temperature with the following equation (Guenther *et al.*, 1993, 1995)

$$I = I_s C_L C_T$$

(Eq. S5.)

where I_s is the standard emission of isoprene (as $\mu\text{g m}^{-2} \text{h}^{-1}$) in standard temperature and light conditions (303 K and $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation). Since the

measurements were taken in standard light intensity, the light dependent factor C_L can be ignored. The temperature dependent factor C_T is described as

$$C_T = \frac{\exp\left(\frac{C_{T1}(T - T_S)}{RT_S T}\right)}{C_{T3} + \exp\left(\frac{C_{T2}(T - T_M)}{RT_S T}\right)}$$

(Eq. S6.)

where C_{T1} (95 kJ mol⁻¹), C_{T2} (230 95 kJ mol⁻¹), C_{T3} (0.961) and T_M (314 K) are empirically determined coefficients (Guenther *et al.*, 1993), R is the universal gas constant (8.314 J K⁻¹ mol⁻¹), T_S (303 K) is the standard leaf temperature and T is the actual leaf temperature (in K). For the actual leaf temperatures, we averaged the leaf temperature measurements taken with the gas analyser over the period when isoprene was being measured on each leaf.

References

- von Caemmerer S, Evans JR, Hudson GS, Andrews TJ. 1994.** The kinetics of ribulose-1,5-bisphosphate carboxylase/oxygenase in vivo inferred from measurements of photosynthesis in leaves of transgenic tobacco. *Planta* **195**: 88–97.
- Dreyer E, Le Roux X, Montpied P, Daudet FA, Masson F. 2001.** Temperature response of leaf photosynthetic capacity in seedlings from seven temperate tree species. *Tree Physiology* **21**: 223–232.
- Farquhar GD, von Caemmerer S, Berry JA. 1980.** A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* **149**: 78–90.
- Guenther A, Hewitt CN, Erickson D, Fall R, Geron C, Graedel T, Harley P, Klinger L, Lerdau M, Mckay WA, *et al.* 1995.** A global model of natural volatile organic compound emissions. *Journal of Geophysical Research* **100**: 8873.
- Guenther AB, Zimmerman PR, Harley PC, Monson RK, Fall R. 1993.** Isoprene and monoterpene emission rate variability: Model evaluations and sensitivity analyses. *Journal of Geophysical Research* **98**: 12609.

Manter DK, Kerrigan J. 2004. A/Ci curve analysis across a range of woody plant species: influence of regression analysis parameters and mesophyll conductance. *Journal of Experimental Botany* **55**: 2581–2588.

Marino G, Aqil M, Shipley B. 2010. The leaf economics spectrum and the prediction of photosynthetic light-response curves: Leaf economics spectrum and light-response. *Functional Ecology* **24**: 263–272.

Morecroft MD, Stokes VJ, Morison JIL. 2003. Seasonal changes in the photosynthetic capacity of canopy oak (*Quercus robur*) leaves: the impact of slow development on annual carbon uptake. *International Journal of Biometeorology* **47**: 221–226.

Rennie S, Adamson J, Anderson R, Andrews C, Bater J, Bayfield N, Beaton K, Beaumont D, Benham S, Bowmaker V, et al. 2017. UK Environmental Change Network (ECN) meteorology data: 1991-2015.

Sharkey TD, Bernacchi CJ, Farquhar GD, Singsaas EL. 2007. Fitting photosynthetic carbon dioxide response curves for C₃ leaves. *Plant, Cell & Environment* **30**: 1035–1040.

Fig. S1 Example of a mesh bag

One of the mesh bags during the spring 2015 used to create the shoot-level experimental manipulations



Fig. S2 Experimental leaves in herbivory addition and mechanical damage -treatments.

Example of a pair of leaves used in the gas exchange measurements. A.) shows a leaf damaged by herbivores and b.) shows a mechanically damaged leaf.









Table S1 Table on leaf area loss at the site and in the experiment

The average level of leaf area lost to herbivory naturally on the two experimental sites, and in the two damage treatments (mechanical damage and herbivore addition). Errors are ± 1 SEM.

	Natural herbivory survey	Experimentally manipulated shoots with herbivore damage	Experimentally manipulated shoots with mechanical damage
Number of shoots surveyed	175	19	19
Leaf area removed per eaten leaf, %	8.45 ± 0.39	14.13 ± 1.91	10.88 ± 1.84
Leaf area lost on the side, %	3.75 ± 0.28	8.22 ± 1.34	6.65 ± 1.63
Leaf area lost on tip, %	2.15 ± 0.28	5.53 ± 1.59	4.09 ± 1.56
Leaf area lost as holes, %	0.17 ± 0.02	0.37 ± 0.34	0.14 ± 0.1

Fig. S3 The average A/C_i response curves per leaf treatment.

The average A/C_i curves as derived from the Farquhar et al. (1980) model for each leaf type (a – Control, b – damaged leaf on a herbivore damaged shoot, c – intact leaf on a herbivore damaged shoot, d – damaged leaf on a mechanically damaged shoot, e – intact leaf on a mechanically damaged shoot). The points show raw measurements of photosynthetic rate (A) plotted against the internal CO_2 concentration (C_i). The orange line represents photosynthetic rate if carboxylation capacity is limiting (V_{cmax}), the green line represents electron transport limited photosynthesis (J_{max}), and the blue line shows photosynthesis under triose phosphate use limitation (TPU). The dashed vertical lines at C_i concentrations of 25 and 45 Pa represent the points at which the limiting factor of the photosynthetic rate was assumed to change (from Rubisco limited into RuBP limited photosynthesis).



— V_{cmax}
— J_{max}
— TPU

Fig. S4 The correlation between the isoprene emission rate and photosynthetic parameters.

Isoprene emission rate correlated positively and significantly with the photosynthetic parameters A_{1000} ($r^2 = 0.37$, $p = 0.001$), V_{cmax} ($r^2 = 0.49$, $p < 0.001$), J_{max} ($r^2 = 0.64$, $p < 0.001$) and TPU ($r^2 = 0.62$, $p < 0.001$) Panel a) shows relationship between the isoprene standard emission factor and A_{1000} (photosynthetic rate at 400 ppm of CO_2 and $1000 \mu mol m^{-2} s^{-1}$ of photosynthetically active radiation), b) the relationship between isoprene emission and carboxylation rate V_{cmax} , c) the relationship between isoprene emission and the mean electron transport rate J_{max} , and d) the relationship between isoprene emission and triose phosphate use efficiency TPU. All the parameters except A_{1000} are corrected for standard temperature. The raw data are shown as points. The solid line shows model estimated mean for a quadratic linear (a) and linear (b, c, d) model, and dotted lines represent 95% confidence intervals. Note that the three parameters extracted from the A/Ci curves (V_{cmax} , J_{max} and TPU) correlate with each other and thus show almost identical relationship with isoprene.




Table S2 Coefficient estimates for the linear mixed effects models

Coefficient estimates for fixed effects of the linear mixed effects models assessing the relationships between variables reflecting photosynthetic capacity, leaf respiration, stomatal conductance or isoprene emissions and explanatory variables of the final model. Shown are estimates from the minimum adequate models for each response variable. Note that the negative slope of leaf temperature on respiration means that temperature has a positive effect on the rate of respiration, which has been measured as negative assimilation rates. For each model, the intercept indicates the mean value of the response variable for a given level of the fixed effect(s) and for a covariate value = 0 (stated in parentheses), and the other effects show the mean change, compared with the intercept, caused by the other fixed factor levels and by a unit change in covariate value.

Response variable	Final model	Effects	Estimate	Std error	t-value
A_{sat}	~ leaf treatment	Intercept (Control)	19.82	1.80	11.03
		Herbivore damaged	-9.03	2.12	-4.26
		Herbivore undamaged	-7.35	2.12	-3.47
		Mechanical damaged	-2.12	2.19	-0.97
		Mechanical undamaged	-3.90	2.12	-1.84
K	~ 1	Intercept	171.19	18.38	9.32
Respiration	~ leaf temperature	Intercept (leafT = 0)	0.2	0.25	1.06
		Leaf temperature	-0.035	0.012	-2.95
V_{cmax}	~ leaf treatment + site + leaf temperature + year	Intercept (Control, Site			
		John Krebs, leafT = 0, year	133.56	23.92	5.58
		= 2015)			
		Herbivore damaged	-20.99	10.29	-2.04
		Herbivore undamaged	-24.20	10.23	-2.37

		Mechanical damaged	3.30	10.46	0.32
		Mechanical undamaged	-5.00	10.64	-0.47
		Site (Wytham Woods)	32.43	14.46	2.24
		Leaf temperature	-2.37	1.07	-2.21
		Year (2016)	-26.15	8.69	-3.01
J _{max}	~ leaf treatment	Intercept (Control, Site	352.42	55.50	6.35
	+ site	John Krebs, leafT = 0)			
	+ leaf	Herbivore damaged	-74.89	26.34	-2.84
	temperature	Herbivore undamaged	-84.97	26.26	-3.24
		Mechanical damaged	-23.03	26.32	-0.88
		Mechanical undamaged	-39.75	26.47	-1.50
		Site (Wytham Woods)	77.59	32.70	2.37
		Leaf temperature	-8.51	2.40	-3.55
TPU	~ site	Intercept (Site John Krebs,	17.83	2.74	6.50
	+ leaf	leafT = 0)			
	temperature	Site (Wytham Woods)	3.61	1.56	2.32
		Leaf temperature	-0.43	0.13	-3.39
Stomatal conductance	~ site	Intercept (Site John Krebs,	270.94	46.68	5.80
	+ leaf	leafT = 0, Control)			
	temperature +	Herbivore damaged	-59.82	22.94	-2.61
	leaf treatment	Herbivore undamaged	-63.88	23.34	-2.74
		Mechanical damaged	-16.95	22.76	-0.74
		Mechanical undamaged	-45.92	22.64	-2.03
		Site (Wytham Woods)	70.29	27.92	2.52
		Leaf temperature	-5.12	1.98	-2.58
Isoprene	~ leaf treatment	Intercept (Control,	912.8	289.9	3.15
	+ Damage %	Damage = 0)			

Herbivore damaged	3707.0	1254.8	2.95
Herbivore undamaged	-285.6	201.3	-1.42
Mechanical damaged	5383.2	844.0	6.38
Mechanical undamaged	1582.2	838.1	1.89
Damage %	-176.8	48.2	-3.67

Table S3 Canopy- and leaf-level effects of herbivory on A1000

Canopy- and leaf-level effects of herbivory on photosynthesis in $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (A_{1000}) The effects are expressed relative to the control treatment values (intact leaves in intact shoots). Errors are ± 1 SEM derived through error propagation.

	Intact leaf, intact shoot (1)	Intact leaf, damaged shoot (2)	Damaged leaf, damaged shoot (3)	Canopy scale total effect
A_{1000}				
Rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)	10.8 ± 1.35	7.1 ± 0.9	7.9 ± 1.1	
Rate (% of intact)	100	65.7 ± 11.7	73.1 ± 13.7	
Indirect effect per unit leaf area %	0	-34.3 ± 11.7	-26.9 ± 13.7	
Leaf scale effect % (direct + indirect, Eq. 1)	0	-34.3 ± 11.7	-33.1 ± 12.9	
Canopy scale effect % (direct + indirect, Eq. 2)				-32.8 ± 9.7

Methods S2 iDirac overview and operation

iDirac is a simple gas chromatograph with a photo-ionisation detector (GC-PID) which was designed to make continuous measurements of isoprene away from the laboratory environment (Bolas et al, in preparation). Samples are collected on a Carboxen absorbent trap which is then heated in a flow of nitrogen and passed into a short pre-column. Once isoprene has passed through onto the main column, the flow is reversed and the rest of the sample vented to minimise contamination. A prepared isoprene mixture is used at regular intervals to provide calibration linked to the scale provided by the National Physical Laboratory. Different volumes of this calibration gas are collected on the absorbent trap to provide calibration across a range of mixing ratios. Blank runs were run before each calibration run to ensure to keep the absorbent trap free of contaminants. Calibration programme was run before (15th - 19th July), in the middle (29th July), and after the campaign (4th-6th November). For the leaf measurements presented here, the air being analysed is the exhaust from a CIRAS-2 (PP-Systems, Hitchin, UK) photosynthesis measurement leaf chamber. Analysis of the chromatograms is done by Igor (IGOR Pro Version 6.3.7.2, WaveMetrics Inc, 1988-2014, www.wavemetrics.com) fitting a linear baseline and a Gaussian curve to the isoprene peak.

The overall sampling time for the high isoprene mixing ratios encountered here is 3 minutes. The detection limit of the instrument depends on the volume of air sampled for a particular experiment. Full uncertainty characterisation is being carried out. However, our preliminary estimate for the samples volumes used here is a few 10s of ppt. This means that the observed variability in the emission rates (see below) is dominated by the variation between experiments and not by instrumental uncertainty. We estimate a precision of $\pm 7.80\%$ based on repetitive measurements of the calibration gas before, during and after the experiment.

Small-scale indirect plant responses to insect herbivory could have major impacts on canopy photosynthesis and isoprene emission

Visakorpi, Kristiina

2018-07-26

Attribution-NonCommercial 4.0 International

Visakorpi K, Gripenberg S, Malhi Y, et al., Small-scale indirect plant responses to insect herbivory could have major impacts on canopy photosynthesis and isoprene emission. *New Phytologist*, Volume 220, Issue 3, November 2018, pp. 799-810

<http://dx.doi.org/10.1111/nph.15338>

Downloaded from CERES Research Repository, Cranfield University