

COMMENT OPEN



Indicate separate contributions of long-lived and short-lived greenhouse gases in emission targets

Myles R. Allen¹✉, Glen P. Peters², Keith P. Shine³, Christian Azar⁴, Paul Balcombe⁵, Olivier Boucher⁶, Michelle Cain⁷, Philippe Ciais⁸, William Collins⁹, Piers M. Forster¹⁰, Dave J. Frame¹¹, Pierre Friedlingstein¹², Claire Fyson¹³, Thomas Gasser¹⁴, Bill Hare¹³, Stuart Jenkins¹⁵, Steven P. Hamburg¹⁶, Daniel J. A. Johansson⁴, John Lynch¹⁵, Adrian Macey¹¹, Johannes Morfeldt¹⁴, Alexander Nauels¹³, Ilissa Ocko¹⁶, Michael Oppenheimer¹⁷, Stephen W. Pacala¹⁷, Raymond Pierrehumbert¹⁵, Joeri Rogelj¹⁸, Michiel Schaeffer¹³, Carl F. Schleussner¹³, Drew Shindell¹⁹, Ragnhild B. Skeie², Stephen M. Smith¹⁵ and Katsumasa Tanaka¹⁸

npj Climate and Atmospheric Science (2022)5:5; <https://doi.org/10.1038/s41612-021-00226-2>

As researchers who have published over recent years on the issue of comparing the climate effects of different greenhouse gases, we would like to highlight a simple innovation that would enhance the transparency of stocktakes of progress towards achieving any multi-decade-timescale global temperature goal. In addition to specifying targets for total CO₂-equivalent emissions of all greenhouse gases, governments and corporations could also indicate the separate contribution to these totals from greenhouse gases with lifetimes around 100 years or longer, notably CO₂ and nitrous oxide, and the contribution from Short-Lived Climate Forcers (SLCFs), notably methane and some hydrofluorocarbons. This separate indication would support an objective assessment of the implications of aggregated emission targets for global temperature, in alignment with the UNFCCC Parties' Decision (4/CMA.1)¹ to provide "information necessary for clarity, transparency and understanding" in nationally determined contributions (NDCs) and long-term low-emission development strategies (LT-LEDs).

While differences remain between us regarding how best to set fair yet ambitious targets for individual emitters^{2–5}, including how any additional information might be used, and the interpretation of the Paris Agreement, it is important to emphasise the high level of agreement on the underlying science of how different greenhouse gases affect global temperature. The 2018 IPCC Special Report on 1.5 °C (SR1.5)⁶ stated "Reaching and sustaining net-zero global anthropogenic CO₂ emissions and declining net non-CO₂ radiative forcing (Planetary energy imbalance resulting directly from human-induced changes.) would halt anthropogenic global warming on multi-decadal timescales (*high confidence*). The maximum temperature reached is then determined by cumulative net global anthropogenic CO₂ emissions up to the time of net zero CO₂ emissions (*high confidence*) and the level of non-CO₂ radiative forcing in the decades prior to the time that maximum temperatures are reached (*medium confidence*)". The IPCC 6th Assessment Report (AR6)⁷ confirmed "limiting human-induced global warming to a specific level requires limiting cumulative CO₂ emissions, reaching at least net zero CO₂ emissions, along with strong reductions in other greenhouse gas emissions".

Parties to the Paris Agreement agreed in Katowice in 2018 (Decision 18/CMA.1)¹ to report past emissions of individual gases separately and use 100-year Global Warming Potentials (GWP₁₀₀)

when aggregating them to CO₂-equivalent (we refer to these here as CO₂-e₁₀₀ emissions). The separate specification of individual gases minimises ambiguity in determining the climate impact of past emissions. NDCs and other future targets are, however, almost always expressed in terms of aggregate CO₂-e₁₀₀ emissions only, for which the implications for global temperature are ambiguous^{8,9}. Separate specification of the contribution from CO₂ helps, but ambiguity in global temperature outcomes remains if targets for non-CO₂ gases comprise a mixture of long-lived climate forcers (LLCFs), such as nitrous oxide, with atmospheric lifetimes around 100 years or longer, and SLCFs, such as methane, most of which have lifetimes shorter than 20 years¹⁰.

Specifying the contributions of all gases individually in future targets as well as the reporting of past emissions would resolve the ambiguity in global temperature outcomes, and would also help quantify non-climate benefits of emission reductions, especially for methane¹¹. Governments and particularly corporations may, however, wish to retain some level of aggregation across gases to allow flexibility in how they achieve their targets. Fortunately, a much less restrictive approach delivers almost all the transparency benefits from a climate perspective. The climate system responds similarly over a broad range of timescales to equal emissions expressed in tonnes of CO₂-e₁₀₀ of all LLCFs, including CO₂¹². Likewise, the net radiative forcing due to SLCFs on multi-decadal timescales is similar to the aggregated rate of SLCF emissions expressed in tonnes of CO₂-e₁₀₀ per year multiplied by the 100-year Absolute Global Warming Potential (AGWP₁₀₀) of CO₂¹³. With this additional information, it is straightforward to express the SR1.5 statement quoted above in terms of CO₂-e₁₀₀ emissions: human-induced warming over any multi-decade time-interval is approximately the sum of (i) aggregate CO₂-e₁₀₀ emissions of LLCFs, including CO₂, multiplied by a constant parameter, the Transient Climate Response to cumulative CO₂ Emissions, or TCRE¹⁴ (the TCRE can alternatively be thought of as the Absolute Global Temperature-Change Potential for a sustained emission of CO₂ divided by the time-horizon, AGTP_y/H¹³); (ii) any change in decadal-average radiative forcing due to SLCFs multiplied by another constant parameter, the Transient Climate Response to Forcing, or TCRF, another name for the "fast" component(s) of the climate response¹⁵; and (iii) a

¹School of Geography and the Environment and Department of Physics, University of Oxford, Oxford, UK. ²CICERO Centre for International Climate Research, Oslo, Norway. ³Department of Meteorology, University of Reading, Reading, UK. ⁴Chalmers University of Technology, Göteborg, Sweden. ⁵Queen Mary University of London, London, UK. ⁶Institute Pierre Simon Laplace, Gif-sur-Yvette, France. ⁷Cranfield University, Bedford, UK. ⁸Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France. ⁹University of Reading, Reading, UK. ¹⁰University of Leeds, Leeds, UK. ¹¹Victoria University of Wellington, Wellington, New Zealand. ¹²University of Exeter, Exeter, UK. ¹³Climate Analytics, Berlin, Germany. ¹⁴International Institute for Applied Systems Analysis (IIASA), Vienna, Austria. ¹⁵University of Oxford, Oxford, UK. ¹⁶Environmental Defence Fund, New York, NY, USA. ¹⁷Princeton University, Princeton, NJ, USA. ¹⁸Imperial College, London, UK. ¹⁹Duke University, Durham, NC, USA. ✉email: myles.allen@ouce.ox.ac.uk

gradual adjustment to average SLCF forcing¹⁶, all evaluated over the same time-interval.

Hence a separate indication of the contributions of LLCFs and SLCFs in emission targets, or equivalently the LLCF contribution to total CO₂-e₁₀₀ emissions, is required to allow for the global temperature outcome to be calculated relatively unambiguously. It is important to note, however, that the evaluation of emission targets at the national or corporate level cannot be undertaken from a physical science perspective alone, but also depends on economic, social, equity and political considerations^{2–5,17}, including responsibility for past warming, capacity for and costs of abatement, and non-climate impacts. Separate specification would also facilitate the use of alternate or flexible emission metrics, which may be useful for achieving a cost-effective emission trajectory over time¹⁸ or addressing specific policy goals such as limiting near-term rates of warming¹⁹. Indicative contributions from LLCF and SLCF abatement would not preclude trade-offs between them, but would clarify the need to monitor the temperature impacts of any such trade-offs over a range of timescales²⁰.

It has long been accepted²¹ that stringent mitigation of both LLCFs and SLCFs is needed to meet any ambitious temperature goal, but making progress on two fronts necessitates monitoring progress on two fronts. Some countries (but very few companies) already specify the contribution of LLCFs and/or SLCFs to total CO₂-e₁₀₀ emissions in NDCs, LT-LEDs and science-based targets (<https://sciencebasedtargets.org/>) communicated under the Greenhouse Gas Protocol. Quantifying the aggregated implications of these targets for future global temperature simply requires a much wider uptake of this practice, representing a simple and achievable innovation that would enhance the transparency of any stocktake of progress towards any global temperature outcome. Separate indication of LLCF and/or SLCF contributions could be communicated by countries as additional information consistent with Decision 4/CMA.1. This does not have to affect any existing or planned NDCs or long-term net zero strategies²² communicated using aggregate CO₂-e₁₀₀.

WHY SEPARATE SPECIFICATION IS SO USEFUL

To quantify the SR1.5 and AR6 statements quoted above, human-induced global temperature change over a multi-decade time-interval Δt , relative to the level of human-induced warming at the beginning of that interval (e.g. the present day or pre-industrial), can be decomposed using the framework articulated above as follows:

$$\Delta T = \kappa_E \bar{E}_C \Delta t + \kappa_F (\Delta F_N + \rho \bar{F}_N \Delta t), \quad (1)$$

where \bar{E}_C and \bar{F}_N are globally aggregated average CO₂ emission-rates and non-CO₂ radiative forcing, respectively (so $\bar{E}_C \Delta t$ is cumulative CO₂ emissions), and ΔF_N is the change in decadal-average non-CO₂ forcing, all evaluated over that interval (the geophysical “Zero Emissions Commitment” is expected to be relatively small over a multi-decade time-interval²³, but this may not be the case on longer timescales). The coefficients κ_E (the TCRE) and κ_F (the TCRF, or “fast” component of the climate response to any forcing change, denoted c_1 in ref. ¹², or sum of fast components²⁴: see supplementary material), are both scenario-independent in the absence of strongly non-linear carbon cycle feedbacks or climate response. The only scenario-dependent coefficient is ρ , the fractional Rate of Adjustment to Constant Forcing (RACF), or the relatively small fractional rate at which forcing needs to decline to maintain stable temperatures. It depends on how fast and how recently F_N has increased (this term represents the delayed adjustment to past forcing increases, so is larger for more recent and rapid increases). If F_N varies only on multi-decadal timescales, $\rho = c_2 / (\kappa_F s_2)$, where c_2 is the “slow” (multi-century) component of the climate sensitivity, and s_2 the

deep ocean thermal adjustment timescale. For representative¹² coefficient values, $\rho \leq 0.3\%$ per year, making this third term usually small.

Aggregate CO₂-e₁₀₀ emissions cannot be used to calculate F_N if these comprise a mixture of LLCFs and SLCFs. Aggregate CO₂-e₁₀₀ emissions of LLCFs, E_L , can, however, be combined unambiguously and have the same impact on global temperature on decade to century timescales as the corresponding quantity of CO₂. Likewise, aggregate CO₂-e₁₀₀ emissions of SLCFs, E_S , multiplied by the AGWP₁₀₀ of CO₂, A_{100} , give SLCF radiative forcing, F_S (A_{100} normally includes a first-order estimate of the impact of carbon cycle feedbacks²⁵ so, for consistency, this should also be included in the GWP₁₀₀ values used to compute E_S).

For emissions reported as CO₂-e₁₀₀ the above expression can therefore be re-written (now grouping all LLCFs with CO₂):

$$\Delta T = \kappa_E \bar{E}_L \Delta t + \kappa_F (\Delta F_S + \rho \bar{F}_S \Delta t), \quad (2)$$

or equivalently, using $F_S = A_{100} E_S$ on multi-decadal timescales,

$$\Delta T = \kappa_E \bar{E}_L \Delta t + \kappa_F A_{100} (\Delta E_S + \rho \bar{E}_S \Delta t). \quad (3)$$

Hence ΔT can be estimated directly using well-known (albeit uncertain) climate system properties if, and only if, total CO₂-e₁₀₀ emissions of long-lived climate forcers, E_L , are specified in emission targets together with total CO₂-e₁₀₀ emissions, $E_L + E_S$; or, equivalently, E_L and E_S are specified separately. ΔT cannot be calculated from the sum of $E_L + E_S$ alone.

This is illustrated by Fig. 1, which shows the impact of LLCF and SLCF emissions, expressed as CO₂-e₁₀₀, on global temperature change over a multi-decade period, relative to the level of warming at the beginning of that period, calculated with a simple climate model¹². Stylised cases of constant (darker shades) and step-change (+10%, lighter shades, and –50%, dotted lines) emissions are shown in panels a and c. Warming due to LLCF emissions (the term $\kappa_E \bar{E}_L \Delta t$ in Eq. (3)) increases linearly with cumulative emissions in all three cases (panel b). Warming due to an ongoing constant emission of an SLCF that started decades before the beginning of this period (the $\kappa_F A_{100} \rho \bar{E}_S \Delta t$ term) also increases linearly (panel d, darker blue) but at a slower rate per tCO₂-e₁₀₀ emitted (by a factor of about 4, because $\kappa_E \approx 4 \times \kappa_F A_{100} \rho$): global temperatures have already partially equilibrated with this constant emission (by how much depends on how long ago these SLCF emissions began, which is why ρ is the only scenario-dependent coefficient in these expressions). Finally, warming due to an increase in SLCF emissions (the $\kappa_F A_{100} \Delta E_S$ term, panel d, lighter blue) is 4–5 times greater than would be expected from the same increase in tCO₂-e₁₀₀ emissions of an LLCF (panel b, lighter red) over the 20 years following the increase ($\kappa_F A_{100} \approx 4.5 \times \kappa_E \times 20$ years). Hence the AR6 statement “expressing methane emissions as CO₂ equivalent emissions using GWP₁₀₀ overstates the effect of constant methane emissions on global surface temperature by a factor of 3–4 ... while understating the effect of any new methane emission source by a factor of 4–5 over the 20 years following the introduction of the new source”²⁶ applies to the impact of global emissions of any SLCF. Any decrease in SLCF emissions also has a much greater impact on temperatures over a multi-decade period per tCO₂-e₁₀₀ avoided than a corresponding decrease in LLCF emissions (red and blue dotted lines) (Fig. 1).

Temperature changes in the figure are calculated using a particular model, LLCF, SLCF and scenario. The figure would, however, appear similar if another model, combination of gases or scenario of prior emissions were used, provided emissions do not change rapidly immediately before the beginning or end of the period shown, because the relationship between emissions and warming expressed in Eq. (3) is generic. Individual terms in Eq. (3), assuming constant coefficients, are shown by the arrows on the right of panels




Fig. 1 Stylised LLCF and SLCF emissions and resulting global temperature change ΔT over a multi-decade period. Darker bands in panels **a** and **c** show, respectively, constant LLCF and SLCF emissions of 1 $\text{tCO}_2\text{-e}_{100}$ per year starting some decades before the interval shown. Pale bands show a 10% increase one-quarter of the way through the interval shown, while dotted lines show a 50% decrease. Resulting temperature changes relative to the start of this interval shown in panels **b** and **d**, calculated using a simple climate model: vertical axes in **b** and **d** are scaled identically to illustrate smaller rate of warming due to constant SLCF emissions and much larger warming impact of any change in SLCF emissions relative to the warming due to identical $\text{CO}_2\text{-e}_{100}$ LLCF emissions. Vertical arrows in the right show predicted contributions to ΔT from the individual terms in Eq. (3): three arrows in panel **b** show cumulative LLCF emissions over this interval multiplied by the TCRE for the three scenarios shown; the lower and upper arrows in panel **d** show, respectively, the predicted warming due to ongoing constant SLCF emissions and additional warming due to the 10% increase. The figure illustrates that Eq. (3) allows reliable, if approximate, prediction of multi-decade warming ΔT if, and only if, LLCF and SLCF emissions are specified separately.

b and **d**. These match the warming calculated by the explicit simple climate model within modelling uncertainties. The figure shows temperature change relative to the start of the period rather than absolute warming because the latter is not determined by Eq. (3) but depends on the prior LLCF and SLCF emissions history (the specific scenario used to generate this figure is shown in full in the Supplementary Information).

Temperature change ΔT over a multi-decade period depends, to first order, only on cumulative emissions of LLCFs $\bar{E}_L \Delta t$, cumulative emissions of SLCFs $\bar{E}_S \Delta t$, and net change in total SLCF emission rates ΔE_S , over that period alone. As the SR1.5 and AR6 emphasised, future warming depends on future emissions. Making use of this information, however, requires both E_L and E_S to be

specified: only specifying the sum $E_L + E_S$ introduces an ambiguity in temperature outcome.

Separate specification also facilitates assessing the implications of different metrics. For example, aggregate CO_2 -equivalent emissions using the 20-year Global Warming Potential (GWP_{20}) can be approximated by $E_L + 3E_S$ if both E_L and E_S are reported as $\text{CO}_2\text{-e}_{100}$, with a slightly higher multiplicative factor (up to 4) if E_S is dominated by forcers with lifetimes of order one year (Table 8.A.1 of ref. ¹² shows that GWP_{20} values are similar to GWP_{100} values for LLCFs and 3 or 4 times GWP_{100} values for gases with lifetimes of order a decade or a year, respectively). Finally, we re-emphasise that these expressions capture our physical understanding of how global emissions of LLCFs and SLCFs collectively determine global

temperature change, and illustrate the utility of separate specification of E_L and E_S . How this understanding is used to inform the assessment of the adequacy of individual emission targets depends on other considerations listed above and cannot be argued from a physical science perspective alone. There will be several other advantages to the additional communication such as being able to estimate air quality co-benefits of mitigation.

DATA AND CODE AVAILABILITY

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study. A self-contained Python notebook to reproduce the figure is provided on https://gitlab.ouce.ox.ac.uk/OMP_climate_pollutants/separate-contributions.

Received: 24 September 2021; Accepted: 6 December 2021;

Published online: 28 January 2022

REFERENCES

- UNFCCC, https://unfccc.int/sites/default/files/resource/cma2018_3_add2_new_advance.pdf (2018).
- Rogelj, J. & Schleussner, C.-F. Unintentional unfairness when applying new greenhouse gas emissions metrics at country level. *Environ. Res. Lett.* **14**, 114039 (2019).
- Cain, M. et al. Comment on 'Unintentional unfairness when applying new greenhouse gas emissions metrics at country level'. *Environ. Res. Lett.* **16**, 068001 (2021).
- Rogelj, J. & Schleussner, C.-F. Reply to Comment on 'Unintentional unfairness when applying new greenhouse gas emissions metrics at country level'. *Environ. Res. Lett.* **16**, 068002 (2021).
- Schleussner, C.-F., Nauels, A., Schaeffer, M., Hare, W. & Rogelj, J. Inconsistencies when applying novel metrics for emissions accounting to the Paris agreement. *Environ. Res. Lett.* **14**, 124055 (2019).
- IPCC, 2018. In *Global Warming of 1.5 °C* (eds. Masson-Delmotte, V. et al.) 32 (World Meteorological Organization, 2018).
- IPCC, 2021. In *Climate Change 2021, the Physical Science Basis* (eds. Masson-Delmotte, V. et al.) (Cambridge University Press, 2021).
- Tanaka, K. & O'Neill, B. C. The Paris Agreement zero-emissions goal is not always consistent with the 1.5 °C and 2 °C temperature targets. *Nat. Clim. Change* **8**, 319–324 (2018).
- Denison, S., Forster, P. M. & Smith, C. J. Guidance on emissions metrics for nationally determined contributions under the Paris Agreement. *Environ. Res. Lett.* **14**, 124002 (2019).
- Smith, S. et al. Equivalence of greenhouse-gas emissions for peak temperature limits. *Nat. Clim. Change* **2**, 535–538 (2012).
- Shindell, D. et al. A climate policy pathway for near- and long-term benefits. *Science* **356**, 493–494 (2017).
- Myhre, G. et al. Anthropogenic and Natural Radiative Forcing. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Stocker, T. F. et al.) Ch. 8 (2013).
- Shine, K. P. et al. Alternatives to the global warming potential for comparing climate impacts of emissions of greenhouse gases. *Clim. Change* **68**, 281–302 (2005).
- Gregory, J. M. et al. Quantifying carbon cycle feedbacks. *J. Clim.* **22**, 5232–5250 (2009).
- Held, I. M. et al. Probing the fast and slow components of global warming by returning abruptly to preindustrial forcing. *J. Clim.* **23**, 2418–2427 (2010).
- Cain, M. et al. Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *npj Clim. Atmos. Sci.* **2**, 29 (2019).
- Rajamani, L. et al. National 'fair shares' in reducing greenhouse gas emissions within the principled framework of international environmental law. *Clim. Policy*, <https://doi.org/10.1080/14693062.2021.1970504> (2021).
- Tanaka, K. et al. Cost-effective implementation of the Paris Agreement using flexible greenhouse gas metrics. *Sci. Adv.* **7**, eabf9020 (2021).
- Ocko, I. et al. Unmask temporal trade-offs in climate policy debates. *Science* **356**, 492–493 (2017).
- Allen, M. R. et al. Ensuring that offsets and other internationally transferred mitigation outcomes contribute effectively to limiting global warming. *Environ. Res. Lett.* **16**, 074009 (2021).
- Shindell, D. et al. Simultaneously mitigating near-term climate change and improving human health and food security. *Science* **335**, 183–189 (2012).
- Rogelj, J., Geden, O., Cowie, A. & Reisinger, A. Net-zero emissions targets are vague: three ways to fix. *Nature* **591**, 365–368 (2021).
- MacDougall, A. H. et al. Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂. *Biogeosciences* **17**, 2987–3016 (2020).
- Tsutsui, J. Quantification of temperature response to CO₂ forcing in atmosphere-ocean general circulation models. *Climatic Change* **140**, 287–305 (2017).
- Gasser, T. et al. Accounting for the climate-carbon feedback in emission metrics. *Earth Syst. Dynam.* **8**, 235–253 (2017).
- Forster, P. et al. In *Climate Change 2021, the Physical Science Basis* (eds. Masson-Delmotte, V. et al.) Ch. 7 (Cambridge University Press, 2021).

ACKNOWLEDGEMENTS

The authors acknowledge support from the FORCeS, 4C and CONSTRAIN projects funded by the European Union's Horizon 2020 research and innovation program under Grant Agreement Nos. 821205, 821003, 820829, respectively; the Wellcome Trust, Our Planet Our Health (Livestock, Environment and People), award number 205212/Z/16/Z; the ESA Earth Observation Programme; and the University of Oxford's Strategic Research Fund.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41612-021-00226-2>.

Correspondence and requests for materials should be addressed to Myles R. Allen.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

Indicate separate contributions of long-lived and short-lived greenhouse gases in emission targets

Allen, Myles R.

2022-01-28

Attribution 4.0 International

Allen MR, Peters GP, Shine KP, et al., (2022) Indicate separate contributions of long-lived and short-lived greenhouse gases in emission targets, *npj Climate and Atmospheric Science*, Volume 5, January 2022, Article number 5

<https://doi.org/10.1038/s41612-021-00226-2>

Downloaded from CERES Research Repository, Cranfield University