

Short communication

On the origin of carbon dioxide released from rewetted soils



F.C. Fraser ^{a,*}, R. Corstanje ^a, L.K. Deeks ^a, J.A. Harris ^a, M. Pawlett ^a, L.C. Todman ^b,
A.P. Whitmore ^b, K. Ritz ^c

^a School of Energy, Environment, and Agrifood, Cranfield University, Bedford, UK

^b Rothamsted Research, Harpenden, UK

^c School of Biosciences, University of Nottingham, Nottingham, UK

ARTICLE INFO

Article history:

Received 22 December 2015

Received in revised form

24 June 2016

Accepted 28 June 2016

Keywords:

Birch effect

Dry:wet cycles

CO₂ flux

Extracellular oxidative metabolism

Soil sterilisation

ABSTRACT

When dry soils are rewetted a pulse of CO₂ is invariably released, and whilst this phenomenon has been studied for decades, the precise origins of this CO₂ remain obscure. We postulate that it could be of chemical (i.e. via abiotic pathways), biochemical (via free enzymes) or biological (via intact cells) origin. To elucidate the relative contributions of the pathways, dry soils were either sterilised (double autoclaving) or treated with solutions of inhibitors (15% trichloroacetic acid or 1% silver nitrate) targeting the different modes. The rapidity of CO₂ release from the soils after the drying:rewetting (DRW) cycle was remarkable, with maximal rates of evolution within 6 min, and 41% of the total efflux over 96 h released within the first 24 h. The complete cessation of CO₂ efflux following sterilisation showed there was no abiotic (dissolution of carbonates) contribution to the CO₂ release on rewetting, and clear evidence for an organismal or biochemical basis to the flush. Rehydration in the presence of inhibitors indicated that there were approximately equal contributions from biochemical (outside membranes) and organismal (inside membranes) sources within the first 24 h after rewetting. This suggests that some of the flux was derived from microbial respiration, whilst the remainder was a consequence of enzyme activity, possibly through remnant respiratory pathways in the debris of dead cells.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Rewetting of a dry soil invariably causes a large flux of carbon dioxide (CO₂) to be rapidly released, which is sometimes referred to as the Birch effect (Birch, 1958, 1960). This phenomenon has been observed both in laboratory incubations (Kieft et al., 1987; Unger et al., 2010; Shi and Marschner, 2014) and in field circumstances using closed chambers (Yan et al., 2014) or eddy covariance towers (Xu et al., 2004). These fluxes have been observed across a wide range of ecotypes (Jarvis et al., 2007; Thomas and Hoon, 2010; Sugihara et al., 2015), but are particularly significant in dryland and Mediterranean ecosystems where they can make up a significant proportion of soil C-emissions (Lee et al., 2004; Hunt et al., 2004; Brito et al., 2013). These drying:rewetting (DRW) induced CO₂ efflux events can even significantly reduce the annual net C gain in Mediterranean forests (Jarvis et al., 2007).

Several theories have been proposed to explain this phenomenon including: (i) the exposure of physically-protected organic matter to microbial metabolism via aggregate dispersion on rewetting (Denef et al., 2001; Wu and Brookes, 2005; Xiang et al.,

2008); (ii) microbial necromass increasing the supply of readily assimilable substrate to the surviving microbial populations (Kieft et al., 1987; Van Gestel et al., 1992; Blazewicz et al., 2013); (iii) increases in the supply of labile organic matter due to the rapid release, on rewetting, of intra-cellular solutes previously concentrated within microbial cells to maintain osmotic balance in response to dehydration (Halverson et al., 2000; Warren, 2014); and (iv) a supply of labile organic C is built up during the dry period prior to rewetting and subsequently quickly metabolised on rewetting. There is a known uncoupling of rates of CO₂ efflux and detectable microbial growth rates after a DRW cycle (Iovieno and Bååth, 2008; Meisner et al., 2015) and microbial populations in such circumstances show little change in their net size (Fierer and Schimel, 2002). However, recent work by Blazewicz et al. (2013) show that despite their unchanging size these populations turnover rapidly in response to a DRW cycle. They also suggest that more cellular derived organic-C is available in soil samples than is turned over in the initial phases after rewetting. This organic-C will contain cellular material including constituents of enzymatic pathways – remnant respiratory pathways – with the potential to carry out reactions leading to CO₂ efflux. Thus it is possible that CO₂

* Corresponding author.

E-mail address: f.c.fraser@cranfield.ac.uk (F.C. Fraser).

release from re-wetted soils is not exclusively derived from respiration pathways occurring in intact microbes. There are also reports of over-estimation of soil respiration rates due to contributions of CO₂ from dissolution of soil carbonates; however, reports are inconsistent and range from 1 to 2% up to 74% of CO₂ efflux from soil being attributed to carbonate dissolution (Biasi et al., 2008; Ramnarine et al., 2012; Schindlbacher et al., 2015). It is as yet unclear how the DRW process may affect carbonate dissolution from soils although Tamir et al. (2011) found that in highly calcareous soils the rate of inorganic CO₂ production was lower in drier samples. However, it is also known that increases in soil OM content can alter the balance of pH, as a result of increased nitrification rates, leading to increase dissolution of carbonates (Tamir et al., 2013). As such an increase in available OM as a result of any of the 4 processes described above (aggregate dispersion, increased necromass, release of intracellular-solutes, or accumulation of labile organic matter) could potentially lead to this phenomenon on rewetting, and an abiotic route to CO₂ production must also be considered.

On this basis we posit that there are three potential sources of CO₂, all of which could contribute to the efflux on rewetting: (i) *abiotic* via carbonate dissolution (Shanhun et al., 2012); (ii) *biochemical*, involving the release of CO₂ from organic matter outside cell membranes and mediated by free or residually-bound enzymes (Maire et al., 2013) (Blankinship et al., 2014); (iii) *organismal*, i.e. microbial respiration via the Krebs cycle carried out within intact organelles or cells (Fig. 1). One potential way to determine the relative contribution of these sources is to probe the phenomenon in soils treated in various ways to block certain of the pathways involved, such as via complete sterilisation (i.e. any form of biochemical or organismal pathway), or to spike the rehydration water with various forms of metabolic inhibitors (i.e. to distinguish biochemical from organismal). We hypothesised that i) the majority of CO₂ released is derived from an organismal source, and hence that CO₂ efflux upon rehydration would be curtailed where organismal pathways were blocked and ii) there would be no significant contribution to the total CO₂ efflux of CO₂ from an abiotic source.

Soils were collected from the top 15 cm of 4 long-term grassland sites in May 2015 (soil parameters shown in Table 1); all soils were sieved to pass a 2 mm mesh, adjusted to 45% water holding capacity (WHC) and pre-incubated at 25 °C for 7 days. Aliquots of the soils (1 g; 3 replicates of each soil) were then exposed to 4 DRW cycles over 28 days, where each cycle consisted of 3 days drying followed by rewetting to 45% WHC using sterile, deionised water. Drying was

standardised by locating the soils in a sealed container in the presence of silica gel. Aliquots of 1.0 g of soil were adopted in order to ensure that penetration of water throughout the soil volume would be rapid. The time-course of CO₂ evolution at 6-min intervals following rewetting was determined independently for each replicate using an automated multi-channel conductimetric respirometer (RABIT, Don Whitley, Shipley, UK; (Butler et al., 2012), for 5 days. To account for any background variation in CO₂ efflux blanks were run alongside soil samples; this involved measuring the signal from empty, sealed cells.

Another set of three replicates was subjected to a further range of treatments, viz. (i) 'Live controls' - involving no sterilisation, DRW as described above; (ii) 'Moist controls' - also unsterilized but with 0.2 mL sterile, deionised water added prior to exposure to DRW - this is a procedural control to account for the fact that liquid was added to the sample prior to drying as described above; (iii) 'Autoclaved', where samples were autoclaved twice at 121 °C at 3.1 bar for 20 min with a 24 h pause between (Systec 3150 EL, Linden, Germany); (iv) 'TCA', with 0.2 mL of 15% trichloroacetic acid (TCA) addition; (v) 'AgNO₃', with 0.2 mL of 1% silver nitrate addition. All amendments and autoclaving were carried out prior to the DRW process described above. The rationale for these treatments (Fig. 1) is that autoclaving would prevent all organismal or biochemical activity by denaturing all proteins - in this circumstance any CO₂ produced would be via abiotic pathways. TCA (15%) would precipitate proteins, including extracellular enzymes (Ladd and Butler, 1972) and as such remove any biochemical source of CO₂. The mechanism of protein precipitation by TCA is unclear but is likely to be due to protein unfolding (Rajalingam et al., 2009) and as such may also affect microbial membranes. AgNO₃ is a known antiseptic and so kills microbes; the precise mode of action is surprisingly poorly understood but the Ag⁺ ions are known to cause physical damage to cells and DNA - separation of cytoplasmic membranes from cell walls and condensing of DNA in both *Escherichia coli* and *Staphylococcus aureus* (Feng et al., 2000). Silver and other heavy metals are also known to bind to thiol groups in proteins resulting in their inactivation (Liau et al., 1997). They also interfere with intra-cellular processes and membranes/cell walls therefore AgNO₃ may also affect some extracellular enzymes (e.g. thiol-proteases). This treatment is designed to primarily inhibit the organismal pathway but is likely to have a lesser effect on biochemical mechanisms - i.e. extracellular enzymes (Fig. 1). Whilst the extent to which these inhibitors operate exclusively on these pathways is unknown (and may be impossible to precisely establish), the rationale is that they will be at least partly informative. However, autoclaving twice unequivocally sterilises soil.

The rapidity of CO₂ release from the soils after the DRW cycle was remarkable, in that we detected maximal rates of evolution after 6 min, and never captured the actual peak as such, only a downward trend from a presumed peak (Fig. 2). Within the first hour following wet-up an average of 5% of the total CO₂ efflux over 96 h was observed and of this approximately 24% occurred within the first 12 min (Fig. 2a–d). Of the total CO₂ efflux measured over 96 h after rewetting, an average of 41% was measured in the first 24 h (Fig. 2e–h); this consistency of effect with - where the same proportion of CO₂ was measured in the first 24 h after each of a series of rewetting events - was also observed by Birch (1958).

A large difference in CO₂ release on rewetting between the wet control and the standard response to DRW was manifest (Fig. 3a). This is likely because the 3-day drying period resulted in different amounts of moisture loss between treatments; those exposed to the prescribed DRW cycle lost 34% of their mass on average over the 3 days of drying, however, the moist controls lost only 16% of their mass on average. This shows that soil dried to a greater extent will give a larger flush of CO₂ on rewetting than a sample of the same




Fig. 1. Three potential sources of CO₂ to account for the flush on rewetting of dry soils and the treatments used to identify the respective contributions of these. Light grey bars in lower panel indicates which potential sources of CO₂ are uninhibited by each treatment, mid-grey shows which sources are potentially inhibited, and dark grey shows those that are 'switched off' by the different treatments.

Table 1
Locations from which soils sampled (latitude and longitude) and associated basic properties.

| Soil | Latitude: Longitude | Sand (%) | Silt (%) | Clay (%) | N (%) | C (%) | C:N | pH | Water-holding capacity (ml g ⁻¹) | Loss on ignition (g g soil ⁻¹) | Microbial biomass C(μg g ⁻¹) | Inorganic C-content (%) |
|------|----------------------|----------|----------|----------|-------|-------|------|-----|--|--|--|-------------------------|
| A | 52.4245°N: -4.0652°W | 7.5 | 53.9 | 38.6 | 0.7 | 7.6 | 10.8 | 5.5 | 0.98 | 0.150 | 2330 | 0.18 |
| B | 53.2222°N: -4.0132°W | 28.7 | 41.8 | 29.4 | 0.8 | 9.5 | 11.4 | 5.1 | 0.10 | 0.169 | 1699 | 0.74 |
| C | 53.0412°N: -4.0445°W | 34.8 | 48.9 | 16.3 | 0.6 | 6.1 | 10.8 | 5.8 | 0.91 | 0.127 | 1407 | 0.74 |
| D | 52.9988°N: -4.4290°W | 75.3 | 24.4 | 0.3 | 0.3 | 3.4 | 10.7 | 5.6 | 0.70 | 0.065 | 739 | 0.26 |




Fig. 2. CO₂ release profiles from unsterilized grassland soil exposed to 4 repeated DRW events (Cycles 1–4); (a–d) CO₂ release measured at 6 min intervals in the first hour after rewetting, (e–h) hourly CO₂ release over the first 24 h after rewetting, (i–l) hourly CO₂ release over the entire 94 h wet period. Means (n = 3) indicated by black line surrounded by confidence bands of ±1 standard error.




Fig. 3. CO₂ efflux rates following rewetting of a dry soil with various solutions; (a) live soil (green) exposed to a DRW cycle compared to all other treatments including a moist control (blue), area outlined in red is shown in greater detail in (b); (b) amplification of y-axis from (a), i.e. CO₂ efflux following a DRW cycle from the moist control (blue), blanks (no soil - brown), autoclaved (orange), 15% TCA (purple) and 1% silver nitrate (grey) treated soils. Lines show mean rates of CO₂ efflux (n = 12 (3 reps each of 4 soils)) surrounded by confidence bands of ±1 standard error.

soil dried less severely (Kieft et al., 1987; Fierer and Schimel, 2002; Unger et al., 2010; Meisner et al., 2015). Those samples treated with 15% TCA and 1% AgNO₃ dried to a greater extent over 3 days than the moist controls (21 and 28% mass loss respectively) and those that were autoclaved lost 45% of their mass on average. Despite these large differences in moisture loss between the moist controls and the inhibitor treated samples (both TCA and AgNO₃) the effect of moisture loss on total CO₂ efflux was found to be non-significant using an analysis of covariance (ANCOVA; p = 0.71), nor was there a significant interaction between inhibitor treatment and moisture loss (p = 0.25). As such, the main effect of inhibitor treatment can be interpreted directly.

Hereafter, responses of inhibitor-treated samples to DRW are compared to that of the moist controls (Fig. 3b). Autoclaving effectively ‘switched off’ CO₂ production after a rewetting event (total CO₂ efflux over 24 h was significantly different between water controls and autoclaved samples and autoclaved totals were not significantly different from blanks (p = 0.01, p = 0.99 respectively, Fig. 3). A preliminary experiment using soil with higher CaCO₃ contents (0.93% compared to 0.48% on average for soils listed in Table 1) showed the same lack of activity after autoclaving and a DRW event (data not presented). These results show that there was effectively no chemical contribution to the CO₂ flush observed after rewetting in these soils. This is in contrast to observations made in

some calcareous, arid soils where CO₂ derived from inorganic–C has been observed to account for 30–75% of the total soil CO₂ efflux (Tamir et al., 2011; Shanahun et al., 2012). As previously stated, these observations have also been made in temperate soils but results are scarce and inconsistent with ranges of 1–2% (Schindlbacher et al., 2015), to 50% (Biasi et al., 2008) all the way up to 74% (Ramnarine et al., 2012) of the total CO₂ flux attributable to inorganic C sources. Notably, none of these studies examined the response to a DRW event although Biasi et al. (2008) noted an effect of water addition in the laboratory. The effect of autoclaving observed in our study is therefore strong evidence for an organismal and/or biochemical origin for the evolved CO₂.

Treating soils with either 15% TCA or 1% AgNO₃ substantially reduced but did not eliminate CO₂ production, compared to the moist control, following a DRW event (Fig. 3b). Inhibition of CO₂ evolution by AgNO₃ was greater than by TCA for the latter half of the measurement period (Fig. 3), although the accumulated total release was not statistically significant in the case of these two inhibitors ($p = 0.98$). This suggests that a greater portion of the CO₂ measured after a DRW event is derived from the organismal pathway. This effect appeared to increase over time with the amount of CO₂ produced hourly by AgNO₃ treated soils decreasing more rapidly over the first 24 h than it did for TCA treated soils this is exemplified by the increasing gap between the confidence bands for AgNO₃ and TCA treated soils after approximately 13 h of incubation in Fig. 3. It is commonly assumed that the majority of CO₂ measured after a DRW event is derived from the organismal pathway, and the effect of AgNO₃ would certainly suggest this. There was also a substantial reduction in CO₂, compared to the moist control, due to the addition of TCA, which suggests that an additional contribution to the CO₂ flux after the DRW event was via the biochemical route. This is consistent with the findings of Maire et al. (2013) who report a 16–48% contribution of an extracellular oxidative metabolism pathway, termed 'EXOMET', to soil CO₂ flux. Blankinship et al. (2014) found only a 26–47% reduction in CO₂ emission from intermediates in the TCA cycle after sterilisation suggesting that these enzymes are still active when cells are dead but not completely dispersed, again noting that neither of these two studies were in response to DRW events. It is known that many enzymes are stable in the soil environment on a long term basis (Burns et al., 2013). Such stability is generally achieved by adsorption onto soil colloids or incorporation with humic complexes (Nannipieri et al., 1996). The effects of adsorption or humic complexing can include inhibition and steric hindrance which can cause a reduction in potential activity of this sizeable enzyme pool by up to 90% (Quiquampoix et al., 2002). If even a small proportion of these enzymes were to be brought into solution after rewetting this could have a large effect on the levels of activity in soils (Stursova and Sinsabaugh, 2008). Significant increases in rates of enzyme activity have been recorded in soils exposed to DRW both during laboratory preparation (Kandeler and Gerber, 1988) and as a result of environmental conditions (Hinojosa et al., 2004) suggesting that portions of the adsorbed enzyme pool are solubilised by the process of rewetting after drying increasing the potential for a biochemically driven response in DRW soils.

Our results demonstrate the apparent immediacy of the Birch effect, and go some way to explaining the pathways by which the CO₂ is evolved, viz. primarily organismal but with a potentially large contribution from the biochemical pathways. We note that for our experiments, these are roughly equivalent in magnitude. Thus we reject the hypothesis that the origin of the CO₂ released following rehydration is predominantly organismal. We have shown that in these temperate soils, unlike in more calcareous, arid systems, there is no contribution of carbonate dissolution even when the intrinsic concentration of CaCO₃ is high. This means that this

effectively instantaneous release of CO₂ is governed by the soil biota. We have shown evidence that not only are intact microbial cells apparently capable of reinstating their high rates of respiration within minutes following rehydration after 3 days of drying, but also that there is a potentially extensive contribution of CO₂ from remnant enzymatic pathways outside of cell membranes.

Acknowledgements

The authors would like to thank the two anonymous reviewers for their insightful comments. This work was supported by the UK Biotechnology and Biological Research Council research grants BB/J000396/1, BB/J000396/2 and BB/J000671/1 (partly supported by UK Department of Environment, Food and Rural Affairs).

References

- Biasi, C., Lind, S.E., Pekkarinen, N.M., Huttunen, J.T., Shurpali, N.J., Hyvönen, N.P., Repo, M.E., Martikainen, P.J., 2008. Direct experimental evidence for the contribution of lime to CO₂ release from managed peat soil. *Soil Biol. Biochem.* 40, 2660–2669.
- Birch, H.F., 1960. Nitrification in soils after different periods of dryness. *Plant Soil* 12, 81–96.
- Birch, H.F., 1958. Pattern of humus decomposition in east african soils. *Nature* 181, 788–788.
- Blankinship, J.C., Becerra, C.A., Schaeffer, S.M., Schimel, J.P., 2014. Separating cellular metabolism from exoenzyme activity in soil organic matter decomposition. *Soil Biol. Biochem.* 71, 68–75.
- Blazewicz, S.J., Schwartz, E., Firestone, M.K., 2013. Growth and death of bacteria and fungi underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. *Ecology* 95, 1162–1172.
- Brito, P., Trujillo, J.L., Morales, D., Jiménez, M.S., Wieser, G., 2013. response of soil CO₂ efflux to simulated precipitation pulses in a Canary Island pine forest at Treeline. *Arid Land Res. Manag.* 27, 178–187.
- Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein, M.D., Weintraub, M.N., Zoppini, A., 2013. Soil enzymes in a changing environment: current knowledge and future directions. *Soil Biol. Biochem.* 58, 216–234.
- Butler, E., Whelan, M.J., Ritz, K., Sakrabani, R., van Egmond, R., 2012. The effect of triclosan on microbial community structure in three soils. *Chemosphere* 89, 1–9.
- Denef, K., Six, J., Paustian, K., Merckx, R., 2001. Importance of macroaggregate dynamics in controlling soil carbon stabilization: short-term effects of physical disturbance induced by dry–wet cycles. *Soil Biol. Biochem.* 33, 2145–2153.
- Feng, Q.L., Wu, J., Chen, G.Q., Cui, F.Z., Kim, T.N., Kim, J.O., 2000. A mechanistic study of the antibacterial effect of silver ions on *Escherichia coli* and *Staphylococcus aureus*. *J. Biomed. Mater. Res.* 52, 662–668.
- Fierer, N., Schimel, J.P., 2002. Effects of drying–rewetting frequency on soil carbon and nitrogen transformations. *Soil Biol. Biochem.* 34, 777–787.
- Halverson, L.J., Jones, T.M., Firestone, M.K., 2000. Release of intracellular solutes by four soil bacteria exposed to dilution stress. *Soil Sci. Soc. Am. J.* 64, 1630.
- Hinojosa, M.B., Carreira, J.A., García-Ruiz, R., Dick, R.P., 2004. Soil moisture pre-treatment effects on enzyme activities as indicators of heavy metal-contaminated and reclaimed soils. *Soil Biol. Biochem.* 36, 1559–1568. *Enzymes in the Environment: Activity, Ecology and Applications*.
- Hunt, J.E., Kelliher, F.M., McSeveny, T.M., Ross, D.J., Whitehead, D., 2004. Long-term carbon exchange in a sparse, seasonally dry tussock grassland. *Glob. Change Biol.* 10, 1785–1800.
- Iovieno, P., Bååth, E., 2008. Effect of drying and rewetting on bacterial growth rates in soil. *FEMS Microbiol. Ecol.* 65, 400–407.
- Jarvis, P., Rey, A., Petsikos, C., Wingate, L., Rayment, M., Pereira, J., Banza, J., David, J., Miglietta, F., Borghetti, M., Manca, G., Valentini, R., 2007. Drying and wetting of Mediterranean soils stimulates decomposition and carbon dioxide emission: the "Birch effect". *Tree Physiol.* 27, 929–940.
- Kandeler, E., Gerber, H., 1988. Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biol. Fertil. Soils* 6, 68–72.
- Kieft, T.L., Soroker, E., Firestone, M.K., 1987. Microbial biomass response to a rapid increase in water potential when dry soil is wetted. *Soil Biol. Biochem.* 19, 119–126.
- Ladd, J.N., Butler, J.H.A., 1972. Short-term assays of soil proteolytic enzyme activities using proteins and dipeptide derivatives as substrates. *Soil Biol. Biochem.* 4, 19–30.
- Lee, X., Wu, H.-J., Sigler, J., Oishi, C., Siccama, T., 2004. Rapid and transient response of soil respiration to rain. *Glob. Change Biol.* 10, 1017–1026.
- Liau, S.Y., Read, D.C., Pugh, W.J., Furr, J.R., Russell, A.D., 1997. Interaction of silver nitrate with readily identifiable groups: relationship to the antibacterial action of silver ions. *Lett. Appl. Microbiol.* 25, 279–283.
- Maire, V., Alvarez, G., Colombet, J., Comby, A., Despinasse, R., Dubreucq, E., Joly, M., Lehours, A.-C., Perrier, V., Shahzad, T., Fontaine, S., 2013. An unknown oxidative metabolism substantially contributes to soil CO₂ emissions. *Biogeosciences* 10,

- 1155–1167.
- Meisner, A., Rousk, J., Bååth, E., 2015. Prolonged drought changes the bacterial growth response to rewetting. *Soil Biol. Biochem.* 88, 314–322.
- Nannipieri, P., Sequi, P., Fusi, P., 1996. Chapter 7–Humus and enzyme activity. In: Piccolo, A. (Ed.), *Humic Substances in Terrestrial Ecosystems*. Elsevier Science B.V., Amsterdam, pp. 293–328.
- Quiquampoix, H., Servagent-Noinville, S., Baron, M., 2002. *Enzymes in the Environment*. Marcel Dekker, pp. 285–306.
- Rajalingam, D., Loftis, C., Xu, J.J., Kumar, T.K.S., 2009. Trichloroacetic acid-induced protein precipitation involves the reversible association of a stable partially structured intermediate. *Protein Sci.* 18, 980–993. A Publication of the Protein Society.
- Ramnarine, R., Wagner-Riddle, C., Dunfield, K.E., Voroney, R.P., 2012. Contributions of carbonates to soil CO₂ emissions. *Can. J. Soil Sci.* 92, 599–607.
- Schindlbacher, A., Borken, W., Djukic, I., Brandstätter, C., Spötl, C., Wanek, W., 2015. Contribution of carbonate weathering to the CO₂ efflux from temperate forest soils. *Biogeochemistry* 124, 273–290.
- Shanhun, F.L., Almond, P.C., Clough, T.J., Smith, C.M.S., 2012. Abiotic processes dominate CO₂ fluxes in Antarctic soils. *Soil Biol. Biochem.* 53, 99–111.
- Shi, A., Marschner, P., 2014. Drying and rewetting frequency influences cumulative respiration and its distribution over time in two soils with contrasting management. *Soil Biol. Biochem.* 72, 172–179.
- Stursova, M., Sinsabaugh, R.L., 2008. Stabilization of oxidative enzymes in desert soil may limit organic matter accumulation. *Soil Biol. Biochem.* 40, 550–553.
- Sugihara, S., Funakawa, S., Kadono, A., Takata, Y., Sawada, K., Fujii, K., Kosaki, T., 2015. In situ short-term dynamics of CO₂ flux and microbial biomass after simulated rainfall in dry croplands in four tropical and continental ecosystems. *Soil Sci. Plant Nutr.* 61, 392–403.
- Tamir, G., Shenker, M., Heller, H., Bloom, P.R., Fine, P., Bar-Tal, A., 2013. Organic N mineralization and transformations in soils treated with animal waste in relation to carbonate dissolution and precipitation. *Geoderma* 209–210, 50–56.
- Tamir, G., Shenker, M., Heller, H., Bloom, P.R., Fine, P., Bar-Tal, A., 2011. Can soil carbonate dissolution lead to overestimation of soil respiration? *Soil Sci. Soc. Am. J.* 75, 1414.
- Thomas, A.D., Hoon, S.R., 2010. Carbon dioxide fluxes from biologically-crusted Kalahari Sands after simulated wetting. *J. Arid Environ.* 74, 131–139.
- Unger, S., Máguas, C., Pereira, J.S., David, T.S., Werner, C., 2010. The influence of precipitation pulses on soil respiration – assessing the “Birch effect” by stable carbon isotopes. *Soil Biol. Biochem.* 42, 1800–1810.
- Van Gestel, M., Ladd, J.N., Amato, M., 1992. Microbial biomass responses to seasonal change and imposed drying regimes at increasing depths of undisturbed topsoil profiles. *Soil Biol. Biochem.* 24, 103–111.
- Warren, C.R., 2014. Response of osmolytes in soil to drying and rewetting. *Soil Biol. Biochem.* 70, 22–32.
- Wu, J., Brookes, P.C., 2005. The proportional mineralisation of microbial biomass and organic matter caused by air-drying and rewetting of a grassland soil. *Soil Biol. Biochem.* 37, 507–515.
- Xiang, S.-R., Doyle, A., Holden, P.A., Schimel, J.P., 2008. Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils. *Soil Biol. Biochem.* 40, 2281–2289. Special Section: *Enzymes in the Environment III*.
- Xu, L., Baldocchi, D.D., Tang, J., 2004. How soil moisture, rain pulses, and growth alter the response of ecosystem respiration to temperature. *Glob. Biogeochem. Cycles* 18, GB4002.
- Yan, L., Chen, S., Xia, J., Luo, Y., 2014. Precipitation regime shift enhanced the rain pulse effect on soil respiration in a semi-arid steppe. *PLoS One* 9, e104217.

On the origin of carbon dioxide released from rewetted soils

Fraser, Fiona

2016-07-06

Attribution 4.0 International

Fraser FC, Corstanje R, Deeks LK, et al., (2016) On the origin of carbon dioxide released from rewetted soils. *Soil Biology and Biochemistry*, Volume 101, October 2016, pp. 1-5

<http://dx.doi.org/10.1016/j.soilbio.2016.06.032>

Downloaded from CERES Research Repository, Cranfield University