

## Concentration and composition of bioaerosol emissions from intensive farms: Pig and Poultry Livestock

T.L. Gladding<sup>a</sup>, C.A. Rolph<sup>a</sup>, C.L. Gwyther<sup>a</sup>, R. Kinnersley<sup>b</sup>, K. Walsh<sup>b</sup>, S. Tyrrel<sup>c</sup>

<sup>a</sup>The Open University, Walton Hall, Milton Keynes. MK7 6AA. [toni.gladding@open.ac.uk](mailto:toni.gladding@open.ac.uk)

<sup>b</sup>The Environment Agency, Horizon House, Deanery Road, Bristol, BS1 5AH

<sup>c</sup>School of Water, Energy and Environment, Cranfield University, Bedfordshire, MK43 0AL

### Abstract

Intensive farming is widespread throughout the UK and yet the health effects of bioaerosols which may be generated by these sites are currently not well researched. A scoping study was established to measure bioaerosols emitted from intensive pig ( $n = 3$ ) and poultry farms ( $n = 3$ ) during the period 2014-2015. The concentration of culturable mesophilic bacteria, Gram-negative bacteria, *Staphylococcus* spp., and fungi selecting for presumptive *Aspergillus fumigatus* were measured using single-stage impaction Andersen samplers, whilst endotoxin and (1→3)- $\beta$ -D-glucan was undertaken using inhalable personal samplers. Particulate matter concentration was determined using an optical particulate monitor. Results showed that culturable bacteria, fungi, presumptive *Staphylococcus aureus* (confirmed only as *Staphylococcus* spp.) and endotoxin concentrations were elevated above background concentrations for distances of up to 250 m downwind of the source. Of all the culturable bioaerosols measured, bacteria and *Staphylococcus* spp. were identified as the most significant, exceeding published or proposed bioaerosol guidelines in the UK. In particular, culturable *Staphylococcus* spp. downwind was at least 61 times higher than background at the boundary and at least 8 times higher 70m downwind on the four farms tested. This research represents a novel dataset of intensive farm emissions within the UK. Future research should exploit the use of innovative culture-independent methods such as next generation sequencing to develop deeper insights into the make-up of microbial communities emitted from intensive farming facilities and which would better inform species of interest from a public health perspective.

### Key Words

Bioaerosols; Intensive Farming; Agriculture; Endotoxin; Air Sampling.

## Introduction

The use of intensive animal farming is widespread throughout the UK. There are 217,000 agricultural holdings within the UK employing approximately 474,000 people (Department of Environment, Food and Rural Affairs (Defra), 2018). There are some five million pigs and 182 million birds, 118 million of which are 'table chicken' e.g. broilers (Defra, 2018), distributed across approximately 10,900 pig farms (AHDB 2018) and 2,500 poultry farms (British Poultry Council, 2018).

Agricultural production and concentrated animal feeding operations can emit a range of pollutants to the atmosphere including odour, greenhouse gases, ammonia, volatile organic compounds and particulate matter (Defra 2018, Douglas et al., 2018, Hayes et al., 2006; Cambra-López et al., 2010; Tilman et al., 2002). Whilst some studies have investigated the effect of particulate matter on the health of livestock workers (Andersen et al., 2004; Cambra-López et al., 2010), less is known about the exposure of the general population to bioaerosol emissions from such farming, either in the UK or elsewhere. In a systematic review to address these knowledge gaps Douglas et al., (2018) reported that a range of papers illustrated farm workers experienced respiratory and gastrointestinal problems and that negative health effects had been linked to surrounding populations and that further research was needed.

This study concentrates on bioaerosols, particulate matter from a biological origin, which are of concern as they can cause infection and an inflammatory response in the lungs, particularly in high risk groups such as the immunosuppressed or those with an existing respiratory disease (Biermann et al., 2013; Bünger et al., 2007; O'Gorman, 2011). Bioaerosols investigated here included microorganisms such as bacteria, fungi and their components endotoxin from the cell wall of Gram-negative bacteria and (1→3)-β-D-glucan from the cell wall of fungi (Gutarowska et al., 2015, Swan et al., 2003, Searl, 2008).

Bioaerosols have previously been associated with farming operations such as manure spreading, where it was concluded they could pose a risk to downwind receptors (Jahne et al., 2015). Elevated concentrations of bacteria, fungi, endotoxin and antibiotic resistant bacteria have also been reported downwind of intensive swine and poultry farms (Defra, 2009; Gibbs et al., 2006; Ko et al., 2008; Schulz et al., 2012). For example, *Staphylococcus aureus* has previously been identified as a potential indicator organism for emissions from chicken broiler houses (Schulz et al., 2011) and occupationally on pig farms (Masclaux et al., 2013). *S. aureus* is a potential human pathogen and antibiotic resistant *S. aureus* has also previously been identified in air samples up to 150 m from pig feeding operations (Gibbs et al., 2006). Whilst much of the existing evidence shows that intensive farming produces bioaerosols, the distances at which concentrations return to background are unclear. A comprehensive study

of poultry farms by Defra (2009) concluded that bioaerosols approached background values at 100 m downwind. However, other studies have identified drug and antibiotic resistant bacteria up to 150 m downwind of the source (Gibbs et al., 2006; Schulz et al., 2012).

It has also been identified that bioaerosol components, such as endotoxin, are present in high concentrations in air samples close to livestock farms (Schulze et al., 2006; Thorne et al., 2009; Ko et al., 2008) and that air pollutants from swine farms can cause acute respiratory symptoms in people living within 1.5 miles of swine operations (Schinasi et al., 2011). Schulze et al., (2006) reported endotoxin concentrations up to 23.2 EU/m<sup>3</sup> in gardens close to intensive farms compared to 0.7 EU/m<sup>3</sup> at a reference site. Endotoxin at concentrations associated with adverse health effects have since been found downwind of swine livestock operations (Thorne et al., 2009; Schinasi et al., 2011).

However, the health risk that livestock farms may pose to the general public remains uncertain. The Health Council of the Netherlands (2012) reported that “there is clear evidence that local residents can be exposed to microorganisms and substances derived from them”. Outbreaks of Q-fever, caused by *Coxiella burnetii*, in both the Netherlands and the UK have been attributed to nearby livestock farms (van der Hoek et al., 2010; Wallensten et al., 2010). Radon et al., (2007) found an increased prevalence of wheezing with an increasing number of animal houses in the surrounding area. In contrast, a systemic review concluded that there were insufficient dose response data to draw conclusions about the impact of animal feeding operations on community health (O’Connor et al., 2010). However, Smit et al., (2012) determined there was an increased pneumonia incidence associated with the presence of poultry within 1 km in adults, but a later paper (Smit et al., 2014) determined there was a statistically significant negative association with farm related PM<sub>10</sub> and all health outcomes.

For regulatory purposes an endotoxin exposure limit for the general public of 30 EU/m<sup>3</sup> has been proposed in the Netherlands (Health Council of the Netherlands, 2010), but the UK has no limit. However, environmental trigger levels have been in force in the UK since 2009 for environmental concentrations of non-speciated bacteria and *Aspergillus fumigatus* at 1000 and 500 CFU/m<sup>3</sup> downwind of waste composting (Drew et al., 2009). There is also the potential that glucan can cause airway inflammation when present above 10 ng/m<sup>3</sup> (Rylander, 1997). For environmental particulate matter, PM<sub>10</sub> has a European air quality target value of 50 µg/m<sup>3</sup> (as a 24 hour running mean) and PM<sub>2.5</sub> a target value of 25 µg/m<sup>3</sup> (Defra 2012).

This study was designed as a first investigation of bioaerosols downwind from intensive farming to evaluate emissions within a regulatory context. Hence bioaerosol sampling was carried out up and downwind of three poultry and three swine farms within the UK to determine particulate matter, endotoxin, bacterial and fungal concentrations in emissions. This

investigation was designed to measure bioaerosols using culture-dependent microbiology, and endotoxin and glucan, at all sites using the same methodology, to increase understanding of the scale and composition of viable bioaerosols associated with intensive livestock farming. The aim of this study was to improve the regulatory science evidence base needed to inform future decision-making and inform risk assessment regarding emissions from these facilities.

## Methods

### Study design

During 2014 and 2015 three chicken and three pig farms covered by the Environmental Permitting Regulations (2010) within the UK were selected for a pilot investigation as seen in Table 1. Sampling was undertaken on 12 separate days – each farm received two visits. Two farms were visited twice in spring 2014, and four farms twice in spring 2015.

**Table 1 Summary of Farm characteristics**

<b>Farm ID</b>	<b>Type</b>	<b>Farm animal Capacity</b>	<b>Ventilation</b>	<b>Activity</b>
1	Chicken layers	590,000	Automatic temperature control	Normal running and deliveries
2	Chicken broilers	205,000	Roof vents, 16 automatic fans per shed	Normal running chickens/mucking out
3	Chicken broilers	150,000	Roof fans	Normal running of chickens/mucking out
4	Pig	7,000	Mix of automatic fans and passive door and roof vents	Movement of pigs onsite
5	Pig	6,480	Side vented fans	Normal running of pigs
6	Pig	3,510	Passive door and roof vents, 1-2 fans on older sheds	Normal running of pigs

Sampling was carried out at chicken farms at the peak of the cycle e.g. all animals were present in the houses and when there was activity on site, pigs were cycled continuously and therefore both represented peak intensive production. Sampling locations were selected at least 5-10 m from the nearest buildings with no overhanging trees. At each site samples were taken upwind (10-50+ m), onsite (0 m), at the downwind boundary (10-31 m) and downwind (40-250 m) consisting of between four and five sampling points sequentially per farm per day. Downwind locations were determined by wind direction (verified by anemometer) and access considerations. All farms were surrounded by open flat farm land though this was often not in the ownership or control of the farms, hence compromises were made with distance. There were no other local sources of bioaerosols at any of the sites.

Each farm was monitored for particulates, viable (selected) microorganisms (following the AfOR Protocol (2009), endotoxin and glucan at each sample point. Viable *Staphylococcus* sp. was carried out on four of the farms (two chicken, two pig) as an indicator of animal origin. Information on replicates is given below.

### **Particulate sampling and enumeration**

An Osiris particulate monitor (Turnkey Instruments, Cheshire UK) was used to measure particulate concentrations. The Osiris uses a light-scattering technique to determine the concentration of airborne particles. The monitor was operated at 0.6 L/min for at least 30 minutes at each sample location concurrently with bioaerosol sampling (each data point consists of a 30 minute run from which is calculated an average). Total suspended particulates (TSP), particulate matter with a diameter less than or equal to 10 µm (PM<sub>10</sub>) and particulate matter with a diameter less than or equal to 2.5 µm (PM<sub>2.5</sub>) was enumerated by this method.

### **Weather Conditions**

Weather conditions such as cloud cover were noted manually. The Osiris was also used to log temperature, relative humidity, and wind speed and wind direction (from an inline anemometer placed at 1.8m) every minute of the sampling period.

### **Sampling and analysis of culturable microorganisms**

Sampling of culturable microorganisms was undertaken using the impaction method with four single-stage Andersen samplers with a hemi-cylindrical baffle in accordance with the Association for Organics Recycling Protocol (AfOR, 2009). Nutrient agar (NA), MacConkey agar No. 3 (MAC) and malt extract agar (MEA) plates were prepared as outlined in the AfOR Protocol (2009) to determine concentrations of culturable bacteria, Gram-negative bacteria and fungi selecting for *Aspergillus fumigatus*. Mannitol salt agar (MSA) was used for the collection of *Staphylococcus* spp. and prepared according to the manufacturer's instructions. All media were obtained from Oxoid Ltd. (Basingstoke, UK) and supplements from Fisher Scientific (Loughborough, UK) and Sigma Aldrich (Dorset, UK). Each petri dish contained 40 mL of media aseptically in advance of sampling and stored at 4°C, and were loaded on-site aseptically and connected to a vacuum pump. An inline-flow meter was used to calibrate the pump to 28.3 L/min. Plates were exposed for 2 minutes (NA, MAC and MSA) or 10 minutes (MEA). Four duplicate samples were taken at each location, with field blanks unloaded without

exposure and control plates were left in a sealed bag onsite. All samples were returned to the laboratory for incubation on the same day as sampling. Plates were incubated for 48 hours and checked again after 3-4 days at 37 °C for bacteria and 40 °C for fungi to favour thermotolerant fungi such as *A. fumigatus* as per the AfOR (2009) protocol.

After incubation, emerging colonies on the plates were counted and positive hole correlation, (Macher, 1989), was applied where colonies exceeded 20 on a plate. Results are expressed as colony forming units per cubic metre of air sampled (cfu/m<sup>3</sup>). The limit of detection (LOD) for the Andersens were 18 cfu/m<sup>3</sup> for bacteria and *Staphylococcus spp.* and 4 CFU/m<sup>3</sup> for fungi. Each data point consists of the average of the four samples (if samples were within 25% of each other). Field blanks were used to confirm that contamination of either the media or handling of the sampler were below 2 colonies per plate.

### **Sampling and analysis of endotoxin and (1→3)-β-D-glucan**

Samples for endotoxin and (1→3)-β-D-glucan were collected in triplicate using pre-sterilised Institute of Occupational Medicine (IOM) personal sampling heads operated as per the AfOR (2009) protocol. IOM cassettes were loaded with 25 mm 0.8 μm polycarbonate filters and operated with a pump at 2.0 L/min for 30 minutes. Samplers were deployed in duplicate and pre-calibrated at 2.0 L/min. Field blanks were loaded but not exposed, and two control cassettes were unopened. Cassettes were returned to the laboratory and stored overnight at 4°C. Material from the filters were then extracted into 5 mL of pyrogen free water (PFW) by shaking at 100 rpm for 30 minutes. Samples were stored at -20° prior to processing.

Endotoxin and (1→3)-β-D-glucan were quantified using a kinetic chromogenic *Limulus* amoebocyte lysate (LAL) assay (ACC Associates of Cape Cod) at 37°C with an automated microplate reader (BioTek ELx808, Swindon, UK) as per manufacturer instructions. Kinetic readings were recorded every 30 seconds for 90 minutes. For endotoxin, five concentrations of control standard endotoxin (CSE) were prepared within the range 50 – 0.005 Endotoxin Units (EU)/mL using serial dilutions. LAL (Pyrotell-T) was reconstituted with glucashield buffer (to prevent glucan interference). The LOD was 0.42 EU/m<sup>3</sup>. For (1→3)-β-D-glucan, six concentrations of glucan standard were prepared in the range 100 – 3.125 pg/mL using serial dilutions. GlucateLL lysate reconstituted with pyrogen free water and pyrosol buffer was added to each well. The LOD was 0.26 ng/m<sup>3</sup>. Each data point for endotoxin and glucan were the result of the average of three filters (where analysis was replicated in triplicate for each filter).

## Statistical analysis

Statistical analysis was performed in R 3.3.0 (R Core Team 2016). To determine whether there were any differences between farm type, ANOVA was performed on log transformed downwind data followed by Tukey's pairwise comparison. The non-parametric Kruskal Wallance test was used on untransformed data to test the significance of the sampling location. This was followed by the post-hoc Dunn test of multiple comparisons using rank sums. Mean impact range was calculated as described by CEN (2015). Where results were below the limit of detection (LOD), it was planned to use half the LOD value for statistical analysis, in practice this was unnecessary for all except viable fungi.

## Results

### Environmental Conditions

Meteorological measurements (day averages) are presented in Table 2.


Farm ID/visit	Type	Weather	Temp (°C)	Relative Humidity (%)	Wind speed (metres per second)
1/1	Chicken layers	Sun	10.6	66.7	0.6
1/2	Chicken layers	Sun	14.6	48.8	2.6
2/1	Chicken broilers	Sun	14.3	54.1	0.7
2/2	Chicken broilers	Sun	12.4	73.9	1.6
3/1	Chicken broilers	Cloudy	16.5	79.4	1.2
3/1	Chicken broilers	Sun	24.1	51.1	1.3
4/1	Pig	Sun	14.8	57.3	5.5
4/2	Pig	Cloudy	14.6	78.9	2.3
5/1	Pig	Sun	24.2	58.0	0.7
5/2	Pig	Cloudy	18.9	91.1	1.6
6/1	Pig	Sun	13.5	49.3	4.9
6/2	Pig	Cloudy	13.9	79.7	3.3

Meteorological measurements showed some variation between sampling visits. Sampling was only undertaken during dry weather. Temperatures ranged from 10.6 °C to 24.2 °C. Average daily wind speeds ranged from 0.6 – 5.5 m/s.

## Bioaerosols


Bioaerosols were detected in all sites as seen in Figure 1. Of the culturable bioaerosols bacteria and *Staphylococcus* spp. were identified as the most significant, exceeding published or proposed bioaerosol guidelines (1000 cfu/m<sup>3</sup> proposed by Drew et al., 2009) and having higher concentrations downwind compared to upwind. *Staphylococcus* spp. were found in significant culturable concentrations (up to 1.4 x 10<sup>4</sup> cfu/m<sup>3</sup>) at farm boundaries and at 70 m downwind (up to 1.9 x 10<sup>3</sup> cfu/m<sup>3</sup>). General fungi, presumptive Gram-negative bacteria and *A. fumigatus* were all detected across all sites but at low concentrations and mostly with no significant increase in concentration in downwind compared to upwind samples (Figure 1). Broadly the source impact range was in most cases between -50m (upwind) and 50-100m (downwind). The incidence of occasional high concentrations upwind of source may suggest that upwind sampling points were too close to source and subject to a source influence, as other sources were not seen within 250m.





**Figure 1** Culturable bioaerosol concentrations from different types of farm. Dashed lines indicate mean upwind concentration plus 2 standard deviations. Note *Staphylococcus spp.* are results from four farms, the remaining graphs are results from all six farms.

The bioaerosol components and particulates tested are displayed in Figure 2 (average of replicates). Endotoxin concentrations exceeded proposed guidelines (30 EU/m<sup>3</sup> (Health Council of the Netherlands, 2010) in both upwind and downwind samples up to 62 m from the source. Values for total suspended particulates (TSP) were below 250 µg/m<sup>3</sup> with the exception of one sampling location at a layer farm where values averaged 379 µg/m<sup>3</sup> at source. There was no significant difference between farm type or between sampling location for any of the particulate measurements taken ( $p > 0.05$ ).



**Figure 2 Bioaerosol components and particulate concentrations from different types of farm. Dashed lines indicate mean upwind concentration plus 2 standard deviations**

Bioaerosols showed a clear reduction in concentration with location. Generally upwind samples had relatively low concentrations of bioaerosol and were significantly different from

source and downwind samples for bacteria and *Staphylococcus* spp. ( $P < 0.05$ ) illustrated in Figure 3. However there was no significant difference in endotoxin across the different locations ( $P > 0.05$ ). The source type had a small impact on the amount of bioaerosol emitted. Significantly less endotoxin was measured from the broilers (8.61 EU/m<sup>3</sup>) compared to layers (52.18 EU/m<sup>3</sup>) but there was no difference for swine compared to broilers or *Staphylococcus* spp. (Table 3). Whilst broiler emissions resulted in apparent higher concentrations of bacteria, there was no significant difference when compared to the other farm types.

**Table 3 Mean bioaerosols concentration by location and source type. Values in parenthesis indicate concentration range and letters indicate groups of statistical similarity.**

	Sampling Location			Source Type (Downwind data)		
	Upwind	Source	Downwind*	Layers	Broilers	Swine
Bacteria (CFU/m <sup>3</sup> )	479(141 – 1608)	3,674 (141 – 10,870) <sup>a</sup>	2631 (124 – 15,600) <sup>a</sup>	2,421 (177 – 10,870) <sup>d</sup>	5,795 (265 – 15,601) <sup>de</sup>	1,246 (124 – 4,329) <sup>e</sup>
<i>Staphylococcus</i> Spp. (CFU/m <sup>3</sup> )	75 (18 – 230)	3995 (177 – 10,390) <sup>b</sup>	2471 (35 – 14,700) <sup>b</sup>	1,608 (71 – 6,343) <sup>f</sup>	7,374 (35 – 14,700) <sup>f</sup>	1,468 (177 – 3,693) <sup>f</sup>
Endotoxin (EU/m <sup>3</sup> )	14.89 (0.60 – 82.71) <sup>c</sup>	33.03 (0.220 – 158.70) <sup>c</sup>	19.71 (0.22 – 82.97) <sup>c</sup>	52.18 (3.03 – 158.70) <sup>g</sup>	8.61 (0.22 – 32.84) <sup>h</sup>	25.58 (0.45 – 82.70) <sup>gh</sup>

\* From on-site to furthest sampling point pooled

All field blanks and controls were clear of contamination and recorded concentrations below the limit of detection.

## Discussion

This is the first data collection of bioaerosols from intensive pig and poultry sites utilising a standardised sampling approach (AfOR Protocol 2009), which has simultaneously sampled for and enumerated culturable microorganisms, endotoxin, and (1→3)-β-D-glucan within the UK. This is a unique dataset regarding emissions from intensive farming, and in a recent systematic review it was reported that studies of exposure downwind from intensive farming were much rarer than occupational in a ratio of 1:21 (Douglas et al., 2018).

This research measured the distance bioaerosols travelled from source before returning to background. Previous work in the UK (Defra 2009) examined culturable bioaerosols, endotoxin and particulates at poultry houses and determined that these bioaerosols were reduced to background within 100 m from source. However, Gibbs et al., (2006) found that concentrations were significantly higher than upwind up to 150 m from source. Our study

265 found that particulate concentrations were elevated close to source at both chicken and pig  
266 farms but returned to background concentrations within 150 m from source (Figure 2).  
However, some concentrations of bacteria and *Staphylococcus* spp. remained elevated above  
background concentrations potentially to 250m. Smit et al., (2014) modelled health outcomes  
based on PM<sub>10</sub>, but in this study it was not clear whether particulate matter was a good  
predictor of bioaerosol concentrations downwind. Douglas et al., (2018) report that farms are  
required to carry out a bioaerosol risk assessment if sensitive receptors (such as people living  
or working nearby) are within 100m, which contrasts to the requirement at waste composting  
sites which is 250m. This study indicates that distances travelled by organisms such as  
*Staphylococcus* spp. were found to be up to 250m, hence farms should be required to carry  
out an assessment to match that currently required of waste composting, though perhaps  
targeted at bacteria, which is discussed further below.

In terms of the composition of the bioaerosols, although there was significant between-site (of  
different farm types) and within-site (of the same farm types) variability, there was evidence  
that culturable bacteria concentrations can be significantly elevated at both chicken and pig  
farms. However, culturable fungi, particularly *A. fumigatus*, are much lower and are therefore  
not likely to pose a significant concern or be a useful indicator organism in these  
circumstances. Ko, et al., (2008) demonstrated bacteria had a stronger downwind signature  
in relation to upwind concentrations compared to fungi at farm sites and Defra (2009) found  
that the emission profile of bacteria was consistently higher than for fungi. This finding  
contrasts with the findings of studies at biowaste sites where fungi are consistently detected  
in higher concentrations (Swan, et al., 2003, Searl, 2008, Gutarowska, et al., 2015) and  
indicates a different emission spectrum at intensive farms compared to organic waste  
processing facilities.

*Staphylococcus* species were elevated at source on site on all four farms studied, in particular  
on broiler and layer chicken sites with concentrations exceeding  $1.4 \times 10^4$  cfu/m<sup>3</sup> in individual  
replicates. The highest upwind/background samples on farms showed *Staphylococcus* spp.  
at concentrations of 230 cfu/m<sup>3</sup> but more often below 100 cfu/m<sup>3</sup>, i.e. downwind were 61 times  
higher than background at the boundary and 8 times higher 70m downwind on the four farms  
tested. It is also possible the upwind samples were on occasion contaminated by the farm  
site due to restrictions in upwind sampling distance, hence the differences may in fact have  
been more marked compared to background. The vast majority of colonies were presumptive  
*Staphylococcus aureus* (as identified by colony colour on MSA) on selective agar. Schulz et  
al., (2012) also found significant differences in the amount of *Staphylococcus* species upwind  
compared to downwind (as MRSA) from farms.

Previous work, and this research, shows that culturable *Staphylococcus* species appear to be a reliable indicator of animal house emissions at both chicken and pig farms. Since there is not an equivalent *Staphylococcus* species dataset for biowaste facilities in England, as they are not considered indicator organisms in the AfOR (2009) protocol, it is not known if *Staphylococcus* species are also indicative of emissions from such sites. It is a limitation of this study that confirmation was not carried out on the presumptive culturable *Staphylococcus* spp. found, and future studies require further microbiological analysis to confirm *Staphylococcus* species.

In addition to the health concerns a pathogen such as *Staphylococcus aureus* poses, it also has the potential to be a source of antibiotic resistance which could contribute to the presence of such organisms within the environment around farms. Previous work identified that *Staphylococcus* species isolated both on pig farms (Masclaux et al., 2013) and downwind of farms (Schulz, et al., 2012) were antibiotic resistant strains such as MRSA. Future work on UK farms should consider this possibility.

For other organisms, culturable Gram-negative bacteria were around the limit of detection around all sites. Generally it appears that culturable Gram-negatives remain low in other settings such as waste management (Gladding and Gwyther 2017) and is not included as a parameter in the replacement to the AfOR Protocol (2009), known as M9 (Environment Agency 2017), within the UK. However, concerns have grown over recent years because of the emergence of antibiotic resistant Enterobacteriaceae among livestock (Seiffert et al., 2013), and further work should perhaps be targeted on these as additional indicators.

Endotoxin appears to be elevated downwind in this study at concentrations exceeding the suggested environmental limit of 30 EU/m<sup>3</sup> at distances of up to 62m, even though culturable Gram-negative bacteria were low. Schulz, et al., (2006) found elevated concentrations of endotoxin in the vicinity of intensive farming, though with some spatial variability. Defra (2009) also reported a significant emission profile of endotoxin from poultry farms as did Thorne, et al., (2009) from swine farming. In contrast, the concentrations of (1→3)-β-D-glucan found in this study are low, which corresponds to the lower signature of culturable fungi. There is no comparable dataset on (1→3)-β-D-glucan emissions from intensive farming in the literature.

Further work is needed regarding dispersion from farms incorporating boundary and distance sampling. The downwind distances sampled in this study were limited by practicalities of access at the sites studied. Future studies should aim to capture distances further downwind, simultaneously to upwind, to assess the extent of the dispersion decay curves and the point at which they approach background concentrations, although it is acknowledged access is often difficult. Finally, this study was limited to targeted viable

microorganisms, but initial scoping work on the non-viable component (not reported here due to limitations in methodological design) indicate that the diversity of species being emitted from intensive pig and poultry farms warrants further investigation.

## Conclusions

Intensive pig and poultry farming has a measurable impact on the bioaerosols concentration downwind of facilities. Viable cultivation revealed that bacteria, culturable fungi, and *Staphylococcus spp.*, and endotoxin, were elevated above background levels at source and for varying distances of up to 250 m downwind. In conclusion, this study found that from a regulatory perspective, the suite of indicator organisms used to monitor biowaste sites is not directly transferable to intensive farms. There is also evidence to suggest that bioaerosol concentrations remain elevated for a greater distance downwind of animal houses than was found in a previous UK study but commensurate with that found at biowaste facilities, indicating that the requirement to carry out a risk assessment downwind should be harmonised between farms and waste sites. Further monitoring work should be targeted at particle size, and distribution and dispersion. Future research should also exploit more innovative culture-independent technology such as next generation sequencing to provide a more comprehensive understanding of the species of interest from a public health perspective to inform new targets for cultivation.

## References

- AfOR (2009) A standardised protocol for the monitoring of bioaerosols at open composting facilities. Northamptonshire. Edited T Gladding. Available at: <http://www.organics-recycling.org.uk/page.php?article=1750&name=Standardised+Protocol+for+Monitoring+Bioaerosols> (Accessed: 5/12/16).
- AHDB (2018) Pig holdings in the UK. Agricultural and Horticultural Development Board. <http://pork.ahdb.org.uk/prices-stats/industry-structure/pig-holdings-in-the-uk/> (accessed 30/10/18)
- Andersen C. I., Von Essen S. G., Smith L. M., Spencer J., Jolie R., Donham, K. J. (2004) Respiratory symptoms and airway obstruction in swine veterinarians: A persistent problem. *American Journal of Industrial Medicine*, 46(4), pp. 386–392
- Biermann J., Merk H. F., Wehrmann W., Klimek L., Wasem J. (2013) Allergic disorders of the respiratory tract - Findings from a large patient sample in the German statutory health insurance system. *Allergo Journal*, 22(6), pp. 366–373

- Bünger J., Schappler-Scheele B., Hilgers R., Hallier, E. (2007) A 5-year follow-up study on respiratory disorders and lung function in workers exposed to organic dust from composting plants. *Int Arch Occup Environ Health*, 80, 306-312
- British Poultry Council (2018) About the British Poultry Council. <https://www.britishpoultry.org.uk/about-bpc/> (accessed 30/10/18)
- Cambra-López M., Aarnink A. J. a, Zhao Y., Calvet S., Torres, A. G. (2010) Airborne particulate matter from livestock production systems: A review of an air pollution problem. *Environmental Pollution*, 158(1), pp. 1–17
- Defra (2018) Agriculture in the United Kingdom 2017 Crown Copyright 2018 <https://www.gov.uk/government/statistics/agriculture-in-the-united-kingdom-2017> (accessed 25/06/19)
- Defra (2012) UK and EU Air Quality Limits (National Air Quality Objectives). [https://uk-air.defra.gov.uk/assets/documents/National\\_air\\_quality\\_objectives.pdf](https://uk-air.defra.gov.uk/assets/documents/National_air_quality_objectives.pdf) (accessed 25/6/16)
- Defra (2009) Characterising poultry dust properties, assessing the human health implications, quantifying emission levels and assessing the potential for abatement, London. <http://sciencesearch.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=14432> (accessed 30/10/18)
- Douglas P., Robertson S., Gay R., Hansell A.L., Gant T.W. (2018) A systematic review of the public health risks from intensive farming. *Int Jnl Hygiene and Env Health* 221:134-173
- Drew G., Deacon L., Pankhurst L., Pollard S. J., Tyrrel S. (2004) Guidance on the evaluation of bioaerosol risk assessments for composting facilities. Published by The Environment Agency.
- Environmental Permitting Regulations (2010) The Environmental Permitting (England and Wales) Regulations 2010. <http://www.legislation.gov.uk/ukdsi/2010/9780111491423/contents> (accessed 30/10/18)
- Environment Agency (2017) M9: Environmental monitoring of bioaerosols at regulated facilities. <https://www.gov.uk/government/publications/m9-environmental-monitoring-of-bioaerosols-at-regulated-facilities> (accessed 30/10/18)
- Gladding T.L., Gwyther C.L. (2017) A Study of the Potential Release of Bioaerosols from Containers as a Result of Reduced Frequency Residual Waste Collection. *Science of the Total Environment*. 576: 481-489
- Gibbs S. G., Green C. F., Tarwater P. M., Mota L. C., Mena K. D., Scarpino P. V. (2006) Isolation of antibiotic-resistant bacteria from the air plume downwind of a swine confined or concentrated animal feeding operation. *Environmental Health Perspectives*, 114(7), pp. 1032–1037

- 402 Gutarowska B., Skora J., Stepień L., Szponar B., Otlewska A., Pielech-Przybylska K. (2015)  
 403 Assessment of microbial contamination within working environments of different types of  
 404 composting plants. *Journal of the Air & Waste Management Association*, 65(4), pp. 466-478
- 405 Hayes E. T., Curran T. P., Dodd, V. A. (2006) Odour and ammonia emissions from intensive pig units  
 406 in Ireland. *Bioresource Technology*, 97(7), pp. 940–948
- 407 Health Council of the Netherlands (2010) Endotoxins. Health-based recommended occupational  
 408 exposure limit, Publication no. 2010/04OSH.
- Health Council of the Netherlands (2012) Health risks associated with livestock farms. 2012/27E., The Hague.
- van der Hoek W., Dijkstra F., Schimmer B., Schneeberger P. M., Vellema P., Wijkmans C., ter Schegget R., Hackert V., van Duynhoven, Y. (2010) Q fever in the Netherlands: an update on the epidemiology and control measures. *Eurosurveillance*, 15(12), pp. 2007–2011
- Jahne M. A., Rogers S. W., Holsen T. M., Grimberg S. J., Ramler, I. P. (2015) Emission and Dispersion of Bioaerosols from Dairy Manure Application Sites: Human Health Risk Assessment. *Environmental Science & Technology*, p. 150730083724002.
- Ko G., Iii O. D. S., Likirdopulos C., Worley-davis, L., Williams, M., Sobsey, M. D. (2008) Investigation of Bioaerosols Released from Swine Farms using Conventional and Alternative Waste Treatment and Management Technologies. *Environmental Science & Technology*, 42(23), pp. 8849–8857
- Macher J. M. (1989) Positive-hole correction of multiple-jet impactors for collecting viable microorganisms. *American Industrial Hygiene Association journal*, 50(11), pp. 561–568
- Masclaux F.G., Sakwinska O., Charrière N., Semaani E., Oppliger A., (2013) Concentration of airborne *Staphylococcus aureus* (MRSA and MSSA), total bacteria, and endotoxins in pig farms. *Ann Occup Hyg.* 57(5):550-7
- O'Connor A. M., Auvermann B., Bickett-Weddle D., Kirkhorn S., Sargeant J. M., Ramirez A., Von Essen S. G. (2010) The association between proximity to animal feeding operations and community health: A systematic review. *PLoS ONE*, 5(3)
- O'Gorman C. M. (2011). Airborne *Aspergillus fumigatus* conidia: a risk factor for aspergillosis. *Fungal Biology Reviews*, 25(3), pp. 151-157
- Radon K., Schulze A., Ehrenstein V., van Strien R. T., Praml G., Nowak, D. (2007) Environmental exposure to confined animal feeding operations and respiratory health of neighboring residents. *Epidemiology (Cambridge, Mass.)*, 18(3), pp. 300–308
- R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.



- Rylander R. (1997) Airborne (1-3)-beta-D-glucan and airway disease in a day-care center before and after renovation. *Archives of microbiology*, 52(4), pp. 281–285
- Schinasi L., Horton R. A., Guidry V. T., Wing S., Marshall S. W., Morland K. B. (2011) Air pollution, lung function, and physical symptoms in communities near concentrated Swine feeding operations. *Epidemiology (Cambridge, Mass.)*, 22(2), pp. 208–215.
- Schulz J., Formosa L., Seedorf J., Hartung, J. (2011) Measurement of culturable airborne staphylococci downwind from a naturally ventilated broiler house. *Aerobiologia*, 27(4), pp. 311–318
- Schulz J., Friese A., Klees S., Tenhagen B., Fetsch A., Rösler U., Hartung, J. (2012) Longitudinal study of the contamination of air and of soil surfaces in the vicinity of pig barns by livestock-associated methicillin-resistant *Staphylococcus aureus*. *Applied and Environmental Microbiology*, 78(16), pp. 5666–5671
- Schulze A., Van Strien R., Ehrenstein V., Schierl R., Küchenhoff H., Radon K. (2006) Ambient endotoxin level in an area with intensive livestock production. *Annals of Agricultural and Environmental Medicine*, 13(1), pp. 87–91
- Searl A. (2008). Exposure-response relationships for bioaerosol emissions from waste treatment processes, Institute of Occupational Medicine, Edinburgh, UK. <http://sciencesearch.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=15140> (Accessed: 30/10/18)
- Seiffert S. N, Hilty M, Perreten V., Endimiani A (2013) Extended-spectrum cephalosporin-resistant Gram-negative organisms in livestock: an emerging problem for human health? *Drug Resistance Updates* Feb-April; 16 (1-2), 22-45
- Smit, L.A., van der Sman-de Beer, F., Opstal-van Winden, A.W., Hooiveld, M., Beekhuizen, J., Wouters, I.M., Yzermans J., Heederik D., 2012. Q fever and pneumonia in an area with a high livestock density: a large population-based study. *PLoS One* 7, e38843.
- Smit, L.A., Hooiveld, M., van der Sman-de Beer, F., Opstal-van Winden, A.W., Beekhuizen, J., Wouters, I.M., (2014) Air pollution from livestock farms, and asthma, allergic rhinitis and COPD among neighbouring residents. *Occup. Environ. Med.* 71, 134–140
- Swan J. R. M., Kelsey A., Crook B. and Gilbert E. J. (2003). Occupational and environmental exposure to bioaerosols from composts and potential health effect - a critical review of published data. Sudbury, UK. <http://www.hse.gov.uk/research/rrpdf/rr130.pdf> (accessed 30/10/18)
- Thorne P. S., Ansley A. C., Perry S. S. (2009) Concentrations of bioaerosols, odors, and hydrogen sulfide inside and downwind from two types of swine livestock operations. *Journal of occupational and environmental hygiene*, 6(4), pp. 211–220
- Tilman D., Cassman K. G., Matson P., Naylor R., Polasky, S. (2002) Agricultural sustainability and intensive production practices. *Nature*, 418(6898), pp. 671–677

Wallensten A., Moore P., Webster H., Johnson C., van der Burgt G., Pritchard G., Ellis-Iversen J., Oliver I. (2010) Q fever outbreak in Cheltenham, United Kingdom, in 2007 and the use of dispersion modelling to investigate the possibility of airborne spread. *Eurosurveillance*, 15(12)

## **Acknowledgements**

The views expressed in this paper are those of the authors, and not necessarily those of the Environment Agency.

Funding: This work was funded by the Environment Agency for England and Wales grant numbers SC130025/1 and SC130025/2.

The authors declare no competing interests.

# Concentration and composition of bioaerosol emissions from intensive farms: pig and poultry livestock

Gladding, T. L.

2020-07-14

Attribution-NonCommercial-NoDerivatives 4.0 International

---

Gladding TL, Rolph CA, Gwyther CL, et al., (2020) Concentration and composition of bioaerosol emissions from intensive farms: pig and poultry livestock. *Journal of Environmental Management*, Volume 272, October 2020, Article number 111052.

<https://doi.org/10.1016/j.jenvman.2020.111052>

*Downloaded from CERES Research Repository, Cranfield University*